Proceedings of the Ninth Prague Topological Symposium
Contributed papers from the symposium held in
Prague, Czech Republic, August 19–25, 2001

ON THE METRIZABILITY OF SPACES WITH A SHARP BASE

CHRIS GOOD, ROBIN W. KNIGHT, AND ABDUL M. MOHAMAD

Abstract. A base $\mathcal{B}$ for a space X is said to be *sharp* if, whenever $x \in X$ and $(B_n)_{n\in\omega}$ is a sequence of pairwise distinct elements of $\mathcal{B}$ each containing x, the collection $\{\bigcap_{j\le n} B_j : n \in \omega\}$ is a local base at x. We answer questions raised by Alleche *et al.* and Arhangel'skiĭ *et al.* by showing that a pseudocompact Tychonoff space with a sharp base need not be metrizable and that the product of a space with a sharp base and $[0, 1]$ need not have a sharp base. We prove various metrization theorems and provide a characterization along the lines of Ponomarev's for point countable bases.

The notion of a uniform base was introduced by Alexandroff who proved that a space (by which we mean T_1 topological space) is metrizable if and only if it has a uniform base and is collectionwise normal [1]. This result follows from Bing's metrization theorem since a space has a uniform base if and only if it is metacompact and developable. Recently Alleche, Arhangel'skiĭ and Calbrix [2] introduced the notions of sharp base and weak development, which fit very naturally into the hierarchy of such strong base conditions including weakly uniform bases (introduced by Heath and Lindgren [11]) and point countable bases (see Figure 1 below). In this paper we look at the question of when a space, with a sharp base is metrizable. In particular, we show that a pseudocompact space with a sharp base need not be metrizable, but generalize various situations where a space with a sharp base is seen to be metrizable.

Definition 1. Let $\mathcal{B}$ be a base for a space X.

2000 *Mathematics Subject Classification.* 54E20, 54E30.
Key words and phrases. Tychonoff space, pseudocompact, special bases, sharp base, metrizability.

Chris Good, Robin W. Knight and Abdul M. Mohamad, *On the metrizability of spaces with a sharp base*, Proceedings of the Ninth Prague Topological Symposium, (Prague, 2001), pp. 125–134, Topology Atlas, Toronto, 2002; `arXiv:math.GN/0204127`.

(1) $\mathcal{B}$ is said to be *sharp* if, whenever $x \in X$ and $(B_n)_{n\in\omega}$ is a sequence of pairwise distinct elements of $\mathcal{B}$ each containing x, the collection $\{\bigcap_{j\le n} B_j : n \in \omega\}$ is a local base at x.
(2) $\mathcal{B}$ is said to be *uniform* if, whenever $x \in X$ and $(B_n)_{n\in\omega}$ is a sequence of pairwise distinct elements of $\mathcal{B}$ each containing x, then $(B_n)_{n\in\omega}$ is a local base at x.
(3) $\mathcal{B}$ is said to be *weakly uniform* if, whenever $\mathcal{B}'$ is an infinite subset of $\mathcal{B}$, then $\bigcap \mathcal{B}'$ contains at most one point.
(4) $\mathcal{B}$ is said to be a *weak development* if $\mathcal{B} = \bigcup_{n\in\omega} \mathcal{B}_n$, each $\mathcal{B}_n$ is a cover of X and, whenever $x \in B_n \in \mathcal{B}_n$for each $n \in \omega$, then $\{\bigcap_{j\le n} B_j : n \in \omega\}$ is a local base at x.

Arhangel'skiĭ *et al.* prove that a space with a sharp base has a point countable sharp base ([2] and [4]) and is metaLindelöf. Moreover a weakly developable space has a G_δ-diagonal and a submetacompact space with a base of countable order is developable [2].

We note in passing that the obvious definition of 'uniform weak developability' (having a base $\mathcal{G} = \bigcup\{\mathcal{G}_n : n \in \omega\}$ such that each G_n is a cover and whenever $x \in G_n \in \mathcal{G}_n$, $\{G_n\}_n$ is a base at x) is simply a restatement of developability. We also note that a space with a σ-disjoint base need not have a sharp base: Bennett and Lutzer [7] construct a first countable (and a Lindelöf) example of a non-metrizable LOTS with σ-disjoint bases (and continuous separating families), which can not have a sharp base by Theorem 2.



FIGURE 1

When is a space with a sharp base metrizable? We summarize the relevant results of [2], [4] and [6] in the following theorem.

Theorem 2. *Let X be a regular space with a sharp base, then X is metrizable if any of the following hold:*
(1) *X is separable;*

(2) *X is locally compact (so a manifold with sharp base is metrizable);*
(3) *X is countably compact;*
(4) *X is pseudocompact and CCC;*
(5) *X is a GO space.*

A space is pseudocompact if every continuous real valued function is bounded. Every (Tychonoff) pseudocompact space with a uniform base is metrizable (see [19], [16] or [18]), whilst a pseudocompact space with a point-countable base need not be metrizable [17]. Moreover pseudocompact Tychonoff spaces with regular G_δ-diagonals are metrizable [14], whilst Mrowka's Ψ space is an example of a pseudocompact, non-metrizable Moore space. So it is natural to ask (see [2] and [4]) whether every pseudocompact space with a sharp base is metrizable. The space P of Example 3 shows that the answer to this question is 'no.' In addition, P answers a number of other questions in the negative: Alleche *et al.* ask whether the product $X \times [0,1]$ has a sharp base if X does; Heath and Lindgren [11] ask whether a space with a weakly uniform base has a G^*_δ-diagonal; and P is another example (see [17] and [20]) of a pseudocompact space with a point countable base that is not compact, and is a non-compact pseudocompact space with a weakly uniform base, answering questions of Peregudov [15].

Example 3. *There exists a Tychonoff, non-metrizable pseudocompact space with a sharp base but without a G^*_δ-diagonal whose product with the closed unit interval does not have a sharp base.*

Proof. Our example is a modification of the example of a non-developable space with a sharp base [2]. We add extra points to a (non-separable) metric space B in such a way that the resulting space is pseudocompact, has a sharp base but is not compact, hence not metrizable.

Let $B = {}^{\omega}\mathfrak{c}$ be the Tychonoff product of countably many copies of the discrete space of size continuum with the usual Baire metric. For each finite partial function $f \in {}^{<\omega}\mathfrak{c}$, let $[f]$ denote the basic open subset of B, $[f] = \{g \in {}^{\omega}\mathfrak{c} : f \subseteq g\}$ (so $[f]$ is the collection of all elements of B which agree with f on $\operatorname{dom} f$). Note that, if $\operatorname{dom} f \subseteq \operatorname{dom} g$, then the two basic open sets $[f]$ and $[g]$ have non-empty intersection if and only if $f \subseteq g$ if and only if $[g] \subseteq [f]$. If $[f] \cap [g] = \varnothing$ then the functions f and g are incompatible (we write $f \perp g$) and neither $f \subseteq g$ nor $g \subseteq f$.

Let
$$\mathcal{S} = \{S \in {}^{\omega}({}^{<\omega}\mathfrak{c}) : S(m) \perp S(n), \text{ for each } m \text{ and } n\},$$
so that each S in $\mathcal{S}$ codes for a sequence of disjoint basic open sets in B. Enumerate $\mathcal{S}$ as $\{S_\alpha : \alpha \in \mathfrak{c}\}$ in such a way that each S in $\mathcal{S}$ occurs $\mathfrak{c}$ times. To ensure that our space is pseudocompact, we recursively add limit points (to some of) these sequences of open sets. These limit points s_α will have basic open neighbourhoods of the form
$$N(\alpha, n) = \{s_\alpha\} \cup \bigcup_{m \geq n} [T_\alpha(m)],$$

where $T_\alpha \in {}^\omega({}^{<\omega}\mathfrak{c})$ is defined depending on S_α.

Suppose that for each $\alpha < \gamma$ we have either defined if possible a sequence $T_\alpha \in {}^\omega({}^{<\omega}\mathfrak{c})$ such that

(1γ) for $i \neq j$, $T_\alpha(i) \perp T_\alpha(j)$,
(2γ) for $\beta < \gamma$, $\beta \neq \alpha$, T_β defined, $\operatorname{ran} T_\alpha \cap \operatorname{ran} T_\beta = \varnothing$, and
(3γ) for $\beta < \gamma$, $\beta \neq \alpha$, T_β defined, if $T_\alpha(i) \supseteq T_\beta(j)$, then $T_\alpha(i') \perp T_\beta(j')$ for all $\langle i', j'\rangle \neq \langle i, j\rangle$

or we have not defined T_α. We now define T_γ.

First note that if $S'_\gamma(i)$ extends $S_\gamma(i)$, then the open set $[S'_\gamma(i)]$ is a subset of $[S_\gamma(i)]$, so any limit of the sequence of open sets $\{[S'_\gamma(i)] : i \in \omega\}$ will also be a limit of the sequence $\{[S_\gamma(i)] : i \in \omega\}$.

Since each $T_\alpha(j)$ is finite, there is some $\delta < \mathfrak{c}$ which is not in $\bigcup\{T_\alpha(j) : \alpha < \gamma, j \in \omega\}$. For each $i \in \omega$, let $S'_\gamma(i) = S_\gamma(i)^\frown\{\delta\}$ extend $S_\gamma(i)$. Then for all $i, j \in \omega$ and $\alpha < \gamma$, $S'_\gamma(i) \nsubseteq T_\alpha(j)$ and $T_\alpha(j) \subseteq S'(i)$ only if $T_\alpha(j) \subseteq S(i)$. Notice that this implies that $[T_\alpha(j)] \nsubseteq [S'_\gamma(i)]$ and that $[S'_\gamma(i)] \subseteq [T_\alpha(j)]$ only if $[S_\gamma(i)] \subseteq [T_\alpha(j)]$.

Case 1: Suppose that there exists some $\alpha < \gamma$ for which T_α was defined, such that for infinitely many $i \in \omega$ there exists some $j \in \omega$ such that $S'_\gamma(i) \supseteq S_\gamma(i) \supseteq T_\alpha(j)$. In this case we do not define T_γ (since infinitely many of the basic open sets $[T_\alpha(j)]$ contain an open set $[S_\gamma(i)]$ and the limit point s_α will deal with the sequence S_γ).

Case 2: Now suppose that Case 1 does not hold and that hence

(*) for each $\alpha < \gamma$ there are at most finitely many i for which $S'_\gamma(i) \supseteq T_\alpha(j)$ for some j.

Suppose further that for each $i \leq k$, we have chosen natural numbers $0 = r_0 < r_1 < \cdots < r_k$ and defined $T_\gamma(i)$ to be $S'_\gamma(r_i)$.

Since each $T_\gamma(i)$ is a finite partial function, there are at most finitely many possible partial functions such that $f \subseteq T_\gamma(i)$ for some $i \leq k$. By condition (2γ) there are at most finitely many $\alpha < \gamma$ with such an f in $\operatorname{ran} T_\alpha$. List these α as $\alpha(1), \ldots \alpha(m)$. By (*), for each $\alpha(m)$, there is a j_m such that for all $i \geq j$, $S'_\gamma(i)$ does not extend any $T_{\alpha(m)}(j)$. Now let $r_{k+1} = \max j_m$ and $T_\gamma(k+1) = S'_\gamma(r_{k+1})$.

We now claim that conditions (1$\mathfrak{c}$), (2$\mathfrak{c}$) and (3$\mathfrak{c}$) hold. Suppose that T_β and T_α were defined for some $\beta < \alpha < \mathfrak{c}$. Condition (1$\mathfrak{c}$) is obvious since each T_α is a subsequence of S'_α each term of which extends the corresponding term of S_α, and S_α is a sequence of pairwise incompatible partial functions. (2$\mathfrak{c}$) holds since, if $\beta < \alpha$, then the extension $S'_\gamma(i)$ was chosen to ensure that $T_\beta(j) \not\supseteq S'_\alpha(i)$ for any j, so in particular $T_\beta(j) \neq T_\alpha(i)$ and $\operatorname{ran} T_\beta \cap \operatorname{ran} T_\alpha$. To see that (3$\mathfrak{c}$) holds, note first that $S'_\alpha(i)$ was chosen so that $S'_\alpha(i) \nsubseteq T_\beta(j)$ for any j, which implies that $T_\alpha(i) \nsubseteq T_\beta(j)$ for any $\langle i, j\rangle$. On the other hand, suppose that i is least such that for some j, $T_\beta(j) \subseteq T_\alpha(i)$. If $k > i$, then $T_\alpha(k) = S'_\alpha(r_k)$ and r_k was chosen precisely so that $S'_\alpha(r_k) \not\supseteq T_\beta(l)$ for any

$l \in \omega$. Moreover, there can be at most one j such that $T_\alpha(i) \supseteq T_\beta(j)$, since by (1$\mathfrak{c}$), $T_\beta(j) \perp T_\beta(l)$, $j \neq l$. This completes the recursion.

Let $L = \{s_\alpha : T_\alpha$ has been defined$\}$ be a set of pairwise distinct points disjoint form B and let $P = B \cup L$. We topologize P by letting B be an open subspace with the usual Baire metric topology and declaring the n^{th} basic open set about the point s_α to be the set $N(\alpha, n) = \{s_\alpha\} \cup \bigcup_{m \geq n} [T_\alpha(m)]$.

If $\mathcal{T}_\alpha = \{[T_\alpha(n)] : n \in \omega\}$, then condition (1$\mathfrak{c}$) ensures that each $\mathcal{T}_\alpha$ is a pairwise disjoint collection, (2$\mathfrak{c}$) ensures that each basic open set $[f]$ occurs in at most one $\mathcal{T}_\alpha$, and (3$\mathfrak{c}$) ensures that if $N(\alpha, n)$ meets $N(\beta, m)$, then $N(\alpha, n) \cap N(\beta, m) = [T_\alpha(j)] \cap [T_\beta(k)]$ for some $j \geq n$ and $k \geq m$.

That P has a sharp base follows exactly as for the example due to Alleche *et al.* Let $\mathcal{B}_B$ be a sharp base for B and let

$$\mathcal{B} = \mathcal{B}_B \cup \{N(\alpha, n) : s_\alpha \in L \text{ and } n \in \omega\}.$$

Suppose $x \in \bigcap_{k \in \omega} B_k$ for some (injective) sequence $\{B_k \in \mathcal{B} : k \in \omega\}$. Since $\mathcal{B}_B$ is a sharp base and $s_\alpha \in N \in \mathcal{B}$ if and only if $N = (\alpha, n)$ for some n, the only case that is not obvious is when $x \in B$ and $B_k = N(\alpha_k, m_k)$ for all but finitely many k. But in this case condition (3$\mathfrak{c}$) implies that, for $n \geq 1$, $\bigcap_{k \leq n} B_k = \bigcap_{k \leq n} [T_{\alpha_k}(j_k)]$. Moreover (2$\mathfrak{c}$) implies that $T_{\alpha_k}(j_k) \neq T_{\alpha_{k'}}(j_{k'})$, so that $\{\bigcap_{k \leq n} B_k : n \in \omega\}$ contains a strictly decreasing subsequence and is therefore a base at x.

Since the set $\{s_\alpha : \alpha \in \mathfrak{c}\}$ is infinite, closed discrete, P is not compact. On the other hand, P is pseudocompact (so P is not metrizable). To see this, suppose that φ is a continuous real-valued function on P taking values in $[n, \infty)$ for each $n \in \omega$. Since B is dense in P, for each $n \in \omega$, there is some x_n in B such that $\varphi(x_n) > n$. By continuity, $\{x_n : n \in \omega\}$ does not have a limit point in B. Since φ is continuous and B is metrizable, there are basic open sets $[f_n]$ for each $n \in \omega$ such that $x_n \in [f_n] \subseteq \varphi^{-1}(n, \infty)$ and $\{[f_n] :\in \omega\}$ is a disjoint collection. But in this case $f_n \perp f_m$ when $n \neq m$ so that $\{f_n : n \in \omega\} = S_\alpha$ for some $\alpha \in \mathfrak{c}$. In which case, either s_α and T_α were defined or s_α was not defined and, for some $\beta < \alpha$, $T_\beta(j) \subseteq S_\alpha(n) = f_n$ for infinitely many n. In the second case, each basic open neighbourhood $N(\beta, n)$ of s_β contains infinitely many of the sets $[f_n]$. In the first case, T_α was chosen so that $T_\alpha(i) \supseteq f_{r_i}$ for each $i \in \omega$, so that $[T_\alpha(i)] \subseteq [f_{r_i}]$. In either case, each neighbourhood of s_β or s_α contains points which take arbitrarily large values under φ, contradicting continuity.

Now suppose for a contradiction that $P \times [0,1]$ has a sharp base. We shall show that this would imply that P has a σ-point finite base, which is impossible since Uspenskiĭ [18] shows that a pseudocompact space with a σ-point finite base is metrizable.

To this end, let $\mathcal{W}$ be a sharp base for $P \times [0,1]$ and let $\mathcal{C}$ be a countable sharp base for $[0,1]$. For each x in L choose W_n^x in $\mathcal{W}$, B_n^x in $\mathcal{B}$ (the sharp base for P), and C_n^x in $\mathcal{C}$ such that $B_n^x \times C_n^x \subseteq W_n^x$, $\{W_n^x : n \in \omega\}$ (and hence $\{B_n^x \times C_n^x : n \in \omega\}$) is a local base at $(x, 1/2)$ and $W_0^x \cap (L \times [0,1]) \subseteq \{x\} \times [0,1]$, which is possible since L is a closed discrete subset of P.

Let

$$\mathcal{B}_C = \{B \in \mathcal{B} : \text{for some } n \in \omega \text{ and some } x \in L,\, B = B_n^x \text{ and } C = C_n^x\}.$$

If $\mathcal{B}_C$ is not point finite then for some y in P, $y \in \bigcap_{j\in\omega} B_j$ for some pairwise distinct $B_j \in \mathcal{B}_C$. By definition, for each j there is some $x_j \in L$ and $n_j \in \omega$ such that $B_j = B_{n_j}^{x_j}$ and $C = C_{n_j}^{x_j}$. But then

$$\{y\} \times C \subseteq \bigcap_{j\in\omega} (B_{n_j}^{x_j} \times C_{n_j}^{x_j}) \subseteq \bigcap_{j\in\omega} W_{n_j}^{x_j}.$$

Since $B_j \neq B_k$, either there is an infinite set $J \subseteq \omega$ such that $x_j \neq x_k$, for distinct $j, k \in J$, or there is an infinite set $K \subseteq \omega$ such that $x_j = x_k = x$ but $n_j \neq n_k$ for some $x \in L$ and distinct $j, k \in K$. In the first case, $\{W_{n_j}^{x_j} : j \in J\}$ is a pairwise distinct subset of the sharp base $\mathcal{W}$ and $\bigcap_{j\in J} W_{n_j}^{x_j}$ contains at most one point. In the second case $\bigcap_{k\in K}(B_{n_k}^{x_k} \times C_{n_k}^{x_k}) = (x, 1/2)$, since $\{B_n^x \times C_n^x : n \in \omega\}$ is a local base at $(x, 1/2)$. In either case, $\{y\} \times C$ contains at most one point, which is not the case, and $\mathcal{B}_C$ is point finite.

Since $\{B_n^x \times C_n^x : n \in \omega\}$ is a local base at $(x, 1/2)$ and $\mathcal{C}$ is countable, $\mathcal{B} = \bigcup_{C\in\mathcal{C}} \mathcal{B}_C$ is a σ-point finite base for points of L. But $P = B \cup L]$ and B is a metric space, so P has a σ-point finite base: a contradiction.

By Theorem 4, P does not have a G_δ^* diagonal, nor indeed is it submetacompact. $\square$

So when is a pseudocompact space with a sharp base metrizable? As mentioned above, a pseudocompact, CCC regular space with a sharp base is metrizable [4, Theorem 21]. Pseudocompact, Moore spaces are CCC. Moreover, in proving that a pseudocompact Tychonoff space with a regular G_δ-diagonal is metrizable, McArthur [14] proves that a pseudocompact space with a G_δ^*-diagonal is developable. Hence we have

Theorem 4. *A pseudocompact regular space X with a sharp base is metrizable if either of the following hold:*

(1) *X is developable, or;*
(2) *X has a G_δ^*-diagonal.*

A pseudocompact space with a G_δ-diagonal is Čech complete [4, Lemma 20], hence Baire, so the following theorem is a strengthening of Theorem 21 of [4]. A space is strongly quasi-complete if there is a map g assigning to each $x \in X$ and $n \in \omega$ an open set $g(n, x)$ containing x such that $\{x_n\}$ clusters at x whenever $\{x, x_n\} \subseteq \bigcap_{i\leq n} g(i, y_i)$. Weakly developable spaces are clearly strongly quasi-complete.

Theorem 5. *A regular, locally CCC, locally Baire space with a sharp base is metrizable.*

Proof. Let X be a regular, locally CCC, locally Baire space with a sharp base. Since X has a weak development, it is strongly quasi-complete. Hodel [12] shows that every regular, quasi-complete CCC Baire space with either a

G_δ-diagonal or a point countable separating open cover is separable. Since X has a sharp base, X has a point countable base, a G_δ-diagonal and is quasi-complete. Hence X is locally separable. But every locally separable regular space with a point countable base is a disjoint union of clopen subspaces each of which has a countable base (see Theorem 7.2 of [10]). Hence X is metrizable. □

Generalising the fact that a countably compact space with a sharp base is metrizable we have:

Theorem 6. *A regular, ω_1-compact space with a sharp base is metrizable.*

Proof. Since X is ω_1-compact, every point-countable open cover of X has a countable subcover (Lemma 7.5, [10]). Since X has a sharp base, it has a point countable base and therefore is Lindelöf. A metacompact space with a sharp base is developable [2] and so a Lindelöf space with a sharp base is metrizable. □

Not surprisingly a monotonically normal space with a sharp base is metrizable (c.f. [6] where it is shown that a GO-space with a sharp base is metrizable).

Theorem 7. *For a monotonically normal space X the following are equivalent:*

(1) *X is metrizable;*
(2) *X has a sharp base;*
(3) *X has a weak development;*
(4) *X is strongly quasi-complete;*
(5) *X has a base of countable order and a G_δ-diagonal.*

Proof. Since $1 \implies 2 \implies 3 \implies 4 \implies 5$ (that 4 implies 5 follows from Theorems 2.2 and 2.3 of [8]), it remains to show that a monotonically normal space with a base of countable order and a G_δ-diagonal is metrizable. By the Balogh-Rudin theorem [5], since a stationary set of a regular cardinal does not have a G_δ-diagonal, a monotonically normal space with a G_δ-diagonal is paracompact. The result then follows since a paracompact space with a base of countable order is metrizable [3]. □

The proof that $P \times [0,1]$ does not have a sharp base does not quite extend to a proof that if the product of a space X with $[0,1]$ has a sharp base then X has a σ-point finite base. The converse however is easily seen to be true.

Proposition 8. *If a space X has a σ-point finite sharp base then $X \times [0,1]$ has a sharp base.*

Proof. Suppose that $\mathcal{B} = \bigcup \mathcal{B}_n$ is a σ-point finite sharp base for X and $\mathcal{C} = \bigcup \mathcal{C}_n$ is a development for $[0,1]$ such that each $\mathcal{C}_{n+1}$ is finite and refines $\mathcal{C}_n$ (so that $\mathcal{C}$ is also a sharp base for $[0,1]$). For each $n \in \omega$ let $\mathcal{W}_n = \{B \times C : B \in \mathcal{B}_n, C \in \mathcal{C}_n\}$ and let $\mathcal{W} = \bigcup_n \mathcal{W}_n$.

Firstly note that $\mathcal{W}$ is a base for $X \times [0,1]$. If (x,r) is in some open set U, choose n and $B \in \mathcal{B}_m$ such that $(x,r) \in B \times st(r, \mathcal{C}_n) \subseteq U$. Now for some $k \geq \max\{m,n\}$, there is $B' \in \mathcal{B}_k$ $x \in B' \subseteq B$. But then, since $\mathcal{C}_k$ refines $\mathcal{C}_n$, if $r \in C \in \mathcal{C}_k$, $B' \times C \in \mathcal{W}_k$ and

$$(x,r) \in B' \times C \subseteq B' \times st(r, \mathcal{C}_k) \subseteq B \times st(r, \mathcal{C}_n) \subset U.$$

Now suppose that $(x,r) \in B_j \times C_j = W_j \in \mathcal{W}$ for distinct W_j, $j \in \omega$. Each $\mathcal{W}_n$ is a point finite family since both $\mathcal{B}_n$ and $\mathcal{C}_n$ are point finite and so both $\{B_j\}_{j \in \omega}$ and $\{C_j\}_{j \in \omega}$ are infinite. Since $\mathcal{B}$ and $\mathcal{C}$ are sharp bases, this implies that $\{\bigcap_{j \leq n} B_j \times C_j : n \in \omega\}$ is a local base at (x,r) and $\mathcal{W}$ is a sharp base as required. □

Ponomarev, see [10], characterized those spaces with a point countable base as precisely the open s-images of metric spaces (a map is an s-map if it has separable fibres). There is a similar characterization for sharp bases.

Theorem 9. *A space X has a sharp base if and only if there is a metric space M with a base $\mathcal{B}$ and a continuous open mapping $f : M \to X$ such that, whenever $x \in X$ and $\{B_n \in \mathcal{B} : n \in \omega\}$ is a pairwise distinct collection, if $f^{-1}(x) \cap B_n \neq \varnothing$ for each $n \in \omega$, then there exists n_0 such that for each $y \in X$, if $f^{-1}(y) \cap B_j \neq \varnothing$, for each $j \leq n_0$, then $f^{-1}(y) \cap B_0 \neq \varnothing$.*

Proof. Suppose that $\mathcal{G}$ is a sharp base for the space X. Let

$$M = \{(G_n) \in \mathcal{G}^\omega : x \in \bigcap_{n \in \omega} G_n \text{ for some } x \in X\}$$

be the subspace of the Baire metric space $\mathcal{G}^\omega$, with metric $d((G_n),(H_n)) = 1/2^k$ where k is least such that $G_n \neq H_n$. Let $f : M \to X$ be defined letting $f((G_n))$ be the unique element of $\bigcap_{n \in \omega} G_n$ and let $\mathcal{B}$ be the base for M consisting of all $1/2^n$-balls about points of M. Then f is easily seen to be a continuous, open mapping onto X and the condition on $\mathcal{B}$ in the statement of the theorem is merely a translation of the fact that $\mathcal{G}$ is a sharp base. □

It is clear from the proof that, in the statement of the theorem, we can take $\mathcal{B}$ to be the collection of $1/2^n$ balls for any n rather than a base for M. Since a space with a sharp base has a point countable sharp base, we can also assume that the map in the statement of the theorem is an s-map. However, it is not immediately clear that we can prove that a space with a sharp base has a point countable base directly from the theorem.

We conclude with some open problems. Since every collectionwise normal Moore space is metrizable, the following is a natural and intriguing question.

Question 1. Is every collectionwise normal space with a sharp base metrizable?

Example 4 of [2] shows that weakly developable, collectionwise normal spaces do not have to be metrizable and the Heath V-space over a Q-set is an example of a normal space with a uniform base that is not metrizable. On the other hand, the answer is 'yes' if the space is also submetacompact (since it is then a Moore space) or a strict p-space. We might also ask whether a perfect, collectionwise normal space with a sharp base is metrizable. It is interesting to note that it is not known whether a collectionwise normal space with a point countable base need be paracompact.

Since the Heath V-space over a Δ-set is countably paracompact but not normal [13], at least consistently a countably paracompact, (Moore) space with a sharp base need not be normal. What about the converse?

Question 2. Is there a Dowker space with a sharp base?

Question 3. Is every perfectly regular space with a sharp base developable? Is every normal space with a sharp base developable? Is every perfectly regular, pseudocompact space with a sharp base metrizable?

Not every Moore space with a weakly uniform base has a uniform base (see [2]) so we ask:

Question 4. Does every Moore space with a sharp base have a uniform base?

Every pseudocompact space with a G_δ-diagonal is Čech complete [4], and every pseudocompact Moore space with a sharp base is metrizable.

Question 5. Is every Čech complete Moore space with a sharp base metrizable? What about Baire instead of Čech complete?

Question 6. If $X \times [0,1]$ has a sharp base, does X have a σ-point finite sharp base?

Question 7. Does the image (or pre-image) of a space with a sharp base under a perfect map (closed and open map, open map with compact, countable or finite fibres) have a sharp base?

Acknowledgements. Much of this research took place whilst the first author held a Universitas 21 Traveling Fellowship at the Department of Mathematics at the University of Auckland. He would like to thank the department for their hospitality and both the department and U21 for their financial support.

References

[1] P. Aleksandrov, *On the metrisation of topological spaces*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astr. Phys. **8** (1960), 135–140. MR 22 #5024

[2] B. Alleche, A. V. Arhangel′skiĭ, and J. Calbrix, *Weak developments and metrization*, Topology Appl. **100** (2000), no. 1, 23–38, Special issue in honor of Howard H. Wicke. MR **2001g:**54029

[3] A. V. Arhangel′skiĭ, *Some metrization theorems*, Uspehi Mat. Nauk **18** (1963), no. 5 (113), 139–145. MR 27 #6242

[4] A. V. Arhangel′skiĭ, W. Just, E. A. Rezniczenko, and P. J. Szeptycki, *Sharp bases and weakly uniform bases versus point-countable bases*, Topology Appl. **100** (2000), no. 1, 39–46, Special issue in honor of Howard H. Wicke. MR **2001f:**54027

[5] Z. Balogh and M. E. Rudin, *Monotone normality*, Topology Appl. **47** (1992), no. 2, 115–127. MR **94b:**54065

[6] Harold Bennett and David Lutzer, *Ordered spaces with special bases*, Fund. Math. **158** (1998), no. 3, 289–299. MR **99i:**54044

[7] ______, *Continuous separating families in ordered spaces and strong base conditions*, Topology Appl. **119** (2002), 305–314.

[8] Raymond F. Gittings, *Concerning quasi-complete spaces*, General Topology and Appl. **6** (1976), no. 1, 73–89. MR 52 #11855

[9] Chris Good, Robin W. Knight, and Abdul M. Mohamad, *On the metrizability of spaces with a sharp base*, To appear in Topology Appl. PII: S0166-8641(01)00300-5, 2002.

[10] Gary Gruenhage, *Generalized metric spaces*, Handbook of set-theoretic topology, North-Holland, Amsterdam, 1984, pp. 423–501. MR **86h:**54038

[11] R. W. Heath and W. F. Lindgren, *Weakly uniform bases*, Houston J. Math. **2** (1976), no. 1, 85–90. MR 52 #15364

[12] R. E. Hodel, *Metrizability of topological spaces*, Pacific J. Math. **55** (1974), 441–459. MR 51 #6747

[13] Robin W. Knight, *δ-sets*, Trans. Amer. Math. Soc. **339** (1993), no. 1, 45–60. MR **94a:**54016

[14] William G. McArthur, *G_δ-diagonals and metrization theorems*, Pacific J. Math. **44** (1973), 613–617. MR 47 #5835

[15] S. A. Peregudov, *On pseudocompactness and other covering properties*, Questions Answers Gen. Topology **17** (1999), no. 2, 153–155. MR **2000h:**54033

[16] Brian M. Scott, *Pseudocompact, metacompact spaces are compact*, Topology Proc. **4** (1979), no. 2, 577–587 (1980), The Proceedings of the 1979 Topology Conference (Ohio Univ., Athens, Ohio, 1979). MR **81m:**54034

[17] D. B. Shakhmatov, *Pseudocompact spaces with a point-countable base*, Dokl. Akad. Nauk SSSR **279** (1984), no. 4, 825–829, English translation in Soviet Math. Dokl. 30 (1984), no. 3, 747–751. MR **86a:**54029

[18] V. V. Uspenskiĭ, *Pseudocompact spaces with a σ-point-finite base are metrizable*, Comment. Math. Univ. Carolin. **25** (1984), no. 2, 261–264. MR **87f:**54035

[19] W. Stephen Watson, *Pseudocompact metacompact spaces are compact*, Proc. Amer. Math. Soc. **81** (1981), no. 1, 151–152. MR **81j:**54042

[20] ______, *A pseudocompact meta-Lindelöf space which is not compact*, Topology Appl. **20** (1985), no. 3, 237–243. MR **87d:**54042

School of Mathematics and Statistics, University of Birmingham, Birmingham, B15 2TT, UK

E-mail address: c.good@bham.ac.uk

Mathematical Institute, University of Oxford, 24-29 St Giles', Oxford OX1 3LB, UK

E-mail address: knight@maths.ox.ac.uk

Department of Mathematics, University of Auckland, Private Bag 92019, Auckland, New Zealand

E-mail address: mohamad@math.auckland.ac.nz