

2
3 Fixed Points Of (ψ, φ) -Almost Weakly Contractive
4 Maps In G-Metric Spaces*

5 Gutti Venkata Ravindranadh Babu[†], Dasari Ratna Babu[‡],
6 Kanuri Nageswara Rao[§], Bendi Venkata Siva Kumar[†]

7 Received 26 August 2013

8 **Abstract**

9 In this paper, we introduce (ψ, φ) -almost weakly contractive maps in G-metric
10 spaces and prove the existence of fixed points. Our Theorem 4 generalizes the
11 result of Aage and Salunke (Theorem 2, [1]). We also extend it to a pair of weakly
12 compatible maps and prove the existence of common fixed points. We provide
examples in support of our results.

13 **1 Introduction and Preliminaries**

14 The development of fixed point theory is based on the generalization of contraction
15 conditions in one direction or/and generalization of ambient spaces of the operator
16 under consideration on the other. Banach contraction principle plays an important role
17 in solving nonlinear equations, and it is one of the most useful results in fixed point
18 theory. In the direction of generalization of contraction conditions, in 1997, Alber
19 and Guerre-Delabriere [3] introduced weakly contractive maps which are extensions
20 of contraction maps and obtained fixed point results in the setting of Hilbert spaces.
21 Rhoades [16] extended this concept to metric spaces. In 2008, Dutta and Choudhury
22 [12] introduced (ψ, φ) -weakly contractive maps and proved the existence of fixed points
23 in complete metric spaces. In 2009, Doric [11] extended it to a pair of maps. For more
24 literature in this direction, we refer to Choudhury, Konar and Rhoades [9], Babu,
25 Nageswara Rao and Alemayehu [4], Sastry, Babu and Kidane [17], Babu and Sailaja
26 [5] and Zhang and Song [19]. In continuation to the extensions of contraction maps,
27 Berinde [7] introduced ‘weak contractions’ as a generalization of contraction maps.
28 Berinde renamed ‘weak contractions’ as ‘almost contractions’ in his later work [8]. For
29 more works on almost contractions and its generalizations, we refer to Babu, Sandhya
30 and Kameswari [6], Abbas, Babu and Alemayehu [2] and the related references cited
31 in these papers.

32 Throughout this paper, we denote $\mathbb{R}_+ = [0, \infty)$ and

33
$$\Psi = \{\psi/\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+ \text{ is continuous on } \mathbb{R}_+, \psi \text{ is nondecreasing,}$$

34
$$\psi(t) > 0 \text{ for } t > 0, \psi(0) = 0\}.$$

*Mathematics Subject Classifications:47H10, 54H25.

†Department of Mathematics, Andhra University, Visakhapatnam-530 003, India

‡Department of Mathematics, PSCMR CET, Vijayawada-520 001, India

§Department of Mathematics, GIET, Rajahmundry, India

35 In the metric space setting Dutta and Choudhury [12] introduced (ψ, φ) -weakly con-
36 tractive maps as follows:

37 DEFINITION 1 ([12]). Let (X, d) be a metric space. Let $T : X \rightarrow X$ be a map. If
38 there exist $\psi, \varphi \in \Psi$ such that

$$39 \quad \psi(d(Tx, Ty)) \leq \psi(d(x, y)) - \varphi(d(x, y))$$

40 for all $x, y \in X$, then T is said to be a (ψ, φ) -weakly contractive map.

41 Dutta and Choudhury [12] proved that every (ψ, φ) -weakly contractive map has a
42 unique fixed point in complete metric spaces. On the other hand, Berinde [7] introduced
43 'weak contractions' as a generalization of contraction maps.

44 DEFINITION 2 ([7]). Let (X, d) be a metric space. A selfmap $T : X \rightarrow X$ is said
45 to be a weak contraction if there exist $\delta \in (0, 1)$ and $L \geq 0$ such that for all $x, y \in X$,

$$46 \quad d(Tx, Ty) \leq \delta d(x, y) + L d(y, Tx).$$

47 Berinde [7] proved that every weak contraction has a fixed point in complete metric
48 spaces and provided an example to show that this fixed point need not be unique. In
49 order to obtain the uniqueness of fixed point, Berinde [7] used the following condition:
50 there exist $\theta \in (0, 1)$ and $L_1 \geq 0$ such that

$$51 \quad d(Tx, Ty) \leq \theta d(x, y) + L_1 d(x, Tx) \text{ for all } x, y \in X \quad (1)$$

52 and proved that every weak contraction together with (1) has a unique fixed point in
53 complete metric spaces, and further posed the following problem: "Find a contractive
54 type condition different from (1), that ensures the uniqueness of fixed point of weak
55 contractions".

56 In this context Babu, Sandhya and Kameswari [6] answered the above problem by
57 introducing 'condition (B)' as follows:

58 DEFINITION 3 ([6]). Let (X, d) be a metric space. A map $T : X \rightarrow X$ is said to
59 satisfy condition (B) if there exist $0 < \delta < 1$ and $L \geq 0$ such that for all $x, y \in X$,

$$60 \quad d(Tx, Ty) \leq \delta d(x, y) + L \min\{d(x, Tx), d(y, Ty), d(x, Ty), d(y, Tx)\}.$$

61 Babu, Sandhya and Kameswari [6] proved that every selfmap T of a complete
62 metric space satisfying condition (B) has a unique fixed point. On the other hand,
63 in the direction of generalization of ambient spaces, in 2005, Mustafa and Sims [15]
64 introduced a new notion namely generalized metric space called G -metric space and
65 studied the existence of fixed points of various types of contraction mappings in G -
66 metric spaces.

67 DEFINITION 4 ([15]). Let X be a nonempty set and let $G : X^3 \rightarrow \mathbb{R}_+$ be a
68 function satisfying:

69 (G1) $G(x, y, z) = 0$ if $x = y = z$,
 70 (G2) $0 < G(x, x, y)$ for all $x, y \in X$, with $x \neq y$,
 71 (G3) $G(x, x, y) \leq G(x, y, z)$ for all $x, y, z \in X$ with $z \neq y$
 72 (G4) $G(x, y, z) = G(x, z, y) = G(y, z, x) = \dots$ (symmetry in all variables) and,
 73 (G5) $G(x, y, z) \leq G(x, a, a) + G(a, y, z)$ for all $x, y, z, a \in X$ (rectangle inequality).

74 Then the function G is called a generalized metric, or, more specially a G -metric on
 75 X , and the pair (X, G) is called a G -metric space.

76 EXAMPLE 1 ([15]). Let (X, d) be a metric space. The mapping $G_s : X^3 \rightarrow \mathbb{R}_+$
 77 defined by

$$78 \quad G_s(x, y, z) = d(x, y) + d(y, z) + d(x, z)$$

79 for all $x, y, z \in X$ is a G -metric and so (X, G_s) is a G -metric space.

80 EXAMPLE 2 ([15]). Let (X, d) be a metric space. The mapping $G_m : X^3 \rightarrow \mathbb{R}_+$
 81 defined by

$$82 \quad G_m(x, y, z) = \max \{d(x, y), d(y, z), d(x, z)\}$$

83 for all $x, y, z \in X$ is a G -metric and so (X, G_m) is a G -metric space.

84 EXAMPLE 3. Let X be a nonempty set. We denote the class of all real valued
 85 bounded functions on X by $B(X)$. For $f \in B(X)$, we define

$$86 \quad \|f\| = \sup \{|f(x)| / x \in X\}.$$

87 Then $(B(X), \|\cdot\|)$ is a normed linear space. We define metric d on $B(X)$ by $d(f, g) =$
 88 $\|f - g\|$ for $f, g \in B(X)$. Now we define generalized metric G on $B(X)$ by

$$89 \quad G(f, g, h) = \|f - g\| + \|g - h\| + \|h - f\|$$

90 for all $f, g, h \in B(X)$. Then clearly G is a generalized metric on $B(X)$. The space
 91 $(B(X), G)$ is a generalized metric space.

92 DEFINITION 5 ([15]). Let (X, G) be a G -metric space and let $\{x_n\}$ be a sequence
 93 of points of X . We say that $\{x_n\}$ is G -convergent to x if $\lim_{n, m \rightarrow \infty} G(x, x_n, x_m) = 0$;
 94 that is, for any $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that $G(x, x_n, x_m) < \epsilon$ for all $n, m \geq N$.
 95 We refer to x as the limit of the sequence $\{x_n\}$.

96 PROPOSITION 1 ([15]). Let (X, G) be a G -metric space. Then for any $x, y, z, a \in$
 97 X we have that:

98 (1) if $G(x, y, z) = 0$, then $x = y = z$.
 99 (2) $G(x, y, z) \leq G(x, x, y) + G(x, x, z)$.

100 (3) $G(x, y, y) \leq 2G(y, x, x)$.
 101 (4) $G(x, y, z) \leq G(x, a, z) + G(a, y, z)$.
 102 (5) $G(x, y, z) \leq \frac{2}{3} (G(x, y, a) + G(x, a, z) + G(a, y, z))$.

103 PROPOSITION 2 ([15]). Let (X, G) be a G -metric space. Then the following
 104 statements are equivalent:

105 (1) $\{x_n\}$ is G -convergent to x .
 106 (2) $G(x_n, x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.
 107 (3) $G(x_n, x, x) \rightarrow 0$ as $n \rightarrow \infty$.

108 DEFINITION 6 ([15]). Let X be a G -metric space. A sequence $\{x_n\}$ is called G -
 109 Cauchy if given $\epsilon > 0$, there is an $N \in \mathbb{N}$ such that $G(x_n, x_m, x_l) < \epsilon$ for all $n, m, l \geq N$;
 110 that is, if $G(x_n, x_m, x_l) \rightarrow 0$ as $n, m, l \rightarrow \infty$.

111 PROPOSITION 3 ([15]). In a G -metric space X , the following two statements are
 112 equivalent:

113 (1) The sequence $\{x_n\}$ is G -Cauchy.
 114 (2) For every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that $G(x_n, x_m, x_m) < \epsilon$ for all $n, m \geq N$.

115 DEFINITION 7 ([15]). A G -metric space X is said to be G -complete (or a complete
 116 G -metric space) if every G -Cauchy sequence in X is G -convergent in X .

117 PROPOSITION 4 ([15]). Let X be a G -metric space. Then the function $G(x, y, z)$
 118 is jointly continuous in all three of its variables.

119 PROPOSITION 5 ([15]). Every G -metric space X defines a metric space (X, d_G)
 120 by

$$121 \quad d_G(x, y) = G(x, y, y) + G(y, x, x) \text{ for all } x, y \in X.$$

122 Mustafa, Obiedat and Awawdeh [14] proved the following result.

123 THEOREM 1 ([14]). Let (X, G) be a complete G -metric space, and let $T : X \rightarrow X$
 124 be a mapping satisfying one of the following conditions:

$$125 \quad G(Tx, Ty, Tz) \leq aG(x, y, z) + bG(x, Tx, Tx) + cG(y, Ty, Ty) + dG(z, Tz, Tz)$$

126 or

$$127 \quad G(Tx, Ty, Tz) \leq aG(x, y, z) + bG(x, x, Tx) + cG(y, y, Ty) + dG(z, z, Tz)$$

128 for all $x, y, z \in X$ where $0 \leq a + b + c + d < 1$. Then T has a unique fixed point (say
 129 u , i.e., $Tu = u$), and T is G -continuous at u .

130 In 2011, Aage and Salunke [1] introduced weakly contractive maps in G -metric
 131 spaces and proved the existence of fixed points in G -metric spaces.

132 DEFINITION 8. Let (X, G) be a G -metric space. Let $T : X \rightarrow X$ be a selfmap of
 133 X . T is said to be a weakly contractive map in G if, there exists $\varphi \in \Psi$ such that

134
$$G(Tx, Ty, Tz) \leq G(x, y, z) - \varphi(G(x, y, z)) \text{ for each } x, y, z \in X. \quad (2)$$

135 THEOREM 2 ([1]). Let (X, G) be a complete G -metric space and let $T : X \rightarrow X$
 136 be a weakly contractive map in G . Then T has a unique fixed point in X .

137 DEFINITION 9 ([10]). Let f and g be two selfmaps on a G -metric space (X, G) .
 138 The mappings f and g are said to be *compatible* if $\lim_{n \rightarrow \infty} G(fgx_n, gfx_n, gfx_n) = 0$
 139 whenever $\{x_n\}$ is a sequence in X such that $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = z$ for some
 140 $z \in X$.

141 DEFINITION 10 ([10,13]). Two maps f and g on a G -metric space (X, G) are said
 142 to be *weakly compatible* if they commute at their coincidence point.

143 Here we note that every pair of compatible maps is weakly compatible but its
 144 converse need not be true (Example 1.4, [10]). Shatanawi [18] proved the following
 145 common fixed point theorem for a pair of weakly compatible maps.

146 THEOREM 3 ([18]). Let X be a G -metric space. Suppose the maps $f, g : X \rightarrow$
 147 X satisfy the following condition: there exists a nondecreasing function $\phi : \mathbb{R}_+ \rightarrow$
 148 \mathbb{R}_+ satisfying $\lim_{n \rightarrow \infty} \phi^n(t) = 0$ for all $t \in (0, \infty)$ such that either

149
$$G(fx, fy, fz) \leq \phi(\max\{G(gx, gy, gz), G(gx, fx, fx), G(gy, fy, fy), G(gz, fz, fz)\}), \quad (3)$$

150 OR

151
$$G(fx, fy, fz) \leq \phi(\max\{G(gx, gy, gz), G(gx, gx, fx), G(gy, gy, fy), G(gz, gz, fz)\})$$

152 for all $x, y, z \in X$. If $f(X) \subseteq g(X)$ and $g(X)$ is a G -complete subspace of X , then
 153 f and g have a unique point of coincidence in X . Moreover, if f and g are weakly
 154 compatible, then f and g have a unique common fixed point.

155 Unfortunately, the example given in support of Theorem 2 by Aage and Salunke
 156 (Example 2, [1]) is false in the sense that the maps T and φ defined in this example
 157 do not satisfy the inequality (2). For, the example considered by Aage and Salunke is
 158 the following.

159 EXAMPLE 4 ([1]). Let $X = [0, 1]$. Define $G : X^3 \rightarrow \mathbb{R}_+$ by

160
$$G(x, y, z) = |x - y| + |y - z| + |z - x| \text{ for all } x, y, z \in X.$$

161 Then (X, G) is a complete G -metric space. The authors defined T on X by $Tx = x - \frac{x^2}{2}$
 162 and $\varphi(t) = \frac{t^2}{2}$, $t \geq 0$. Let us choose $x = 1$, $y = \frac{1}{2}$ and $z = \frac{1}{4}$. Then $G(Tx, Ty, Tz) = \frac{9}{16}$,
 163 $G(x, y, z) = \frac{3}{2}$ and $\varphi(G(x, y, z)) = \frac{9}{8}$. Hence

$$164 \quad \frac{9}{16} = G(Tx, Ty, Tz) \not\leq G(x, y, z) - \varphi(G(x, y, z)) = \frac{3}{8}.$$

165 Also the inequality (2) fails to hold at $x = \frac{1}{2}$, $y = \frac{1}{3}$ and $z = 0$. Hence T and φ do not
 166 satisfy the inequality (2) so that T is not a weakly contractive map with this φ , even
 167 though T has a fixed point 0.

168 The following is a suitable example in support of Theorem 2.

169 EXAMPLE 5. Let $X = [0, 1]$. We define $G : X^3 \rightarrow \mathbb{R}_+$ by

$$170 \quad G(x, y, z) = \begin{cases} 0, & \text{if } x = y = z \\ \max\{x, y, z\}, & \text{otherwise.} \end{cases}$$

171 Then (X, G) is a complete G -metric space. Let $Tx = \frac{x^2}{2}$ and $\varphi(t) = \frac{t^2}{4}$. Without loss
 172 of generality, we assume that $x > y > z$. Then

$$173 \quad G(Tx, Ty, Tz) = \max\{Tx, Ty, Tz\} = \frac{x^2}{2},$$

$$174 \quad G(x, y, z) = \max\{x, y, z\} = x \text{ and } \varphi(G(x, y, z)) = \frac{x^2}{4}.$$

176 Now, $G(x, y, z) - \varphi(G(x, y, z)) = x - \frac{x^2}{4}$. Therefore $G(Tx, Ty, Tz) = \frac{x^2}{2} < x - \frac{x^2}{4} =$
 177 $G(x, y, z) - \varphi(G(x, y, z))$. Hence T satisfies the inequality (2) so that T is a weakly
 178 contractive map. Thus by Theorem 2, we have T has a unique fixed point and it is 0
 179 in X .

180 Motivated by the ' (ψ, φ) -weakly contractive maps' introduced by Dutta and Choudhury [12], 'almost weak contractions' of Berinde [7, 8] and 'condition (B)' of Babu, Sandhya and Kameswari [6] in metric space setting, in this paper we introduce ' (ψ, φ) -almost
 181 weakly contractive maps' in G -metric spaces and prove the existence of fixed points in
 182 complete G -metric spaces. The importance of the class of (ψ, φ) -almost weakly con-
 183 tractive maps is that this class properly includes the class of weakly contractive maps
 184 studied by Aage and Salunke [1] so that the class of (ψ, φ) -almost weakly contractive
 185 maps is larger than the class of weakly contractive maps, which is illustrated in Ex-
 186 ample 6. Hence, the results obtained on the existence of fixed points of (ψ, φ) -almost
 187 weakly contractive maps generalize the results of Aage and Salunke [1].

188 In the following, we introduce (ψ, φ) -almost weakly contractive maps.

189 DEFINITION 11. Let (X, G) be a G -metric space and let T be a selfmap of X . If
 190 there exist ψ and φ in Ψ and $L \geq 0$ such that

$$191 \quad \psi(G(Tx, Ty, Tz)) \leq \psi(G(x, y, z)) - \varphi(G(x, y, z)) + L m(x, y, z) \quad (4)$$

194 for all $x, y, z \in X$, where

$$195 \quad m(x, y, z) = \min\{G(Tx, x, x), G(Tx, y, y), G(Tx, z, z), G(Tx, y, z)\},$$

196 then we call T is a (ψ, φ) -almost weakly contractive map on X .

197 We observe that if ψ is the identity map and $L = 0$ in (4) then T is a weakly
198 contractive map. Hence the class of all weakly contractive maps is contained in the
199 class of all (ψ, φ) -almost weakly contractive maps. Further, every (ψ, φ) -almost weakly
200 contractive map need not be a weakly contractive map (Example 6).

201 In Section 2, we prove the existence of fixed points of (ψ, φ) -almost weakly contractive
202 maps in G -metric spaces. Our main result (Theorem 4) generalizes the result of
203 Aage and Salunke (Theorem 2, [1]). We also extend it to a pair of weakly compatible
204 maps and prove the existence of common fixed points. Corollaries and examples in
205 support of our results are provided in Section 3.

206 2 Main Results

207 The following is the main result of this paper.

208 THEOREM 4. Let (X, G) be a complete G -metric space and let T be a (ψ, φ) -
209 almost weakly contractive map. Then T has a unique fixed point in X .

210 PROOF. Let $x_0 \in X$. We define the sequence $\{x_n\}$ by $x_n = T(x_{n-1})$, $n = 1, 2, \dots$
211 If $x_{n+1} = x_n$ for some $n \in \mathbb{N}$, then trivially x_n a fixed point of T . Suppose $x_{n+1} \neq x_n$
212 for all $n \in \mathbb{N}$. We now consider

$$213 \quad \begin{aligned} \psi(G(x_n, x_{n+1}, x_{n+1})) &= \psi(G(Tx_{n-1}, Tx_n, Tx_n)) \\ 214 &\leq \psi(G(x_{n-1}, x_n, x_n)) - \varphi(G(x_{n-1}, x_n, x_n)) \\ 215 &\quad + Lm(x_{n-1}, x_n, x_n), \end{aligned}$$

216 where $m(x_{n-1}, x_n, x_n) = 0$ so that

$$217 \quad \psi(G(x_n, x_{n+1}, x_{n+1})) \leq \psi(G(x_{n-1}, x_n, x_n)) - \varphi(G(x_{n-1}, x_n, x_n)). \quad (5)$$

218 By using the property of φ , we have

$$219 \quad \psi(G(x_n, x_{n+1}, x_{n+1})) < \psi(G(x_{n-1}, x_n, x_n)) \text{ for } n = 1, 2, \dots \quad (6)$$

220 Now, by applying the nondecreasing property of ψ , it follows that

$$221 \quad G(x_n, x_{n+1}, x_{n+1}) \leq G(x_{n-1}, x_n, x_n) \text{ for } n = 1, 2, \dots$$

222 Therefore $\{G(x_n, x_{n+1}, x_{n+1})\}$ is a monotone decreasing sequence of nonnegative reals
223 and hence there exists $r \geq 0$ such that $G(x_n, x_{n+1}, x_{n+1}) \rightarrow r$ as $n \rightarrow \infty$. Now, on
224 letting $n \rightarrow \infty$ in the inequality (5), we have $\psi(r) \leq \psi(r) - \varphi(r)$ so that $\varphi(r) \leq 0$.
225 Since $\varphi(r) \geq 0$, it follows that $\varphi(r) = 0$ so that $r = 0$.

$$226 \quad \text{i.e., } G(x_n, x_{n+1}, x_{n+1}) \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (7)$$

227 We now prove that the sequence $\{x_n\}$ is Cauchy.

228 On the contrary, if $\{x_n\}$ is not Cauchy, then there exists an $\epsilon > 0$ for which we can
229 find subsequences $\{x_{n_k}\}, \{x_{m_k}\}$ of $\{x_n\}$ with $n_k > m_k \geq k$ such that

$$230 \quad G(x_{n_k}, x_{m_k}, x_{m_k}) \geq \epsilon. \quad (8)$$

231 Corresponding to each m_k , we can choose n_k such that it is the smallest integer with
232 $n_k > m_k$ and satisfying (8). Then, we have

$$233 \quad G(x_{n_k}, x_{m_k}, x_{m_k}) \geq \epsilon \text{ and } G(x_{n_k-1}, x_{m_k}, x_{m_k}) < \epsilon. \quad (9)$$

234 We now prove the following three identities:

$$235 \quad (i) \lim_{k \rightarrow \infty} G(x_{n_k}, x_{m_k}, x_{m_k}) = \epsilon.$$

$$236 \quad (ii) \lim_{k \rightarrow \infty} G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1}) = \epsilon.$$

$$237 \quad (iii) \lim_{k \rightarrow \infty} G(x_{n_k}, x_{m_k-1}, x_{m_k-1}) = \epsilon.$$

238 From (9), we have $G(x_{n_k}, x_{m_k}, x_{m_k}) \geq \epsilon$ so that

$$239 \quad \epsilon \leq \liminf_{k \rightarrow \infty} G(x_{n_k}, x_{m_k}, x_{m_k}). \quad (10)$$

240 Also,

$$241 \quad G(x_{n_k}, x_{m_k}, x_{m_k}) \leq G(x_{n_k}, x_{n_k-1}, x_{n_k-1}) + G(x_{n_k-1}, x_{m_k}, x_{m_k}) \\ 242 \quad < G(x_{n_k}, x_{n_k-1}, x_{n_k-1}) + \epsilon,$$

243 and hence

$$244 \quad \limsup_{k \rightarrow \infty} G(x_{n_k}, x_{m_k}, x_{m_k}) \leq \epsilon. \quad (11)$$

245 From (10) and (11), we have

$$246 \quad \epsilon \leq \liminf_{k \rightarrow \infty} G(x_{n_k}, x_{m_k}, x_{m_k}) \leq \limsup_{k \rightarrow \infty} G(x_{n_k}, x_{m_k}, x_{m_k}) \leq \epsilon$$

247 so that $\lim_{k \rightarrow \infty} G(x_{n_k}, x_{m_k}, x_{m_k})$ exists and

$$248 \quad \epsilon = \liminf_{k \rightarrow \infty} G(x_{n_k}, x_{m_k}, x_{m_k}) \leq \limsup_{k \rightarrow \infty} G(x_{n_k}, x_{m_k}, x_{m_k}) = \epsilon.$$

249 Hence

$$250 \quad \lim_{k \rightarrow \infty} G(x_{n_k}, x_{m_k}, x_{m_k}) = \epsilon. \quad (12)$$

251 Therefore (i) holds. Also,

$$252 \quad G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1}) \leq G(x_{n_k-1}, x_{m_k}, x_{m_k}) + G(x_{m_k}, x_{m_k-1}, x_{m_k-1}) \\ 253 \quad \leq G(x_{n_k-1}, x_{n_k}, x_{n_k}) + G(x_{n_k}, x_{m_k}, x_{m_k}) \\ 254 \quad + G(x_{m_k}, x_{m_k-1}, x_{m_k-1}).$$

255 On taking limit superior as $k \rightarrow \infty$ and using (7) and (12), we get

$$256 \quad \limsup_{k \rightarrow \infty} G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1}) \leq \epsilon. \quad (13)$$

257 Now,

$$258 \quad G(x_{n_k}, x_{m_k}, x_{m_k}) \leq G(x_{n_k}, x_{n_k-1}, x_{n_k-1}) + G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1}) \\ 259 \quad + G(x_{m_k-1}, x_{m_k}, x_{m_k}).$$

260 Hence, we have that

$$261 \quad G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1}) \geq G(x_{n_k}, x_{m_k}, x_{m_k}) - G(x_{n_k}, x_{n_k-1}, x_{n_k-1}) \\ 262 \quad - G(x_{m_k-1}, x_{m_k}, x_{m_k}).$$

263 Now on taking limit inferior both sides, and using (7) and (12), we get

$$264 \quad \epsilon \leq \liminf_{k \rightarrow \infty} G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1}). \quad (14)$$

265 Thus from (13) and (14), we have

$$266 \quad \epsilon \leq \liminf_{k \rightarrow \infty} G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1}) \leq \limsup_{k \rightarrow \infty} G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1}) \leq \epsilon.$$

267 Hence it follows that

$$268 \quad \lim_{k \rightarrow \infty} G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1}) = \epsilon. \quad (15)$$

269 Therefore (ii) holds. Let us now prove (iii). From (8), we have

$$270 \quad \epsilon \leq G(x_{n_k}, x_{m_k}, x_{m_k}) \leq G(x_{n_k}, x_{m_k-1}, x_{m_k-1}) + G(x_{m_k-1}, x_{m_k}, x_{m_k}).$$

271 This implies that

$$272 \quad G(x_{n_k}, x_{m_k-1}, x_{m_k-1}) \geq \epsilon - G(x_{m_k-1}, x_{m_k}, x_{m_k})$$

273 and

$$274 \quad \liminf_{k \rightarrow \infty} G(x_{n_k}, x_{m_k-1}, x_{m_k-1}) \geq \epsilon. \quad (16)$$

275 Now,

$$276 \quad G(x_{n_k}, x_{m_k-1}, x_{m_k-1}) \leq G(x_{n_k}, x_{n_k-1}, x_{n_k-1}) + G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1})$$

277 and hence using (7) and (ii), we get

$$278 \quad \limsup_{k \rightarrow \infty} G(x_{n_k}, x_{m_k-1}, x_{m_k-1}) \leq \epsilon. \quad (17)$$

279 Now, from (16) and (17), we have

$$280 \quad \epsilon \leq \liminf_{k \rightarrow \infty} G(x_{n_k}, x_{m_k-1}, x_{m_k-1}) \leq \limsup_{k \rightarrow \infty} G(x_{n_k}, x_{m_k-1}, x_{m_k-1}) \leq \epsilon$$

281 so that $G(x_{n_k}, x_{m_k-1}, x_{m_k-1}) \rightarrow \epsilon$ as $k \rightarrow \infty$. Therefore (iii) holds.

282 Now

$$\begin{aligned}
 283 \quad \psi(G(x_{n_k}, x_{m_k}, x_{m_k})) &= \psi(G(Tx_{n_k-1}, Tx_{m_k-1}, Tx_{m_k-1})) \\
 284 &\leq \psi(G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1})) - \varphi(G(x_{n_k-1}, x_{m_k-1}, x_{m_k-1})) \\
 285 &\quad + L m(x_{n_k-1}, x_{m_k-1}, x_{m_k-1}).
 \end{aligned}$$

286 On letting $k \rightarrow \infty$ and using (i)–(iii) and (7), we get

$$287 \quad \psi(\epsilon) \leq \psi(\epsilon) - \varphi(\epsilon) < \psi(\epsilon),$$

288 which is a contradiction. Therefore $\{x_n\}$ is a G -Cauchy sequence. Since X is complete,
289 there exists $p \in X$ such that $\{x_n\}$ is G -convergent to p . We now consider

$$\begin{aligned}
 290 \quad \psi(G(x_n, Tp, Tp)) &= \psi(G(Tx_{n-1}, Tp, Tp)) \\
 291 &\leq \psi(G(x_{n-1}, p, p)) - \varphi(G(x_{n-1}, p, p)) + L m(x_{n-1}, p, p).
 \end{aligned}$$

292 On letting $n \rightarrow \infty$, we have $\varphi(G(p, Tp, Tp)) \leq 0$ so that we must have $Tp = p$.
293 Therefore p is a fixed point of T in X .

294 **Uniqueness:** Suppose T has two fixed points p and q in X with $p \neq q$. Now, we
295 consider

$$\begin{aligned}
 296 \quad \psi(G(p, q, q)) &= \psi(G(Tp, Tq, Tq)) \\
 297 &\leq \psi(G(p, q, q)) - \varphi(G(p, q, q)) + L m(p, q, q) \\
 298 &= \psi(G(p, q, q)) - \varphi(G(p, q, q)) < \psi(G(p, q, q)),
 \end{aligned}$$

299 which is a contradiction. Therefore $\psi(G(p, q, q)) = 0$ so that $G(p, q, q) = 0$ and hence
300 that $p = q$. Thus, p is the unique fixed point of T in X . Hence the theorem follows.

301 We now prove a common fixed point theorem for a pair of weakly compatible maps.

302 **THEOREM 5.** Let (X, G) be a complete G -metric space and let T and S be two
303 selfmaps on (X, G) . Assume that $T(X) \subseteq S(X)$, S is continuous, and there exist
304 $\psi, \varphi \in \Psi$ and $L \geq 0$ such that

$$305 \quad \psi(G(Tx, Ty, Tz)) \leq \psi(M(x, y, z)) - \varphi(M(x, y, z)) + L m(x, y, z), \quad (18)$$

306 where

$$307 \quad M(x, y, z) = \max\{G(Sx, Sy, Sz), G(Sx, Tx, Tx), G(Sy, Ty, Ty), G(Sz, Tz, Tz)\}$$

308 and

$$309 \quad m(x, y, z) = \min\{G(Tx, Sx, Sx), G(Tx, Sy, Sy), G(Tx, Sz, Sz), G(Tx, Sy, Sz)\}$$

310 for $x, y, z \in X$. Then T and S have a unique common fixed point in X provided T and
311 S are weakly compatible maps.

312 PROOF. Let $x_0 \in X$ be arbitrary. Since $T(X) \subseteq S(X)$, we can choose $\{x_n\} \subseteq X$
 313 such that $T(x_n) = S(x_{n+1}) = y_n$ (say), $n = 0, 1, 2, \dots$. Let $n \geq 1$ be an integer. Then
 314 by using the inequality (18) we have

$$\begin{aligned} 315 \quad \psi(G(y_n, y_{n+1}, y_{n+1})) &= \psi(G(Tx_n, Tx_{n+1}, Tx_{n+1})) \\ 316 \quad &\leq \psi(M(x_n, x_{n+1}, x_{n+1})) - \varphi(M(x_n, x_{n+1}, x_{n+1})) \\ 317 \quad &\quad + Lm(x_n, x_{n+1}, x_{n+1}), \end{aligned}$$

318 where

$$\begin{aligned} 319 \quad M(x_n, x_{n+1}, x_{n+1}) &= \max\{G(Sx_n, Sx_{n+1}, Sx_{n+1}), G(Sx_n, Tx_n, Tx_n), \\ 320 \quad &\quad G(Sx_{n+1}, Tx_{n+1}, Tx_{n+1}), G(Sx_{n+1}, Tx_{n+1}, Tx_{n+1})\} \\ 321 \quad &= G(Sx_n, Sx_{n+1}, Sx_{n+1}) \end{aligned}$$

322 and

$$\begin{aligned} 323 \quad m(x_n, x_{n+1}, x_{n+1}) & \\ 324 \quad &= \min\{G(Tx_n, Sx_n, Sx_n), G(Tx_n, Sx_{n+1}, Sx_{n+1}), \\ 325 \quad &\quad G(Tx_n, Sx_{n+1}, Sx_{n+1}), G(Tx_n, Sx_{n+1}, Sx_{n+1})\} \\ 326 \quad &= \min\{G(y_n, y_{n-1}, y_{n-1}), G(y_n, y_n, y_n), G(y_n, y_n, y_n), G(y_n, y_n, y_n)\} \\ 327 \quad &= 0 \end{aligned}$$

328 since $G(y_n, y_n, y_n) = 0$. This implies that

$$329 \quad \psi(G(y_n, y_{n+1}, y_{n+1})) \leq \psi(G(y_{n-1}, y_n, y_n)) - \varphi(G(y_{n-1}, y_n, y_n)). \quad (19)$$

330 Hence, from the inequality (18), if $y_m = y_{m+1}$ for some m , then it follows that $y_n = y_m$
 331 for all $n \geq m$ so that $\{y_n\}$ is Cauchy. Therefore, without loss of generality, we assume
 332 that $y_n \neq y_{n+1}$ for all $n = 0, 1, 2, \dots$. Now, from (18), we have

$$333 \quad \psi(G(y_n, y_{n+1}, y_{n+1})) < \psi(G(y_{n-1}, y_n, y_n)).$$

334 Hence by the nondecreasing nature of ψ , it follows that

$$335 \quad G(y_n, y_{n+1}, y_{n+1}) \leq G(y_{n-1}, y_n, y_n) \text{ for all } n = 1, 2, \dots$$

336 Therefore $\{G(y_n, y_{n+1}, y_{n+1})\}$ is a monotone decreasing sequence of nonnegative reals.
 337 So, there exists $r \geq 0$ such that $G(y_n, y_{n+1}, y_{n+1}) \rightarrow r$ as $n \rightarrow \infty$. Now, from the
 338 inequality (19), we have

$$339 \quad \psi(G(y_n, y_{n+1}, y_{n+1})) \leq \psi(G(y_{n-1}, y_n, y_n)) - \varphi(G(y_{n-1}, y_n, y_n)).$$

340 On letting $n \rightarrow \infty$, we have $\psi(r) \leq \psi(r) - \varphi(r)$ so that $\varphi(r) \leq 0$. Since $\varphi(r) \geq 0$, it
 341 follows that $\varphi(r) = 0$ so that $r = 0$.

$$342 \quad \text{i.e., } G(y_n, y_{n+1}, y_{n+1}) \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (20)$$

343 We now prove that the sequence $\{y_n\}$ is Cauchy. If we suppose that $\{y_n\}$ is not
 344 Cauchy, then there exists an $\epsilon > 0$ and there exist subsequences $\{y_{n_k}\}, \{y_{m_k}\}$ of $\{y_n\}$
 345 with $n_k > m_k \geq k$ such that

$$346 \quad G(y_{n_k}, y_{m_k}, y_{m_k}) \geq \epsilon. \quad (21)$$

347 Corresponding to each m_k , we can choose n_k such that it is the smallest integer with
 348 $n_k > m_k$ and satisfying (21). Then, we have

$$349 \quad G(y_{n_k}, y_{m_k}, y_{m_k}) \geq \epsilon \text{ and } G(y_{n_k-1}, y_{m_k}, y_{m_k}) < \epsilon. \quad (22)$$

350 Now the following identities follow as in the proof of Theorem 4.

- 351 (i) $\lim_{k \rightarrow \infty} G(y_{n_k}, y_{m_k}, y_{m_k}) = \epsilon.$
- 352 (ii) $\lim_{k \rightarrow \infty} G(y_{n_k-1}, y_{m_k-1}, y_{m_k-1}) = \epsilon.$
- 353 (iii) $\lim_{k \rightarrow \infty} G(y_{n_k}, y_{m_k-1}, y_{m_k-1}) = \epsilon.$

354 We now consider

$$\begin{aligned} 355 \quad \psi(G(y_{n_k}, y_{m_k}, y_{m_k})) &= \psi(G(Tx_{n_k}, Tx_{m_k}, Tx_{m_k})) \\ 356 \quad &\leq \psi(M(x_{n_k}, x_{m_k}, x_{m_k})) - \varphi(M(x_{n_k}, x_{m_k}, x_{m_k})) \\ 357 \quad &\quad + Lm(x_{n_k}, x_{m_k}, x_{m_k}), \end{aligned}$$

358 where

$$\begin{aligned} 359 \quad M(x_{n_k}, x_{m_k}, x_{m_k}) &= \max\{G(Sx_{n_k}, Sx_{m_k}, Sx_{m_k}), G(Sx_{n_k}, Tx_{n_k}, Tx_{n_k}), \\ 360 \quad &\quad G(Sx_{m_k}, Tx_{m_k}, Tx_{m_k}), G(Sx_{m_k}, Tx_{m_k}, Tx_{m_k})\} \\ 361 \quad &= G(Sx_{n_k}, Sx_{m_k}, Sx_{m_k}) = G(y_{n_k-1}, y_{m_k-1}, y_{m_k-1}) \end{aligned}$$

362 and

$$\begin{aligned} 363 \quad m(x_{n_k}, x_{m_k}, x_{m_k}) &= \min\{G(Tx_{n_k}, Sx_{n_k}, Sx_{n_k}), G(Tx_{n_k}, Sx_{m_k}, Sx_{m_k}), \\ 364 \quad &\quad G(Tx_{n_k}, Sx_{m_k}, Sx_{m_k}), G(Tx_{n_k}, Sx_{m_k}, Sx_{m_k})\}. \end{aligned}$$

365 Therefore,

$$\begin{aligned} 366 \quad &\psi(G(y_{n_k}, y_{m_k}, y_{m_k})) \\ 367 \quad &\leq \psi(G(y_{n_k-1}, y_{m_k-1}, y_{m_k-1})) - \varphi(G(y_{n_k-1}, y_{m_k-1}, y_{m_k-1})) \\ 368 \quad &\quad + L \min\{G(y_{n_k}, y_{n_k-1}, y_{n_k-1}), G(y_{n_k}, y_{m_k-1}, y_{m_k-1}), \\ 369 \quad &\quad G(y_{n_k}, y_{m_k-1}, y_{m_k-1}), G(y_{n_k}, y_{m_k-1}, y_{m_k-1})\}. \end{aligned}$$

370 On letting $k \rightarrow \infty$ and using (i)–(iii) and (20), we get

$$371 \quad \psi(\epsilon) \leq \psi(\epsilon) - \varphi(\epsilon) < \psi(\epsilon),$$

372 which is a contradiction. Therefore $\{y_n\}$ is a G -Cauchy sequence. Since X is complete,
 373 there exists $z \in X$ such that $\lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} Tx_n = \lim_{n \rightarrow \infty} Sx_{n+1} = z$. We

now prove that z is a common fixed point of T and S . Since S is continuous, we have $\lim_{n \rightarrow \infty} STx_n = \lim_{n \rightarrow \infty} SSx_n = Sz$. Further, since S and T are weakly compatible, we have $\lim_{n \rightarrow \infty} G(TSx_n, STx_n, STx_n) = 0$, which implies $\lim_{n \rightarrow \infty} TSx_n = Sz$. Now, from (18), we have

$$\begin{aligned} \psi(G(TSx_n, Tx_n, Tx_n)) &\leq \psi(M(Sx_n, x_n, x_n)) - \varphi(M(Sx_n, x_n, x_n)) \\ &\quad + Lm(Sx_n, x_n, x_n), \end{aligned} \quad (23)$$

where

$$\begin{aligned} M(Sx_n, x_n, x_n) &= \max\{G(SSx_n, Sx_n, Sx_n), G(SSx_n, TSx_n, TSx_n), \\ &\quad G(Sx_n, Tx_n, Tx_n), G(Sx_n, Tx_n, Tx_n)\} \\ &= G(SSx_n, Sx_n, Sx_n) \end{aligned}$$

and

$$\begin{aligned} m(Sx_n, x_n, x_n) &= \min\{G(TSx_n, SSx_n, SSx_n), G(TSx_n, Sx_n, Sx_n), \\ &\quad G(TSx_n, Sx_n, Sx_n), G(TSx_n, Sx_n, Sx_n)\}. \end{aligned}$$

Therefore, from (23), we have

$$\begin{aligned} \psi(G(TSx_n, Tx_n, Tx_n)) &\leq \psi(G(SSx_n, Sx_n, Sx_n)) - \varphi(G(SSx_n, Sx_n, Sx_n)) \\ &\quad + L \min\{G(TSx_n, SSx_n, SSx_n), G(TSx_n, Sx_n, Sx_n), \\ &\quad G(TSx_n, Sx_n, Sx_n), G(TSx_n, Sx_n, Sx_n)\}. \end{aligned}$$

On letting $n \rightarrow \infty$, we get

$$\psi(G(Sz, z, z)) \leq \psi(G(Sz, z, z)) - \varphi(G(Sz, z, z)),$$

which implies that $\varphi(G(Sz, z, z)) \leq 0$ so that we must have $Sz = z$. Now, we consider

$$\psi(G(Tx_n, Tz, Tz)) \leq \psi(M(x_n, z, z)) - \varphi(M(x_n, z, z)) + Lm(x_n, z, z), \quad (24)$$

where

$$M(x_n, z, z) = \max\{G(Sx_n, Sz, Sz), G(Sx_n, Tx_n, Tx_n), G(Sz, Tz, Tz), G(Sz, Tz, Tz)\}$$

and

$$m(x_n, z, z) = \min\{G(Tx_n, Sx_n, Sx_n), G(Tx_n, Sz, Sz), G(Tx_n, Sz, Sz), G(Tx_n, Sz, Sz)\}.$$

Also, we have

$$\lim_{n \rightarrow \infty} M(x_n, z, z) = G(z, Tz, Tz) \text{ and } \lim_{n \rightarrow \infty} m(x_n, z, z) = 0, \quad (25)$$

since $\lim_{n \rightarrow \infty} G(TSx_n, SSx_n, SSx_n) = 0$. Now, on letting $n \rightarrow \infty$ in (24) and using (25), we get

$$\psi(G(z, Tz, Tz)) \leq \psi(G(z, Tz, Tz)) - \varphi(G(z, Tz, Tz)).$$

Then $\varphi(G(z, Tz, Tz)) \leq 0$. Therefore, $\varphi(G(z, Tz, Tz)) = 0$ so that $G(z, Tz, Tz) = 0$. Therefore, $Tz = z$. Thus z is a common fixed point of T and S . Uniqueness of common fixed point of T and S follows from the inequality (18). This completes the proof of the theorem.

408 3 Corollaries and Examples

409 In this section, we draw some corollaries from the main results of Section 2 and provide
 410 examples in support of our results. The following is an example in support of Theorem
 411 4.

412 EXAMPLE 6. Let $X = [0, 1]$. We define $G : X^3 \rightarrow \mathbb{R}_+$ by

$$413 G(x, y, z) = \begin{cases} 0, & \text{if } x = y = z \\ \max\{x, y, z\}, & \text{otherwise.} \end{cases}$$

414 Then (X, G) is a complete G -metric space. We define $T : X \rightarrow X$ by

$$415 T(x) = \begin{cases} \frac{1}{2} & \text{if } x = 0, \\ 2x & \text{if } 0 < x < \frac{1}{2}, \\ 1 & \text{if } \frac{1}{2} \leq x \leq 1. \end{cases}$$

416 We define ψ and φ on \mathbb{R}_+ by $\psi(t) = \frac{t^2}{3}$ and $\varphi(t) = \frac{t^2}{2}$. Then, it is easy to verify that
 417 T satisfies the inequality (4) with $L = 1$. i.e., T is a (ψ, φ) -almost weakly contractive
 418 map. Thus T satisfies all the hypothesis of Theorem 4 and 1 is the unique fixed point
 419 of T . Here we observe that T is not a continuous map.

420 Further, we observe that T is not a weakly contractive map. For, let us choose $x =$
 421 $\frac{1}{3}$ and $y = z = 0$. Then $G(Tx, Ty, Tz) = \frac{2}{3}$, $G(x, y, z) = \frac{1}{3}$ and $\varphi(G(x, y, z)) = \varphi(\frac{1}{3})$.
 422 Hence,

$$423 \frac{2}{3} = G(Tx, Ty, Tz) \not\leq G(x, y, z) - \varphi(G(x, y, z)) = \frac{1}{3} - \varphi(\frac{1}{3}) \text{ for any } \varphi \in \Psi.$$

424 Hence T does not satisfy the inequality (2) for any $\varphi \in \Psi$ so that T is not a weakly
 425 contractive map in G -metric space. Thus Theorem 2 is not applicable.

426 Further, this example suggests that the class of (ψ, φ) -almost weakly contractive
 427 maps is larger than the class of weakly contractive maps in G -metric spaces.

428 REMARK 1. Theorem 2 follows as a corollary to Theorem 4 by choosing ψ as the
 429 identity map and $L = 0$. Hence Example 6 suggests that Theorem 4 is a generalization
 430 of Theorem 2.

431 COROLLARY 1. Let (X, G) be a complete G -metric space and let T and S be
 432 two selfmaps on (X, G) . Assume that $T(X) \subseteq S(X)$, S is continuous, and there exist
 433 $\psi, \varphi \in \Psi$ such that

$$434 \psi(G(Tx, Ty, Tz)) \leq \psi(M(x, y, z)) - \varphi(M(x, y, z)), \quad (26)$$

435 where

$$436 M(x, y, z) = \max\{G(Sx, Sy, Sz), G(Sx, Tx, Tx), G(Sy, Ty, Ty), G(Sz, Tz, Tz)\}$$

437 for $x, y, z \in X$. Then T and S have a unique common fixed point in X provided T and
 438 S are weakly compatible maps.

439 PROOF. Follows from Theorem 5 by choosing $L = 0$.

440 The following is an example in support of Corollary 1.

441 EXAMPLE 7. Let $X = [0, 1]$. We define $G : X^3 \rightarrow \mathbb{R}_+$ by

442
$$G(x, y, z) = \begin{cases} 0, & \text{if } x = y = z, \\ \max\{x, y, z\}, & \text{otherwise.} \end{cases}$$

443 Then (X, G) is a complete G -metric space. We define $T, S : X \rightarrow X$ and ψ, φ on \mathbb{R}_+ by

444
$$T(x) = \frac{x^2}{2}, S(x) = \frac{x}{4}(5 - x), \psi(t) = \frac{4t^2}{3} \text{ and } \varphi(t) = \frac{t^2}{3}$$

445 for all $x \in X$ and $t \in \mathbb{R}_+$. Then clearly $T(X) \subseteq S(X)$. Without loss of generality, we
446 assume that $x > y > z$. Then

447
$$G(Tx, Ty, Tz) = \max\{Tx, Ty, Tz\} = \max\{\frac{x^2}{2}, \frac{y^2}{2}, \frac{z^2}{2}\} = \frac{x^2}{2}.$$

448 Also

449
$$G(Sx, Sy, Sz) = \max\{Sx, Sy, Sz\} = \max\{\frac{x}{4}(5 - x), \frac{y}{4}(5 - y), \frac{z}{4}(5 - z)\} = \frac{x}{4}(5 - x).$$

450 Now, we consider

451
$$\begin{aligned} \psi(G(Tx, Ty, Tz)) &= \frac{x^4}{3} \leq \frac{x^2}{16}(5 - x)^2 = [G(Sx, Sy, Sz)]^2 \\ 452 &\leq [M(x, y, z)]^2 = \frac{4}{3}[M(x, y, z)]^2 - \frac{1}{3}[M(x, y, z)]^2 \\ 453 &= \psi(M(x, y, z)) - \varphi(M(x, y, z)). \end{aligned}$$

454 Therefore,

455
$$\psi(G(Tx, Ty, Tz)) \leq \psi(M(x, y, z)) - \varphi(M(x, y, z))$$

456 so that the inequality (26) of Corollary 1 holds. Thus, T and S satisfy all the hypotheses
457 of Corollary 1 and 0 is the unique common fixed point of T and S .

458 REMARK 2. We observe that the φ that is used in Theorem 5 is different from ϕ
459 that is used in Theorem 3.

460 **Acknowledgment.** The authors are sincerely thankful to the anonymous referee
461 for his/her valuable suggestions which helped us to improve the quality of the paper.

462 References

463 [1] C. T. Aage and J. N. Salunke, Fixed points for weak contractions in G -metric
464 spaces, Appl. Math. E-Notes, 12(2012), 23–28.

465 [2] M. Abbas, G. V. R. Babu and G. N. Alemayehu, On common fixed points of weakly
466 compatible mappings satisfying ‘Generalized condition (B)’, Filomat 25(2011), 9–
467 19.

468 [3] Ya. I. Alber and Guerre-Delabriere, Principles of weakly contractive maps in Hilbert
469 spaces, New results in Operator Theory and its applications (I. Gohberg, and Yu.
470 Lyubich, Eds.), Oper. Theory Adv. Appl., 98(1997), 7–22.

471 [4] G. V. R. Babu, K. Nageswara Rao and G. N. Alemayehu, Common fixed points of
472 two pairs of generalized weakly contractive maps, Adv. Studies in Contemporary
473 Math., 20(2010), 575–594.

474 [5] G. V. R. Babu and P. D. Sailaja, A fixed point theorem of generalized weakly
475 contractive maps in orbitally complete metric spaces, Thai J. Math., 9(2011),
476 1–10.

477 [6] G. V. R. Babu, M. L. Sandhya and M. V. R. Kameswari, A note on a fixed point
478 theorem of Berinde on weak contractions, Carpath. J. Math., 24(2008), 8–12.

479 [7] V. Berinde, Approximating fixed points of weak contractions using the Picard
480 iteration, Nonlinear Anal. Forum, 9(2004), 43–53.

481 [8] V. Berinde, General constructive fixed point theorems for Cirić-type almost con-
482 tractions in metric spaces, Carpath. J. Math., 24(2008), 10–19.

483 [9] B. S. Choudhury, P. Konar, B. E. Rhoades and N. Metiya, Fixed point therorems
484 for generalized weakly contractive mappings, Nonlinear Anal. Theory, Methods
485 and Appl., 74(2011), 2116–2126.

486 [10] B. S. Choudhury, Sanjay Kumar, Asha and Krishnapada Das, Some fixed point
487 theorems in G -metric spaces, Math. Sci. Lett., 1(2012), 25–31.

488 [11] D. Doric, Common fixed point for generalized (ψ, φ) -weak contractions, Appl.
489 Math. Lett., 22(2009), 1896–1900.

490 [12] P. N. Dutta and B. S. Choudhury, A generalisation of contraction principle in
491 metric spaces, Fixed Point Theory and Appl., (2008), Article ID 406368, 8 pages.

492 [13] G. Jungck and B. E. Rhoades, Fixed points for set valued functions without
493 continuity, Indian J. Pure and Appl. Math., 29(1998), 227–238.

494 [14] Z. Mustafa, H. Obiedat and F. Awawdeh, Some fixed point theorem for mapping
495 on complete G -metric spaces, Fixed Point Theory and Appl., (2008) Article ID
496 189870.

497 [15] Z. Mustafa and B. Sims, A new approach to generalized metric spaces, J. Nonlinear
498 and Convex Anal., 7(2006), 289–297.

499 [16] B. E. Rhoades, Some theorems on weakly contractive maps, Nonlinear Anal.,
500 47(2001), 2683–2693.

501 [17] K. P. R. Sastry, G. V. R. Babu and K. T. Kidane, A common fixed point of gen-
502 eralized (ψ, ϕ) -weakly contractive maps where ϕ is nondecreasing (not necessarily
503 continuous or lower semicontinuous), Advances in Fixed Point Theory, 2(2012),
504 203–223.

505 [18] W. Shatanawi, Common fixed point result for two self-maps in G -metric spaces,
506 Mat. Vesnik, 65(2013), 143–150.

507 [19] Q. Zhang and Y. Song, Fixed point theory for generalized φ -weak contractions,
508 Appl. Math. Lett., 22(2009), 75–78.