

ASYMPTOTIC PROPERTIES OF MILD SOLUTIONS OF NONAUTONOMOUS EVOLUTION EQUATIONS WITH APPLICATIONS TO RETARDED DIFFERENTIAL EQUATIONS

GABRIELE GÜHRING AND FRANK RÄBIGER

Received 18 May 1999

We investigate the asymptotic properties of the inhomogeneous nonautonomous evolution equation $(d/dt)u(t) = Au(t) + B(t)u(t) + f(t)$, $t \in \mathbb{R}$, where $(A, D(A))$ is a Hille-Yosida operator on a Banach space X , $B(t)$, $t \in \mathbb{R}$, is a family of operators in $\mathcal{L}(\overline{D(A)}, X)$ satisfying certain boundedness and measurability conditions and $f \in L^1_{\text{loc}}(\mathbb{R}, X)$. The solutions of the corresponding homogeneous equations are represented by an evolution family $(U_B(t, s))_{t \geq s}$. For various function spaces \mathcal{F} we show conditions on $(U_B(t, s))_{t \geq s}$ and f which ensure the existence of a unique solution contained in \mathcal{F} . In particular, if $(U_B(t, s))_{t \geq s}$ is p -periodic there exists a unique bounded solution u subject to certain spectral assumptions on $U_B(p, 0)$, f and u . We apply the results to nonautonomous semilinear retarded differential equations. For certain p -periodic retarded differential equations we derive a characteristic equation which is used to determine the spectrum of $(U_B(t, s))_{t \geq s}$.

1. Introduction

Consider the inhomogeneous nonautonomous evolution equation

$$\frac{d}{dt}u(t) = A(t)u(t) + f(t), \quad t \in \mathbb{R}, \quad (1.1)$$

where $A(t)$, $t \in \mathbb{R}$, are (unbounded, linear) operators on a Banach space X and $f \in L^1_{\text{loc}}(\mathbb{R}, X)$. Assume that the homogeneous equation

$$\frac{d}{dt}u(t) = A(t)u(t), \quad t \in \mathbb{R}, \quad (1.2)$$

is well posed in the sense that the solutions of (1.2) define a uniquely determined evolution family $(U(t, s))_{t \geq s}$ of bounded operators on X . In that case solutions $u : \mathbb{R} \rightarrow X$ of the integral equation

$$u(t) = U(t, s)u(s) + \int_s^t U(t, \sigma)f(\sigma)d\sigma, \quad t \geq s, \quad (1.3)$$

Copyright © 1999 Hindawi Publishing Corporation

Abstract and Applied Analysis 4:3 (1999) 169–194

1991 Mathematics Subject Classification: 34C25, 34C27, 34C28, 34G10, 47D06, 47H15

URL: <http://aaa.hindawi.com/volume-4/S1085337599000214.html>

can be interpreted as mild solutions of (1.1). It has been shown in [5, 22, 23] that for each $f \in C_b(\mathbb{R}, X)$ (respectively, $C_0(\mathbb{R}, X)$) equation (1.1) has a *unique* mild solution $u \in C_b(\mathbb{R}, X)$ (respectively, $C_0(\mathbb{R}, X)$) if and only if the evolution family $(U(t, s))_{t \geq s}$ has an exponential dichotomy (see also [12, 24] when the operators $A(t)$, $t \in \mathbb{R}$, are bounded). For a detailed account of the numerous other results in this direction we refer to [7, 22].

Now assume that (1.2) is p -periodic, that is, $A(t + p) = A(t)$, $t \in \mathbb{R}$. It has been shown in [6, 19, 27, 30] that under a certain spectral condition (*nonresonance condition*) on the monodromy operator $U(p, 0)$ and the inhomogeneity f there is a p -periodic (respectively, almost periodic) mild solution of (1.1) provided that f has the same property. Moreover, u is unique subject to certain spectral assumptions. If $(U(t, s))_{t \geq s}$ has an exponential dichotomy, then the nonresonance condition is always satisfied and we obtain existence and uniqueness of a p -periodic (respectively, almost periodic) mild solution of (1.1) for every p -periodic (respectively, almost periodic) inhomogeneity f . We point out that in [10, 31] related results are discussed for Volterra equations (see also [32]).

In the present paper, we study the modified equation

$$\frac{d}{dt}u(t) = (A + B(t))u(t) + f(t), \quad t \in \mathbb{R}, \quad (1.4)$$

where $(A, D(A))$ is a Hille-Yosida operator on the Banach space X , $B(t)$, $t \in \mathbb{R}$, is a family of operators in $\mathcal{L}(\overline{D(A)}, X)$, and $f \in L^1_{\text{loc}}(\mathbb{R}, X)$. We stress that, in general, $X_0 = \overline{D(A)}$ is a proper subspace of X from which the main difficulties arise. Our approach is based on the theory of extrapolation spaces associated with the operator A (see [Section 2](#) and [26]). In particular, it is used in our definition of mild solutions of (2.3). Moreover, it allows to show that under a certain boundedness and measurability condition on the family $B(t)$, $t \in \mathbb{R}$, there is a (unique) evolution family $(U_B(t, s))_{t \geq s}$ on X_0 associated with the homogeneous equation

$$\frac{d}{dt}u(t) = (A + B(t))u(t), \quad t \in \mathbb{R}, \quad (1.5)$$

(cf. [9, 33]). The evolution family $(U_B(t, s))_{t \geq s}$ is used to derive another representation of the mild solutions of (2.3) (see [Theorem 2.2](#)). This representation is crucial for the investigations in [Section 3](#). There we extend the above-mentioned results on the existence and uniqueness of mild solutions of (1.1) satisfying a particular asymptotic behavior to mild solutions of (2.3). We point out that in the autonomous case, that is, $B(t) = B$, similar results are obtained in [2]. In [Section 4](#), we discuss asymptotic properties of mild solutions of the semilinear nonautonomous equation

$$\frac{d}{dt}u(t) = (A + B(t))u(t) + F(t, u(t)), \quad t \in \mathbb{R}, \quad (1.6)$$

where the nonlinearity $F : \mathbb{R} \times X_0 \rightarrow X$ satisfies a standard Lipschitz condition. In [Section 5](#), the advantage of our approach becomes visible when we study inhomogeneous nonautonomous retarded differential equations

$$\frac{d}{dt}w(t) = Cw(t) + K(t)w_t + h(t), \quad t \in \mathbb{R}, \quad (1.7)$$

on a Banach space Y . A standard procedure (cf. [16, 33, 39]) allows to transform (5.1) into an equation of the form (2.3) on a different Banach space. Now the results of Section 3 can be applied to investigate asymptotic properties of mild solutions of (5.1). For a special periodic retarded differential equation we derive a characteristic equation which makes it easier to verify the spectral conditions in our results (see Theorem 5.9). Finally, we point out that in finite dimensions asymptotic properties of solutions of inhomogeneous retarded differential equations have been studied in [37] under the assumption that the corresponding homogeneous equation admits an exponential dichotomy (see also [17, Section 6.6.2], [29]).

2. Mild solutions and extrapolation spaces

We first recall some properties of Hille-Yosida operators and extrapolation spaces. For more details we refer to [26] and the references therein. Throughout the whole paper X denotes a Banach space and $(A, D(A))$ is a *Hille-Yosida operator* on X , that is, A is linear and the *resolvent set* $\rho(A)$ of A contains a half-line (ω, ∞) such that

$$M = \sup \{ \|(\lambda - \omega)^n R(\lambda, A)^n\| : \lambda > \omega; n \in \mathbb{N} \} < \infty, \quad (2.1)$$

where $R(\lambda, A) = (\lambda - A)^{-1}$ is the resolvent of A at λ . It is well known that the part A_0 of A in $X_0 = \overline{D(A)}$ generates a C_0 -semigroup $(T_0(t))_{t \geq 0}$ on X_0 and that $\|T_0(t)\| \leq M e^{\omega t}$, $t \geq 0$. For $\lambda \in \rho(A_0)$ the resolvent $R(\lambda, A_0)$ is the restriction of $R(\lambda, A)$ to X_0 .

Typical examples of Hille-Yosida operators appearing in partial differential equations can be found, for example, in [11], see also Section 5. On X_0 we introduce the norm $\|x\|_{-1} = \|R(\lambda_0, A_0)x\|$, where $\lambda_0 \in \rho(A)$ is fixed. A different choice of $\lambda_0 \in \rho(A)$ leads to an equivalent norm. The completion X_{-1} of X_0 with respect to $\|\cdot\|_{-1}$ is called the *extrapolation space* of X_0 with respect to A . The *extrapolated semigroup* $(T_{-1}(t))_{t \geq 0}$ consists of the unique continuous extensions $T_{-1}(t)$ of the operators $T_0(t)$, $t \geq 0$, to X_{-1} . The semigroup $(T_{-1}(t))_{t \geq 0}$ is strongly continuous and its generator A_{-1} is the unique continuous extension of A_0 to $\mathcal{L}(X_0, X_{-1})$. Moreover, X is continuously embedded in X_{-1} and $R(\lambda, A_{-1})$ is the unique continuous extension of $R(\lambda, A)$ to X_{-1} for $\lambda \in \rho(A)$. Finally, A_0 and A are the parts of A_{-1} in X_0 and X , respectively. It follows from [26, Proposition 3.3], that for $f \in L^1_{\text{loc}}(\mathbb{R}, X)$ and $t \geq s$

$$\begin{aligned} & \int_s^t T_{-1}(t-\sigma) f(\sigma) d\sigma \in X_0, \\ & (t, s) \mapsto \int_s^t T_{-1}(t-\sigma) f(\sigma) d\sigma \text{ is continuous,} \\ & \left\| \int_s^t T_{-1}(t-\sigma) f(\sigma) d\sigma \right\| \leq M_1 \int_s^t e^{\omega(t-\sigma)} \|f(\sigma)\| d\sigma \quad \text{for a constant } M_1 \geq 1. \end{aligned} \quad (2.2)$$

We consider the inhomogeneous nonautonomous evolution equation

$$\frac{d}{dt} u(t) = (A + B(t))u(t) + f(t), \quad t \in \mathbb{R}, \quad f \in L^1_{\text{loc}}(\mathbb{R}, X), \quad (2.3)$$

where $(A, D(A))$ is a Hille-Yosida operator on the Banach space X and $B(t) \in \mathcal{L}(X_0, X)$, $t \in \mathbb{R}$, is a family of operators such that $t \mapsto B(t)x$ is strongly measurable for every $x \in X_0$ and $\|B(\cdot)\| \leq b(\cdot)$ for a function $b \in L^1_{\text{loc}}(\mathbb{R})$. For our purposes the notion of a mild solution of (2.3) is most useful. We point out that our definition of a mild solution coincides with that given in [8], the F -solutions in [11], the weak solutions in [13] and the integral solutions in [39].

Definition 2.1. If $f \in L^1_{\text{loc}}(\mathbb{R}, X)$ and $T \geq s$, then $u = u(\cdot, f) \in C([s, T], X_0)$ is called a *mild solution* of (2.3) on $[s, T]$ if

$$u(t) = T_0(t-s)u(s) + \int_s^t T_{-1}(t-\sigma)(B(\sigma)u(\sigma) + f(\sigma))d\sigma \quad \text{for } t \in [s, T]. \quad (2.4)$$

A function $u = u(\cdot, f) \in C(\mathbb{R}, X_0)$ that satisfies (2.4) for all $t \geq s$ in \mathbb{R} is called a *mild solution on \mathbb{R}* of (2.3).

Under our assumptions on A and $B(t)$, $t \in \mathbb{R}$, it follows that for $f \in L^1_{\text{loc}}(\mathbb{R}, X)$ and $s \in \mathbb{R}$ there is a unique mild solution $u = u(\cdot, f, s, x) \in C([s, \infty), X_0)$ of

$$\frac{d}{dt}u(t) = (A + B(t))u(t) + f(t), \quad t \geq s, \quad u(s) = x \in X_0, \quad (2.5)$$

(cf. [15] or Theorem 2.2). Mild solutions of the homogeneous equation

$$\frac{d}{dt}v(t) = (A + B(t))v(t), \quad t \in \mathbb{R}, \quad (2.6)$$

have another representation. For that we need the following notion. A family $(U(t, s))_{t \geq s}$ in $\mathcal{L}(X_0)$ is called an *evolution family* on X_0 if $U(t, t) = Id$, $U(t, r)U(r, s) = U(t, s)$ for $t \geq r \geq s$ and $(t, s) \mapsto U(t, s)x$ is continuous for $t \geq s$ and $x \in X_0$. It is known (cf. [9, Theorem 2.3], [33, Theorem 2.3], where a slightly more special situation is considered) that there exists a unique evolution family $(U_B(t, s))_{t \geq s}$ on X_0 that satisfies the variation-of-parameters formula

$$U_B(t, s)x = T_0(t-s)x + \int_s^t T_{-1}(t-\sigma)B(\sigma)U_B(\sigma, s)x d\sigma, \quad t \geq s, x \in X_0. \quad (2.7)$$

Thus $t \mapsto U_B(t, s)x$ is the unique mild solution on $[s, \infty)$ of the initial value problem

$$\frac{d}{dt}u(t) = (A + B(t))u(t), \quad t \geq s, \quad u(s) = x \in X_0. \quad (2.8)$$

Gronwall's inequality (cf. [1, Corollary II.6.2]), the estimate in (2.2), and (2.7) imply

$$\|U_B(t, s)\| \leq M e^{\omega(t-s) + M_1 \int_s^t b(\sigma) d\sigma}, \quad t \geq s, \quad (2.9)$$

for the constants $M, M_1 \geq 1$. In particular, if $\|B(\cdot)\|$ is bounded from above by a function $b \in L^1_{\text{loc}, u}(\mathbb{R})$, that is, $\|b\|_{1, \text{loc}, u} = \sup_{t \in \mathbb{R}} \int_{t-1}^t |b(\sigma)| d\sigma < \infty$, then the evolution family $(U_B(t, s))_{t \geq s}$ is *exponentially bounded*, that is, $\|U_B(t, s)\| \leq N e^{\beta(t-s)}$ for $t \geq s$ and constants $N \geq 1$, $\beta \in \mathbb{R}$. In the following result we give a representation of mild solutions of (2.3) in terms of the evolution family $(U_B(t, s))_{t \geq s}$. A special case has been discussed in [16, Theorem 3.6].

THEOREM 2.2. Let $f \in L^1_{\text{loc}}(\mathbb{R}, X)$, $s \in \mathbb{R}$, and $x \in X_0$. Then there is a unique mild solution $u \in C([s, \infty), X_0)$ of (2.5) given by

$$u(t) = U_B(t, s)x + \lim_{\lambda \rightarrow \infty} \int_s^t U_B(t, \sigma)\lambda R(\lambda, A)f(\sigma)d\sigma \quad \text{for } t \geq s. \quad (2.10)$$

Moreover, $\lim_{\lambda \rightarrow \infty} \int_s^t U_B(t, \sigma)\lambda R(\lambda, A)f(\sigma)d\sigma \in X_0$ exists uniformly for $t \geq s$ in compact sets in \mathbb{R} .

Proof. Let $\lambda > \omega$ and set

$$w_\lambda(t, s) = \int_s^t U_B(t, \sigma)\lambda R(\lambda, A)f(\sigma)d\sigma, \quad t \geq s. \quad (2.11)$$

Then (2.7) leads to

$$\begin{aligned} w_\lambda(t, s) &= \int_s^t T_0(t - \sigma)\lambda R(\lambda, A)f(\sigma)d\sigma \\ &\quad + \int_s^t \left(\int_\sigma^t T_{-1}(t - \tau)B(\tau)U_B(\tau, \sigma)\lambda R(\lambda, A)f(\sigma)d\tau \right) d\sigma \\ &= \lambda R(\lambda, A_0) \int_s^t T_{-1}(t - \sigma)f(\sigma)d\sigma \\ &\quad + \int_s^t \left(\int_s^\tau T_{-1}(t - \tau)B(\tau)U_B(\tau, \sigma)\lambda R(\lambda, A)f(\sigma)d\sigma \right) d\tau \\ &= \lambda R(\lambda, A_0) \int_s^t T_{-1}(t - \sigma)f(\sigma)d\sigma + \int_s^t T_{-1}(t - \sigma)B(\sigma)w_\lambda(\sigma, s)d\sigma, \quad t \geq s. \end{aligned} \quad (2.12)$$

If $z(t, s) = \int_s^t T_{-1}(t - \sigma)f(\sigma)d\sigma$, $t \geq s$, then by (2.2) for $\lambda, \mu > \omega$

$$\begin{aligned} \|w_\lambda(t, s) - w_\mu(t, s)\| &\leq \|(\lambda R(\lambda, A_0) - \mu R(\mu, A_0))z(t, s)\| \\ &\quad + M_1 \int_s^t e^{\omega(t-\sigma)}b(\sigma) \|w_\lambda(\sigma, s) - w_\mu(\sigma, s)\| d\sigma. \end{aligned} \quad (2.13)$$

From (2.2) it follows that z is a continuous mapping into X_0 . Hence

$$\lim_{\lambda, \mu \rightarrow \infty} \|(\lambda R(\lambda, A_0) - \mu R(\mu, A_0))z(t, s)\| = 0 \quad (2.14)$$

uniformly for $t \geq s$ in compact intervals. Thus if $\epsilon > 0$ and $I \subseteq \mathbb{R}$ is a compact interval, then by (2.13) there is a constant \tilde{M} depending only on the length of I such that

$$\|w_\lambda(t, s) - w_\mu(t, s)\| \leq \epsilon + \tilde{M} \int_s^t b(\sigma) \|w_\lambda(\sigma, s) - w_\mu(\sigma, s)\| d\sigma \quad (2.15)$$

for $t \geq s$ in I and $\lambda, \mu > \omega$ sufficiently large. An application of Gronwall's inequality (see [1, Corollary II.6.2]) leads to the estimate

$$\|w_\lambda(t, s) - w_\mu(t, s)\| \leq \epsilon e^{\tilde{M} \int_s^t b(\sigma)d\sigma} \quad (2.16)$$

for $t \geq s$ in I and $\lambda, \mu > \omega$ sufficiently large. Hence $w(t, s) = \lim_{\lambda \rightarrow \infty} w_\lambda(t, s)$ exists uniformly for $t \geq s$ in compact intervals.

Since A is a Hille-Yosida operator it follows from the definition of w_λ that $\sup\{\|w_\lambda(t, s)\| : \lambda > \omega + 1; t \geq s \text{ in } I\} < \infty$. Hence, by (2.12) and Lebesgue's dominated convergence theorem, we have

$$w(t, s) = \int_s^t T_{-1}(t-\sigma) B(\sigma) w(\sigma, s) d\sigma + \int_s^t T_{-1}(t-\sigma) f(\sigma) d\sigma, \quad t \geq s. \quad (2.17)$$

Now consider the function

$$u(t) = U_B(t, s)x + \lim_{\lambda \rightarrow \infty} \int_s^t U_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma = U_B(t, s)x + w(t, s), \quad t \geq s. \quad (2.18)$$

By (2.17) and (2.7), we obtain

$$\begin{aligned} u(t) &= U_B(t, s)x + \int_s^t T_{-1}(t-\sigma) B(\sigma) w(\sigma, s) d\sigma + \int_s^t T_{-1}(t-\sigma) f(\sigma) d\sigma \\ &= T_0(t-s)x + \int_s^t T_{-1}(t-\sigma) B(\sigma) (U_B(\sigma, s)x + w(\sigma, s)) d\sigma \\ &\quad + \int_s^t T_{-1}(t-\sigma) f(\sigma) d\sigma \\ &= T_0(t-s)x + \int_s^t T_{-1}(t-\sigma) (B(\sigma)u(\sigma) + f(\sigma)) d\sigma. \end{aligned} \quad (2.19)$$

Hence u is a mild solution of (2.3).

If $\tilde{u} \in C([s, \infty), X_0)$ is another mild solution of (2.5) we obtain

$$u(t) - \tilde{u}(t) = \int_s^t T_{-1}(t-\sigma) B(\sigma) (u(\sigma) - \tilde{u}(\sigma)) d\sigma, \quad t \geq s, \quad (2.20)$$

and an application of Gronwall's inequality yields $u = \tilde{u}$. \square

Remark 2.3. If in [Theorem 2.2](#) we assume that $f \in L^1_{\text{loc}}(\mathbb{R}, X_0)$, then the function $u \in C([s, \infty], X_0)$ is a mild solution of (2.5) if and only if

$$u(t) = U_B(t, s)x + \int_s^t U_B(t, \sigma) f(\sigma) d\sigma, \quad t \geq s. \quad (2.21)$$

[Theorem 2.2](#) has the following immediate consequence.

COROLLARY 2.4. *If $f \in L^1_{\text{loc}}(\mathbb{R}, X)$, then $u \in C(\mathbb{R}, X_0)$ is a mild solution of (2.3) if and only if*

$$u(t) = U_B(t, s)u(s) + \lim_{\lambda \rightarrow \infty} \int_s^t U_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma \quad \text{for } t \geq s. \quad (2.22)$$

In our next result we improve the convergence of the integrals considered in [Theorem 2.2](#). By $\text{BUC}_r(\mathbb{R}, X)$ we denote the space of bounded, uniformly continuous functions f from \mathbb{R} into X such that f has relatively compact range.

PROPOSITION 2.5. *Let $\|B(\cdot)\| \leq b(\cdot)$ for some $b \in L^1_{\text{loc},u}(\mathbb{R})$ and let $f \in \text{BUC}_r(\mathbb{R}, X)$. Then, for fixed $s > 0$, the limit*

$$\lim_{\lambda \rightarrow \infty} \int_{t-s}^t U_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma \quad (2.23)$$

exists uniformly for t in \mathbb{R} .

Proof. We claim that the function

$$\psi : \mathbb{R} \longrightarrow X_0 : t \longmapsto z(t, t-s) = \int_{t-s}^t T_{-1}(t-\sigma) f(\sigma) d\sigma = \int_0^s T_{-1}(\sigma) f(t-\sigma) d\sigma \quad (2.24)$$

has relatively compact range. In fact, fix $\epsilon > 0$. There exists $\delta = s/n > 0$ for an $n \in \mathbb{N}$ and a function $g : \mathbb{R} \rightarrow X$ such that g is constant on each interval $[k\delta, (k+1)\delta], k \in \mathbb{Z}$, the range of g is contained in a finite set $K \subseteq X$, and $\|f - g\|_\infty \leq \epsilon$. From [\(2.2\)](#) it follows that the mapping

$$(r, x) \longmapsto \int_0^r T_{-1}(\sigma) x d\sigma \quad (2.25)$$

from $\mathbb{R}_+ \times X$ into X_0 is continuous. The range of

$$\phi : \mathbb{R} \longrightarrow X_0 : t \longmapsto \int_0^s T_{-1}(\sigma) g(t-\sigma) d\sigma \quad (2.26)$$

is contained in $K_0 = \{n T_0(\tau) \int_0^r T_{-1}(\sigma) x d\sigma : 0 \leq \tau, r \leq s; x \in K\}$, and hence, K_0 is compact. On the other hand, by [\(2.2\)](#), there is a constant N independent of $t \in \mathbb{R}$ such that

$$\left\| \int_0^s T_{-1}(\sigma) (f(t-\sigma) - g(t-\sigma)) d\sigma \right\| \leq N \int_0^s \|f(t-\sigma) - g(t-\sigma)\| d\sigma \leq N s \epsilon. \quad (2.27)$$

Thus the range of ψ is contained in $K_0 + N s \epsilon B_{X_0}$, where B_{X_0} denotes the closed unit ball of X_0 . In particular, the range of ψ is totally bounded, which proves the claim.

Since ψ has relatively compact range we obtain

$$\lim_{\lambda \rightarrow \infty} (\lambda R(\lambda, A_0) - \mu R(\mu, A_0)) z(t, t-s) = 0 \quad \text{uniformly for } t \in \mathbb{R}. \quad (2.28)$$

If $w_\lambda(t, t-s) = \int_{t-s}^t U_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma$, $t \in \mathbb{R}$, then as in the proof of [Theorem 2.2](#) we derive from [\(2.28\)](#) and [\(2.13\)](#) that

$$\lim_{\lambda, \mu \rightarrow \infty} \|w_\lambda(t, t-s) - w_\mu(t, t-s)\| = 0 \quad (2.29)$$

uniformly for $t \in \mathbb{R}$. This completes the proof. \square

The following lemma will be used in [Section 3](#).

LEMMA 2.6. *Let $f \in L^1_{\text{loc}}(\mathbb{R}, X)$ and let $u \in C(\mathbb{R}, X_0)$ be a mild solution of (2.3). If $\phi \in C^1(\mathbb{R})$, then ϕu is a mild solution of (1.4) with f replaced by $\phi' u + \phi f$.*

Proof. If $t \geq s$, then the representation of u obtained in [Theorem 2.2](#) leads to

$$\begin{aligned}
& \int_s^t U_B(t, \sigma) \phi'(\sigma) u(\sigma) d\sigma \\
&= \int_s^t U_B(t, \sigma) \phi'(\sigma) U_B(\sigma, s) u(s) d\sigma \\
&+ \lim_{\lambda \rightarrow \infty} \int_s^t U_B(t, \sigma) \phi'(\sigma) \int_s^\sigma U_B(\sigma, \tau) \lambda R(\lambda, A) f(\tau) d\tau d\sigma \\
&= (\phi(t) - \phi(s)) U_B(t, s) u(s) \\
&+ \lim_{\lambda \rightarrow \infty} \int_s^t \int_\tau^t \phi'(\sigma) U_B(t, \tau) \lambda R(\lambda, A) f(\tau) d\sigma d\tau \\
&= \phi(t) \left(U_B(t, s) u(s) + \lim_{\lambda \rightarrow \infty} \int_s^t U_B(t, \tau) \lambda R(\lambda, A) f(\tau) d\tau \right) \\
&- U_B(t, s) \phi(s) u(s) - \lim_{\lambda \rightarrow \infty} \int_s^t U_B(t, \tau) \lambda R(\lambda, A) \phi(\tau) f(\tau) d\tau.
\end{aligned} \tag{2.30}$$

Another application of [Theorem 2.2](#) establishes the result. \square

3. Asymptotic properties of solutions of inhomogeneous equations

In this section, we discuss conditions on the evolution family $(U_B(t, s))_{t \geq s}$ and the inhomogeneity $f \in L^1_{\text{loc}}(\mathbb{R}, X)$ which ensure that (2.3) has a (unique) mild solution u with a prescribed asymptotic behavior. For the rest of the paper we impose the following condition on the perturbation $(B(t))_{t \in \mathbb{R}}$.

(B) $\|B(\cdot)\| \leq b(\cdot)$ for some $b \in L^1_{\text{loc}, u}(\mathbb{R})$.

Note that (B) implies exponential boundedness of the evolution family $(U_B(t, s))_{t \geq s}$ (see (2.9)).

At first we discuss the case where $(U_B(t, s))_{t \geq s}$ has an exponential dichotomy. We recall the following notion (see [12, 18, 21, 23, 24, 25, 36]).

Definition 3.1. An evolution family $(U(t, s))_{t \geq s}$ on the Banach space Z has an *exponential dichotomy* with constants $\alpha > 0$, $L \geq 1$ if there exists a bounded, strongly continuous family of projections $(P(t))_{t \in \mathbb{R}} \subseteq \mathcal{L}(Z)$ such that for $t \geq s$

- (i) $P(t)U(t, s) = U(t, s)P(s)$,
- (ii) the map $U_1(t, s) : (Id - P(s))Z \rightarrow (Id - P(t))Z : z \mapsto U(t, s)z$ is invertible,
- (iii) $\|U(t, s)z\| \leq Le^{-\alpha(t-s)}\|z\|$ for $z \in P(s)Z$,
- (iv) $\|U_1(t, s)^{-1}z\| \leq Le^{-\alpha(t-s)}\|z\|$ for $z \in (Id - P(t))Z$.

In that case the family $(\Gamma(t, s))_{(t, s) \in \mathbb{R}^2} \subseteq \mathcal{L}(Z)$ given by

$$\Gamma(t, s) = \begin{cases} P(t)U(t, s)P(s), & t \geq s, \\ -(Id - P(t)) [U_1(s, t)]^{-1} (Id - P(s)), & t < s, \end{cases} \quad (3.1)$$

is called the corresponding *Green's operator function*.

Remark 3.2. It is shown in [38, Lemma VI.9.15] that $(t, s) \mapsto [U_1(t, s)]^{-1} (Id - P(t))$ is strongly continuous for $t \geq s$.

The existence of an exponential dichotomy for the evolution family $(U(t, s))_{t \geq s}$ on the Banach space Z allows to connect asymptotic properties of the solution $u(\cdot, f) \in C(\mathbb{R}, Z)$ of the integral equation

$$u(t, f) = U(t, s)u(s, f) + \int_s^t U(t, \tau)f(\tau)d\tau, \quad t \geq s, \quad (3.2)$$

with asymptotic properties of the function $f \in C(\mathbb{R}, Z)$. We recall the following result in [22, Theorem 2.1], (see also [23, Section 10.2, Theorem 1], [5, Theorem 4]). By $C_b(\mathbb{R}, Z)$ we denote the set of all bounded, continuous, Z -valued functions on \mathbb{R} , and $C_0(\mathbb{R}, Z)$ is the space of all functions in $C_b(\mathbb{R}, Z)$ vanishing at $\pm\infty$.

THEOREM 3.3. *Let $(U(t, s))_{t \geq s}$ be an exponentially bounded evolution family on the Banach space Z and let $\mathcal{F}(\mathbb{R}, Z)$ be the space $C_0(\mathbb{R}, Z)$ or $C_b(\mathbb{R}, Z)$. Then $(U(t, s))_{t \geq s}$ has an exponential dichotomy if and only if for every $f \in \mathcal{F}(\mathbb{R}, Z)$ there exists a unique solution $u(\cdot, f) \in \mathcal{F}(\mathbb{R}, Z)$ of (3.2). In that case $u(\cdot, f)$ is given by*

$$u(t, f) = \int_{-\infty}^{\infty} \Gamma(t, \sigma)f(\sigma)d\sigma, \quad t \in \mathbb{R}. \quad (3.3)$$

We will show a corresponding result on asymptotic properties of the mild solutions of the inhomogeneous equation (2.3). We stress that in our case the evolution family $(U_B(t, s))_{t \geq s}$ given by equation (2.7) consists of operators on the Banach space X_0 whereas the inhomogeneity f has values in the larger space X . The following lemma plays a central role. By $L_{\text{loc}, u}^1(\mathbb{R}, X)$ we denote the space of uniformly locally integrable functions from \mathbb{R} into X equipped with the norm $\|f\|_{1, \text{loc}, u} = \sup_{t \in \mathbb{R}} \int_{t-1}^t \|f(\sigma)\| d\sigma$.

LEMMA 3.4. *Assume that $(U_B(t, s))_{t \geq s}$ has an exponential dichotomy with constants $\alpha > 0$, $L \geq 1$, and projections $(P_B(t))_{t \geq 0}$. For $f \in L_{\text{loc}, u}^1(\mathbb{R}, X)$ and $\lambda > \omega$ define $u_\lambda(\cdot, f) \in C(\mathbb{R}, X_0)$ by*

$$u_\lambda(t, f) = \int_{-\infty}^{\infty} \Gamma_B(t, \sigma)\lambda R(\lambda, A)f(\sigma)d\sigma, \quad t \in \mathbb{R}, \quad (3.4)$$

where $(\Gamma_B(t, s))_{(t, s) \in \mathbb{R}^2}$ is the Green's operator function corresponding to $(U_B(t, s))_{t \geq s}$. Then

- (i) $\|u_\lambda(\cdot, f)\| \leq C\|f\|_{1, \text{loc}, u}$ for a constant C independent of $\lambda \geq \omega + 1$ and f .

- (ii) $(u_\lambda(\cdot, f))$ is uniformly convergent on compact intervals in \mathbb{R} as $\lambda \rightarrow \infty$.
- (iii) If $f \in \text{BUC}_r(\mathbb{R}, X)$, then $(u_\lambda(\cdot, f))$ is uniformly convergent on \mathbb{R} as $\lambda \rightarrow \infty$.

Proof. Let $Q_B(t) = Id - P_B(t)$, $t \in \mathbb{R}$. Since $(U_B(t, s))_{t \geq s}$ has an exponential dichotomy and A is a Hille-Yosida operator we obtain for $t \in \mathbb{R}$ and $\lambda \geq \omega + 1$

$$\begin{aligned}
\|u_\lambda(t, f)\| &\leq \int_t^\infty \left\| [U_B(\sigma, t)]^{-1} Q_B(\sigma) \lambda R(\lambda, A) f(\sigma) \right\| d\sigma \\
&\quad + \int_{-\infty}^t \|U_B(t, \sigma) P_B(\sigma) \lambda R(\lambda, A) f(\sigma)\| d\sigma \\
&\leq \sum_{k \geq 0} L e^{-\alpha k} \|\lambda R(\lambda, A)\| \int_{t+k}^{t+k+1} \|Q_B(\sigma)\| \|f(\sigma)\| d\sigma \quad (3.5) \\
&\quad + \sum_{k \geq 0} L e^{-\alpha k} \|\lambda R(\lambda, A)\| \int_{t-k-1}^{t-k} \|P_B(\sigma)\| \|f(\sigma)\| d\sigma \\
&\leq C \|f\|_{1, \text{loc}, u},
\end{aligned}$$

where C is a constant independent of f . This proves assertion (i) and the continuity of u_λ follow.

In order to show (ii) note that

$$u_\lambda(t, f) = U_B(t, s) u_\lambda(s, f) + \int_s^t U_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma \quad \text{for } t \geq s \quad (3.6)$$

(see [22, Proof of Proposition 1.2]). For $\lambda, \mu > \omega + 1$, $t \in \mathbb{R}$, and $r > 0$ we have

$$\begin{aligned}
&\|P_B(t)(u_\lambda(t, f) - u_\mu(t, f))\| \\
&\leq \|U_B(t, t-r) P_B(t-r)(u_\lambda(t-r, f) - u_\mu(t-r, f))\| \\
&\quad + \left\| P_B(t) \int_{t-r}^t U_B(t, \sigma) (\lambda R(\lambda, A) - \mu R(\mu, A)) f(\sigma) d\sigma \right\| \quad (3.7) \\
&\leq L e^{-\alpha r} C_1 + \left\| P_B(t) \int_{t-r}^t U_B(t, \sigma) (\lambda R(\lambda, A) - \mu R(\mu, A)) f(\sigma) d\sigma \right\|,
\end{aligned}$$

where $C_1 = \sup\{\|P_B(t)\| \|u_\lambda(t, f) - u_\mu(t, f)\| : t \in \mathbb{R}; \lambda, \mu > \omega + 1\}$. By Theorem 2.2, $\lim_{\lambda \rightarrow \infty} \lambda \int_s^t U_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma$ exists uniformly for $t \geq s$ in compact intervals in \mathbb{R} . Thus, if in (3.7) we choose $r > 0$ sufficiently large and then consider $\lambda, \mu \rightarrow \infty$ we obtain

$$\lim_{\lambda, \mu \rightarrow \infty} \|P_B(t)(u_\lambda(t, f) - u_\mu(t, f))\| = 0 \quad (3.8)$$

uniformly for t in compact intervals in \mathbb{R} .

On the other hand, for $\lambda, \mu > \omega + 1$, $t \in \mathbb{R}$, and $r > 0$ we obtain

$$\begin{aligned} & \left\| [U_B(t+r, t)]^{-1} Q_B(t+r)(u_\lambda(t+r, f) - u_\mu(t+r, f)) \right\| \\ & \geq \|Q_B(t)(u_\lambda(t, f) - u_\mu(t, f))\| \\ & \quad - \left\| [U_B(t+r, t)]^{-1} Q_B(t+r) \int_t^{t+r} U_B(t+r, \sigma) (\lambda R(\lambda, A) - \mu R(\mu, A)) f(\sigma) d\sigma \right\|. \end{aligned} \quad (3.9)$$

Thus

$$\begin{aligned} & \|Q_B(t)(u_\lambda(t, f) - u_\mu(t, f))\| \\ & \leq L e^{-\alpha r} \left(C_2 + \left\| Q_B(t+r) \int_t^{t+r} U_B(t+r, \sigma) (\lambda R(\lambda, A) - \mu R(\mu, A)) f(\sigma) d\sigma \right\| \right), \end{aligned} \quad (3.10)$$

where $C_2 = \sup\{\|Q_B(t)\| \|u_\lambda(t, f) - u_\mu(t, f)\| : t \in \mathbb{R}; \lambda, \mu > \omega + 1\}$. As above, if we choose $r > 0$ sufficiently large and apply [Theorem 2.2](#) we obtain

$$\lim_{\lambda, \mu \rightarrow \infty} \|Q_B(t)(u_\lambda(t, f) - u_\mu(t, f))\| = 0 \quad (3.11)$$

uniformly for t in compact intervals of \mathbb{R} . Assertion (ii) is now an immediate consequence of (3.8) and (3.11).

Finally, if $f \in \text{BUC}_r(\mathbb{R}, X)$, then (3.7) and (3.10) together with [Proposition 2.5](#) imply that

$$\begin{aligned} & \lim_{\lambda, \mu \rightarrow \infty} \|P_B(t)(u_\lambda(t, f) - u_\mu(t, f))\| = 0, \\ & \lim_{\lambda, \mu \rightarrow \infty} \|Q_B(t)(u_\lambda(t, f) - u_\mu(t, f))\| = 0, \end{aligned} \quad (3.12)$$

uniformly for $t \in \mathbb{R}$. This proves (iii). \square

We come to our first main result. It is an analogue of [Theorem 3.3](#) and connects asymptotic properties of mild solutions of (2.3) with the existence of an exponential dichotomy for the evolution family $(U_B(t, s))_{t \geq s}$. In the special case where $B(t) = B$ is constant a similar result has been shown in [2] by completely different methods.

THEOREM 3.5. *The following assertions are equivalent.*

- (i) *The evolution family $(U_B(t, s))_{t \geq s}$ has an exponential dichotomy.*
- (ii) *For every $f \in L^1_{\text{loc}, u}(\mathbb{R}, X)$ there is a unique mild solution $u \in C_b(\mathbb{R}, X_0)$ of (2.3).*
- (iii) *For every $f \in C_b(\mathbb{R}, X)$ there is a unique mild solution $u \in C_b(\mathbb{R}, X_0)$ of (2.3).*
- (iv) *For every $f \in C_0(\mathbb{R}, X)$ there is a unique mild solution $u \in C_0(\mathbb{R}, X_0)$ of (2.3).*

In that case the function $u = u(\cdot, f)$ is given by

$$u(t, f) = \lim_{\lambda \rightarrow \infty} \int_{-\infty}^{\infty} \Gamma_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma, \quad t \in \mathbb{R}, \quad (3.13)$$

where $(\Gamma_B(t, s))_{(t, s) \in \mathbb{R}^2}$ is the Green's operator function corresponding to $(U_B(t, s))_{t \geq s}$.

Proof. (i) \Rightarrow (ii). Assume that $(U_B(t, s))_{t \geq s}$ has an exponential dichotomy and let $f \in L^1_{\text{loc}, u}(\mathbb{R}, X)$. [Lemma 3.4](#) implies that the limit function $u = u(\cdot, f)$ in [\(3.13\)](#) is defined and $u \in C_b(\mathbb{R}, X)$. We claim that $u(\cdot, f)$ is a mild solution of [\(2.3\)](#). In fact, if $t \geq s$, then

$$\begin{aligned}
& u(t, f) - U_B(t, s)u(s, f) \\
&= \lim_{\lambda \rightarrow \infty} \lambda \left(\int_{-\infty}^{\infty} \Gamma_B(t, \sigma) R(\lambda, A) f(\sigma) d\sigma - \int_{-\infty}^s U_B(t, \sigma) P_B(\sigma) R(\lambda, A) f(\sigma) d\sigma \right. \\
&\quad + \int_s^t U_B(t, \sigma) Q_B(\sigma) R(\lambda, A) f(\sigma) d\sigma \\
&\quad \left. + \int_t^{\infty} [U_B(\sigma, t)]^{-1} Q_B(\sigma) R(\lambda, A) f(\sigma) d\sigma \right) \\
&= \lim_{\lambda \rightarrow \infty} \lambda \left(\int_{-\infty}^{\infty} \Gamma_B(t, \sigma) R(\lambda, A) f(\sigma) d\sigma - \int_{-\infty}^t U_B(t, \sigma) P_B(\sigma) R(\lambda, A) f(\sigma) d\sigma \right. \\
&\quad + \int_s^t U_B(t, \sigma) R(\lambda, A) f(\sigma) d\sigma \\
&\quad \left. + \int_t^{\infty} [U_B(\sigma, t)]^{-1} Q_B(\sigma) R(\lambda, A) f(\sigma) d\sigma \right) \\
&= \lim_{\lambda \rightarrow \infty} \int_s^t U_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma.
\end{aligned} \tag{3.14}$$

By [Theorem 2.2](#), $u(\cdot, f)$ is a mild solution of [\(2.3\)](#). To show that $u(\cdot, f)$ is the only mild solution of [\(2.3\)](#) belonging to $C_b(\mathbb{R}, X)$ we can assume that $f \equiv 0$ and repeat the arguments in [\[22, proof of Proposition 1.2\]](#).

Since $C_b(\mathbb{R}, X) \subseteq L^1_{\text{loc}, u}(\mathbb{R}, X)$ the implication (ii) \Rightarrow (iii) is obvious.

(iii) \Rightarrow (iv). From the definition of a mild solution it follows immediately that the operator G assigning to each $f \in C_b(\mathbb{R}, X)$ the unique mild solution $u = u(\cdot, f) \in C_b(\mathbb{R}, X_0)$ of [\(2.3\)](#) is closed. Hence, G is bounded. Now let $f \in C_0(\mathbb{R}, X)$. We have to show that also $u(\cdot, f) \in C_0(\mathbb{R}, X)$. Let $n \in \mathbb{N}$ and choose $t_n > n$ such that $\sup_{|t| > t_n - n} \|f(t)\| < 1/n$. For $|t| > t_n$ choose $\phi_t \in C^1(\mathbb{R})$ such that $0 \leq \phi_t \leq 1$, $\phi_t(t) = 1$, $\text{supp } \phi_t \subseteq [t-n, t+n]$, and $\|\phi_t'\| \leq 2/n$. By [Lemma 2.6](#), $G(\phi_t' u + \phi_t f) = \phi_t u$. Hence

$$\|\phi_t u\|_{\infty} \leq \|G\| \|\phi_t' u + \phi_t f\|_{\infty} \leq n^{-1} \|G\| (2\|u\|_{\infty} + 1). \tag{3.15}$$

In particular, $\|u(t)\| = \|\phi_t(t)u(t)\| \leq n^{-1} \|G\| (2\|u\|_{\infty} + 1)$ for $|t| > t_n$. Hence $u \in C_0(\mathbb{R}, X_0)$. Since $C_0(\mathbb{R}, X_0) \subseteq C_0(\mathbb{R}, X)$ implication (iv) \Rightarrow (i) follows from [Theorem 3.3](#). \square

Remark 3.6. The arguments in the proof of (iii) \Rightarrow (iv) can be used to simplify parts of the proof of [\[22, Theorem 2.1\]](#) considerably.

Now we assume that the evolution family $(U_B(t, s))_{t \geq s}$ is *p-periodic*, in the sense that there exists $p > 0$ such that $U_B(t + p, s + p) = U_B(t, s)$ for $t \geq s$. From formula

(2.7) we see that $(U_B(t, s))_{t \geq s}$ is p -periodic provided that $t \mapsto B(t)$ is p -periodic, that is, $B(t) = B(t + p)$. We call $U_B(p, 0)$ the *monodromy operator* of the evolution family $(U_B(t, s))_{t \geq s}$. On $C(\mathbb{R}, X_0)$ we define the operator T by

$$Th(t) = U_B(t, t - p)h(t - p), \quad h \in C(\mathbb{R}, X_0), t \in \mathbb{R}. \quad (3.16)$$

If $u \in C(\mathbb{R}, X_0)$ is a mild solution of (2.3), then the representation formula for u obtained in [Theorem 2.2](#) leads to

$$(Id - T)u(t) = \lim_{\lambda \rightarrow \infty} \int_{t-p}^t U_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma, \quad t \in \mathbb{R}. \quad (3.17)$$

We need the notion of the *spectrum* $\text{sp}(f)$ of a Banach space-valued function $f : \mathbb{R} \rightarrow Z$ (cf. [\[3, 20, 23, 32, 35\]](#)). If $f \in C_b(\mathbb{R}, Z)$ we set

$$\begin{aligned} \text{sp}(f) = \{ \xi \in \mathbb{R} : \text{for every } \epsilon > 0 \text{ there exists } \phi \in L^1(\mathbb{R}), \\ \text{such that } \text{supp}(\hat{\phi}) \subseteq [\xi - \epsilon, \xi + \epsilon] \text{ and } \phi \star f \neq 0 \}, \end{aligned} \quad (3.18)$$

where $\hat{\phi}$ denotes the Fourier transform of ϕ and $\phi \star f$ is the convolution of ϕ and f . Moreover, we set

$$\Sigma_p(f) = \overline{\text{sp}(f) + (2\pi/p)\mathbb{Z}} \subseteq \mathbb{R}. \quad (3.19)$$

We obtain the following extension of [\[6, Theorem 3.8\]](#).

THEOREM 3.7. *Assume that the evolution family $(U_B(t, s))_{t \geq s}$ is p -periodic. Let $f \in C_b(\mathbb{R}, X)$ and suppose that $\sigma(U_B(p, 0)) \cap \{e^{i\eta p} : \eta \in \text{sp}(f)\} = \emptyset$. Then*

(a) *There is at most one mild solution $u \in C_b(\mathbb{R}, X_0)$ of (2.3) such that $\text{sp}(u) \subseteq \Sigma_p(f)$.*

(b) *Let $\mathcal{F}(\mathbb{R}, X_0)$ be a closed, translation-invariant subspace of $\text{BUC}(\mathbb{R}, X_0)$ such that $s \mapsto e^{2\pi ins/p} Rh(s)$ belongs to $\mathcal{F}(\mathbb{R}, X_0)$ whenever $h \in \mathcal{F}(\mathbb{R}, X_0)$, $R \in \mathcal{L}(X_0)$, and $n \in \mathbb{Z}$. Suppose that $f \in \text{BUC}_r(\mathbb{R}, X)$ such that $\lambda R(\lambda, A) f(\cdot) \in \mathcal{F}(\mathbb{R}, X_0)$ for $\lambda > \omega$. Then there exists a mild solution $u \in \mathcal{F}(\mathbb{R}, X_0)$ of (2.3), and u has relatively compact range.*

Proof. In order to prove (a) consider

$$\mathcal{M} = \{h \in C_b(\mathbb{R}, X_0) : \text{sp}(h) \subseteq \Sigma_p(f)\}. \quad (3.20)$$

In [\[6, proof of Theorem 3.8\]](#) it is shown that the operator T defined in (3.16) maps \mathcal{M} into itself and the restriction $T|_{\mathcal{M}}$ of T to \mathcal{M} is bounded and satisfies $1 \in \rho(T|_{\mathcal{M}})$. The invertibility of $Id - T|_{\mathcal{M}}$ and (3.17) show that there is at most one mild solution u of (2.3) contained in \mathcal{M} .

For the proof of (b) let

$$\mathcal{N} = \{h \in \mathcal{F}(\mathbb{R}, X_0) : \text{sp}(h) \subseteq \Sigma_p(f)\}. \quad (3.21)$$

In [\[6, proof of Theorem 3.8\]](#) it is shown that \mathcal{N} is T -invariant and $1 \in \rho(T|_{\mathcal{N}})$. For $\lambda > \omega$ set $f_\lambda = \lambda R(\lambda, A) f(\cdot)$. Note that $\text{sp}(f_\lambda) \subseteq \text{sp}(f)$. By [\[6, Theorem 3.8\]](#) for each $\lambda > \omega$

there is a (unique) mild solution $u_\lambda \in \mathcal{F}(\mathbb{R}, X_0)$ of (1.4) with f_λ instead of f such that $\text{sp}(u_\lambda) \subseteq \Sigma_p(f_\lambda) \subseteq \Sigma_p(f)$, and u_λ has relatively compact range. Let

$$w_\lambda(t) = \int_{t-p}^t U_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma, \quad t \in \mathbb{R}, \lambda > \omega. \quad (3.22)$$

Since $f \in \text{BUC}_r(\mathbb{R}, X)$ [Proposition 2.5](#) implies that $w(t) = \lim_{\lambda \rightarrow \infty} w_\lambda(t)$ exists uniformly for t in \mathbb{R} . From (3.17) we obtain $(Id - T_{|\mathcal{N}})u_\lambda = w_\lambda$, $\lambda > \omega$. In particular, $w_\lambda \in \mathcal{N}$ for $\lambda > \omega$, and $u_\lambda = (Id - T_{|\mathcal{N}})^{-1}w_\lambda$ converges uniformly to $u = (Id - T_{|\mathcal{N}})^{-1}w \in \mathcal{N}$ as $\lambda \rightarrow \infty$. From [Theorem 2.2](#) and the fact that each u_λ is a mild solution of (1.4) with f replaced by f_λ it follows that the limit function u is a mild solution of (2.3). Moreover, since each u_λ has relatively compact range also u has relatively compact range. This completes the proof. \square

Recall that a function $h \in \text{BUC}(\mathbb{R}, Z)$ is *almost periodic* if the set of translates $\{h(\cdot + t) : t \in \mathbb{R}\}$ is relatively compact in $\text{BUC}(\mathbb{R}, Z)$. By $AP(\mathbb{R}, Z)$ we denote the space of almost periodic, Z -valued functions. [Theorem 3.7](#) has the following immediate consequence (cf. [\[6, Corollary 3.9\]](#)).

COROLLARY 3.8. *Assume that $(U_B(t, s))_{t \geq s}$ is p -periodic. Let $f \in AP(\mathbb{R}, X)$, and suppose that $\sigma(U_B(p, 0)) \cap \{e^{i\eta p} : \eta \in \text{sp}(f)\} = \emptyset$. Then there is a unique $u \in C_b(\mathbb{R}, X_0)$ such that u is a mild solution of (2.3) and $\text{sp}(u) \subseteq \Sigma_p(f)$. Moreover, $u \in AP(\mathbb{R}, X_0)$.*

Let $S^1 = \{\lambda \in \mathbb{C} : |\lambda| = 1\}$ be the unit circle.

COROLLARY 3.9. *If the evolution family $(U_B(t, s))_{t \geq s}$ is p -periodic, then the following assertions are equivalent.*

- (i) $S^1 \subseteq \rho(U_B(p, 0))$.
- (ii) *For every $f \in AP(\mathbb{R}, X)$ there is a unique mild solution $u \in AP(\mathbb{R}, X_0)$ of (2.3).*

Proof. Note that (i) is equivalent to the existence of an exponential dichotomy for $(U_B(t, s))_{t \geq s}$ (see [\[19, Theorem 3.2.2\]](#), [\[18, Theorem 7.2.3\]](#)). Hence if (i) is satisfied and $f \in AP(\mathbb{R}, X)$, the existence of a mild solution $u \in AP(\mathbb{R}, X_0)$ of (2.3) follows from [Corollary 3.8](#), whereas the uniqueness is a consequence of [Theorem 3.5](#). The converse implication (ii) \Rightarrow (i) follows immediately from [\[27, Lemma 4\]](#). \square

By $P_p(\mathbb{R}, Z)$ we denote the space of p -periodic, continuous, Z -valued functions on \mathbb{R} .

COROLLARY 3.10. *If the evolution family $(U_B(t, s))_{t \geq s}$ is p -periodic, then the following assertions are equivalent:*

- (i) $1 \in \rho(U_B(p, 0))$.
- (ii) *For every $f \in P_p(\mathbb{R}, X)$, there exists a unique mild solution $u \in P_p(\mathbb{R}, X_0)$ of (2.3).*

Proof. Assume that (i) is satisfied. If $f \in P_p(\mathbb{R}, X)$, then $\text{sp}(f) \subseteq (2\pi/p)\mathbb{Z}$ (see [32, Example 0.1]). Hence, by Theorem 3.7, there is a unique mild solution $u \in P_p(\mathbb{R}, X_0)$ of (2.3). The implication (ii) \Rightarrow (i) follows immediately from [19, Theorem 3.3.4] (see also [27, Proposition 1]). \square

Remark 3.11. The operator T_{P_p} on $P_p(\mathbb{R}, X_0)$ satisfies $T_{P_p}h(t) = U_B(t, t-p)h(t)$, $h \in P_p(\mathbb{R}, X_0)$, $t \in \mathbb{R}$. Moreover, $1 \in \rho(U_B(p, 0))$ implies $1 \in \rho(U_B(t+p, t))$ for all $t \in \mathbb{R}$ (see [18, Lemma 7.2.2]). From this we obtain $(Id - T_{P_p})^{-1}h(t) = (Id - U(t, t-p))^{-1}h(t)$, $h \in P_p(\mathbb{R}, X_0)$, $t \in \mathbb{R}$. In particular, by (3.17), the mild solution u obtained in Corollary 3.10(ii) has the representation

$$u(t) = \lim_{\lambda \rightarrow \infty} (Id - U_B(t, t-p))^{-1} \int_{t-p}^t U_B(t, \sigma) \lambda R(\lambda, A) f(\sigma) d\sigma, \quad t \in \mathbb{R}. \quad (3.23)$$

4. The semilinear equation

In this section, we apply the results of Section 3 to the semilinear equation

$$\frac{d}{dt}u(t) = (A + B(t))u(t) + F(t, u(t)), \quad t \in \mathbb{R}, \quad (4.1)$$

where A and $B(t)$, $t \in \mathbb{R}$, are as in the previous sections and $F : \mathbb{R} \times X_0 \rightarrow X$ is jointly continuous and Lipschitz continuous in the second variable with Lipschitz constant l independent of t and x . Moreover, we assume that $t \mapsto F(t, 0)$ is a bounded function on \mathbb{R} . Our definition of a mild solution of (4.1) is similar to Definition 2.1.

Definition 4.1. A function $u \in C(\mathbb{R}, X_0)$ is called a *mild solution* of (4.1) if

$$u(t) = T_0(t-s)u(s) + \int_s^t T_{-1}(t-\sigma)(B(\sigma)u(\sigma) + F(\sigma, u(\sigma))) d\sigma \quad \text{for } t \geq s. \quad (4.2)$$

The following conditions will be needed.

(H1) The evolution family $(U_B(t, s))_{t \geq s}$ has an exponential dichotomy with constants $\alpha > 0$, $L \geq 1$, and projections $(P_B(t))_{t \in \mathbb{R}}$, and $l < \alpha/2LC$, where $C = \sup_{t \in \mathbb{R}} \times \sup_{\lambda > \omega} \{\|\lambda P_B(t)R(\lambda, A)\|, \|\lambda(Id - P_B(t))R(\lambda, A)\|\} < \infty$.

(H2) The evolution family $(U_B(t, s))_{t \geq s}$ is p -periodic, $1 \in \rho(U_B(p, 0))$, and $l < (\tilde{C}pC)^{-1}$, where $C = \sup_{t \in \mathbb{R}} \|(Id - U(t, t-p))^{-1}\|$ and $\tilde{C} = \sup_{t \in \mathbb{R}} \|U(t, t-p)\|$.

THEOREM 4.2. *If condition (H1) holds, then there exists exactly one mild solution $u \in C_b(\mathbb{R}, X_0)$ of (4.1).*

Proof. For $f \in C_b(\mathbb{R}, X_0)$ set

$$Sf(t) = \lim_{\lambda \rightarrow \infty} \int_{-\infty}^{\infty} \Gamma_B(t, \sigma) \lambda R(\lambda, A) F(\sigma, f(\sigma)) d\sigma, \quad t \in \mathbb{R}. \quad (4.3)$$

By [Lemma 3.4](#) and the boundedness of $F(\cdot, 0)$, S is well defined and maps $C_b(\mathbb{R}, X_0)$ into itself. If $f, g \in C_b(\mathbb{R}, X_0)$, then

$$\begin{aligned} \|Sf - Sg\|_\infty &= \sup_{t \in \mathbb{R}} \left\| \lim_{\lambda \rightarrow \infty} \int_{-\infty}^{\infty} \Gamma_B(t, \sigma) \lambda R(\lambda, A) (F(\sigma, f(\sigma)) - F(\sigma, g(\sigma))) d\sigma \right\| \\ &\leq \sup_{t \in \mathbb{R}} CL \int_{-\infty}^{\infty} e^{-\alpha|t-\sigma|} l \|f - g\|_\infty d\sigma \leq \frac{2CL}{\alpha} l \|f - g\|_\infty. \end{aligned} \quad (4.4)$$

By our assumption $(2CL/\alpha)l < 1$. Hence S is a contraction, and by Banach's fixed point theorem there is a unique function $u \in C_b(\mathbb{R}, X_0)$ such that

$$u(t) = \lim_{\lambda \rightarrow \infty} \int_{-\infty}^{\infty} \Gamma(t, \sigma) \lambda R(\lambda, A) F(\sigma, u(\sigma)) d\sigma, \quad t \in \mathbb{R}. \quad (4.5)$$

[Theorem 3.5](#) implies that u is the unique mild solution of [\(4.1\)](#) contained in $C_b(\mathbb{R}, X_0)$. \square

In the same way, the following two results can be derived from [Theorem 3.5](#) and [Corollary 3.9](#), respectively.

PROPOSITION 4.3. *Assume that condition (H1) holds and that $\lim_{t \rightarrow \pm\infty} F(t, y) = 0$ uniformly for y in compact sets in X_0 . Then there exists exactly one mild solution $u \in C_0(\mathbb{R}, X_0)$ of [\(4.1\)](#).*

PROPOSITION 4.4. *Assume that condition (H1) holds and that the evolution family $(U_B(t, s))_{t \geq s}$ is p -periodic. If $F(\cdot, x)$ is almost periodic uniformly for x in compact sets in X_0 , that is, for every compact set K in X_0 and every sequence (t_n) in \mathbb{R} there is a subsequence (s_n) of (t_n) such that $(F(t + s_n, x))$ converges uniformly for (t, x) in $\mathbb{R} \times K$, then there is exactly one mild solution $u \in AP(\mathbb{R}, X_0)$ of [\(4.1\)](#).*

The following result is the semilinear version of [Corollary 3.10](#).

THEOREM 4.5. *Assume that condition (H2) holds and that $F(t + p, x) = F(t, x)$ for every $t \in \mathbb{R}$ and every $x \in X_0$. Then there exists exactly one mild solution $u \in P_p(\mathbb{R}, X_0)$ of [\(4.1\)](#).*

Proof. For $f \in P_p(\mathbb{R}, X_0)$ set

$$Sf(t) = \lim_{\lambda \rightarrow \infty} (Id - U_B(t, t-p))^{-1} \int_{t-p}^t U(t, \sigma) \lambda R(\lambda, A) F(\sigma, f(\sigma)) d\sigma, \quad t \in \mathbb{R}. \quad (4.6)$$

By [Proposition 2.5](#) and [Remark 3.11](#), S is well-defined and maps $P_p(\mathbb{R}, X_0)$ into itself. If $f, g \in P_p(\mathbb{R}, X_0)$, then

$$\begin{aligned}
\|Sf - Sg\|_\infty &= \sup_{t \in \mathbb{R}} \left\| \lim_{\lambda \rightarrow \infty} \left(Id - U_B(t, t-p) \right)^{-1} \right. \\
&\quad \left. \times \int_{t-p}^t U_B(t, \sigma) \lambda R(\lambda, A) (F(\sigma, f(\sigma)) - F(\sigma, g(\sigma))) d\sigma \right\| \quad (4.7) \\
&\leq \tilde{C} p C l \|f - g\|_\infty.
\end{aligned}$$

Since $\tilde{C} p C l < 1$, the map S is contractive and there is a unique function $v \in P_p(\mathbb{R}, X_0)$ such that

$$v(t) = \lim_{\lambda \rightarrow \infty} \left(Id - U_B(t, t-p) \right)^{-1} \int_{t-p}^t U_B(t, \sigma) \lambda R(\lambda, A) F(\sigma, v(\sigma)) d\sigma, \quad t \in \mathbb{R}. \quad (4.8)$$

By [Corollary 3.10](#), there is a unique mild solution $u \in P_p(\mathbb{R}, X_0)$ of [\(1.4\)](#) where f is replaced by the function $F(\cdot, v(\cdot))$. The representation of u obtained in [Remark 3.11](#) shows that $u = v$, and hence v is a mild solution of [\(4.1\)](#). On the other hand, it follows from [\(3.17\)](#) and [Remark 3.11](#) that each p -periodic mild solution of [\(4.1\)](#) satisfies [\(4.8\)](#). Hence v is the only p -periodic mild solution of [\(4.1\)](#). \square

5. Nonautonomous retarded differential equations

In this section, we apply the results obtained for [\(1.4\)](#) to retarded differential equations. Throughout the whole section Y is a fixed Banach space. We consider the inhomogeneous nonautonomous retarded differential equation

$$\frac{d}{dt} w(t) = C w(t) + K(t) w_t + h(t), \quad t \in \mathbb{R}, \quad (5.1)$$

where $(C, D(C))$ is a Hille-Yosida operator on Y and $h \in L^1_{\text{loc}}(\mathbb{R}, Y)$. The part C_0 of C on $Y_0 = \overline{D(C)}$ generates a C_0 -semigroup $(S_0(t))_{t \geq 0}$ on Y_0 , and by $(S_{-1}(t))_{t \geq 0}$ we denote the corresponding extrapolated C_0 -semigroup on the extrapolation space Y_{-1} . We set $E = C([-p, 0], Y_0)$, $p > 0$, and for a function $w \in C(\mathbb{R}, Y_0)$ we define $w_t \in E$ by $w_t(r) = w(t+r)$, $r \in [-p, 0]$. Finally, we assume that $K(t)$, $t \in \mathbb{R}$, is a family of operators in $\mathcal{L}(E, Y)$ such that $t \mapsto K(t)\phi$ is strongly measurable for every $\phi \in E$, and $\|K(\cdot)\| \leq d(\cdot)$ for a function $d \in L^1_{\text{loc}, u}(\mathbb{R})$. We define mild solutions of [\(5.1\)](#) as follows (cf. [\[4, 15, 16, 28, 34, 40\]](#)).

Definition 5.1. If $h \in L^1_{\text{loc}}(\mathbb{R}, Y)$, then $w = w(\cdot, h) \in C(\mathbb{R}, Y_0)$ is called a *mild solution* of [\(5.1\)](#) if

$$w(t) = S_0(t-s)w(s) + \int_s^t S_{-1}(t-\sigma) (K(\sigma)w_\sigma + h(\sigma)) d\sigma \quad \text{for } t \geq s. \quad (5.2)$$

Remark 5.2. If C is the generator of a C_0 -semigroup on Y , then the above definition of a mild solution coincides with that given in [\[15, 28, 34, 40\]](#).

In [16] (see also [33, 39]) it is shown how (5.1) can be transformed into an equation of the form of (1.4). For this we set $X = Y \times E$ and consider the equation

$$\frac{d}{dt}u(t) = Au(t) + B(t)u(t) + f(t), \quad t \in \mathbb{R}, \quad (5.3)$$

where $A : D(A) \rightarrow X$ is the linear operator on X given by

$$\begin{aligned} A \begin{pmatrix} 0 \\ \phi \end{pmatrix} &= \begin{pmatrix} -\phi'(0) + C\phi(0) \\ \phi' \end{pmatrix}, \\ D(A) &= \left\{ \begin{pmatrix} 0 \\ \phi \end{pmatrix} \in \{0\} \times E : \phi \in C^1([-p, 0], Y_0), \phi(0) \in D(C) \right\}, \end{aligned} \quad (5.4)$$

$B(t) \in \mathcal{L}(\{0\} \times E, X)$, $t \in \mathbb{R}$, is defined by

$$B(t) \begin{pmatrix} 0 \\ \phi \end{pmatrix} = \begin{pmatrix} K(t)\phi \\ 0 \end{pmatrix}, \quad (5.5)$$

and $f(\cdot) = \begin{pmatrix} h(\cdot) \\ 0 \end{pmatrix}$. It is shown in [33] that A is a Hille-Yosida operator on X , and the C_0 -semigroup $(T_0(t))_{t \geq 0}$ generated by the part A_0 of A in $X_0 = \overline{D(A)} = \{0\} \times E$ is given by

$$(T_0(t)|_E \phi)(r) = \begin{cases} \phi(t+r) & \text{if } t+r \leq 0, \\ S_0(t+r)\phi(0) & \text{if } t+r > 0. \end{cases} \quad (5.6)$$

We recall the following results obtained in [16, Theorem 5.3 and Proposition 5.4].

PROPOSITION 5.3. (a) *If for $u \in C(\mathbb{R}, E)$ the map $t \mapsto \begin{pmatrix} 0 \\ u(t) \end{pmatrix}$ is a mild solution of (2.3), then $t \mapsto u(t)(0)$ is a mild solution of (5.1) and $u(t)(\xi) = u(t+\xi)(0)$ for $t \in \mathbb{R}$ and $\xi \in [-p, 0]$.*

(b) *If $w \in C(\mathbb{R}, Y_0)$ is a mild solution of (5.1), then $t \mapsto \begin{pmatrix} 0 \\ w_t \end{pmatrix}$ is a mild solution of (2.3).*

PROPOSITION 5.4. *If $(U_B(t, s))_{t \geq s}$ is the evolution family on E determined by the variation-of-parameters formula*

$$\begin{pmatrix} 0 \\ U_B(t, s)\phi \end{pmatrix} = T_0(t-s) \begin{pmatrix} 0 \\ \phi \end{pmatrix} + \int_s^t T_{-1}(t-\sigma)B(\sigma) \begin{pmatrix} 0 \\ U_B(\sigma, s)\phi \end{pmatrix} d\sigma, \quad (5.7)$$

$t \geq s$, $\phi \in E$, then each mild solution $w \in C(\mathbb{R}, Y_0)$ of (5.1), with $h(t) = 0$ for all t , satisfies

$$w_t = U_B(t, s)w_s \quad \text{for } t \geq s. \quad (5.8)$$

Furthermore, if $\phi \in E$ and $t \geq s$, then

$$(U_B(t, s)\phi)(\xi) = \begin{cases} S_0(t+\xi-s)\phi(0) \\ \quad + \int_s^{t+\xi} S_{-1}(t+\xi-\sigma)K(\sigma)U_B(\sigma, s)\phi d\sigma, & t+\xi \geq s, \\ \phi(t+\xi-s), & t+\xi \leq s. \end{cases} \quad (5.9)$$

Asymptotic properties of the mild solutions of (5.1) are connected with properties of the evolution family $(U_B(t, s))_{t \geq s}$ on E in the following way.

THEOREM 5.5. *Assume that the evolution family $(U_B(t, s))_{t \geq s}$ defined by (5.7) has an exponential dichotomy. Then*

- (a) *For every $h \in L_{\text{loc}, u}^1(\mathbb{R}, Y)$ (in particular, for every $h \in C_b(\mathbb{R}, Y)$) there exists a unique mild solution $w \in C_b(\mathbb{R}, Y_0)$ of (5.1).*
- (b) *For every $h \in C_0(\mathbb{R}, Y)$ there exists a unique mild solution $w \in C_0(\mathbb{R}, Y_0)$ of (5.1).*

Proof. If h is in $L_{\text{loc}, u}^1(\mathbb{R}, Y)$ (respectively, $C_0(\mathbb{R}, Y)$), then $f : \mathbb{R} \rightarrow X$ defined by $f(t) = \begin{pmatrix} h(t) \\ 0 \end{pmatrix}$ is in $C_b(\mathbb{R}, X)$ (respectively, $C_0(\mathbb{R}, X)$). [Proposition 5.3](#) shows that there is a one-to-one correspondence between the mild solutions $w \in C(\mathbb{R}, Y_0)$ of (5.1) and the mild solutions $u \in C(\mathbb{R}, X_0)$ of (2.3), and w is in $C_b(\mathbb{R}, Y_0)$ (respectively, $C_0(\mathbb{R}, Y_0)$) if and only if u is in $C_b(\mathbb{R}, X_0)$ (respectively, $C_0(\mathbb{R}, X_0)$). An application of [Theorem 3.5](#) and proves the theorem. \square

In the same way the following results can be derived from [Corollary 3.9](#) and [Corollary 3.10](#).

THEOREM 5.6. *If the evolution family $(U_B(t, s))_{t \geq s}$ defined by (5.7) is p -periodic and $S^1 \subseteq \rho(U_B(p, 0))$, then for every $h \in AP(\mathbb{R}, Y)$ there is a unique mild solution $w \in AP(\mathbb{R}, Y_0)$ of (5.1).*

THEOREM 5.7. *If the evolution family $(U_B(t, s))_{t \geq s}$ defined by (5.7) is p -periodic and $1 \in \rho(U_B(p, 0))$, then for every $h \in P_p(\mathbb{R}, Y)$ there is a unique mild solution $w \in P_p(\mathbb{R}, Y_0)$ of (5.1).*

To give more concrete results we impose the following additional condition on the family $K(t)$, $t \in \mathbb{R}$.

- (K) Each operator $K(t)$, $t \in \mathbb{R}$, is of the form

$$K(t)\phi = \hat{K}(t)\phi(-p), \quad \phi \in E, \quad (5.10)$$

where $\hat{K}(t)$, $t \in \mathbb{R}$, is a p -periodic family in $\mathcal{L}(Y_0, Y)$ such that $t \mapsto \hat{K}(t)y$ is strongly measurable for all $y \in Y_0$, and $\|\hat{K}(\cdot)\| \leq d(\cdot)$ for some $d \in L_{\text{loc}}^1(\mathbb{R})$.

If condition (K) holds, then the evolution family $(U_B(t, s))_{t \geq s}$ is p -periodic. Now, we want to determine the spectrum $\sigma(U_B(p, 0))$ of the monodromy operator $U_B(p, 0)$. To that purpose we consider for each $\lambda \in \mathbb{C}$ the evolution family $(V_K^\lambda(t, s))_{t \geq s}$ on Y_0 determined by the integral equation

$$V_K^\lambda(t, s)y = S_0(t-s)y + \int_s^t S_{-1}(t-\sigma)e^{-\lambda p}\hat{K}(\sigma)V_K^\lambda(\sigma, s)y\,d\sigma \quad (5.11)$$

for $y \in Y_0$ and $t \geq s$. The existence of $(V_K^\lambda(t, s))_{t \geq s}$ is guaranteed by the same reasons as for the evolution family $(U_B(t, s))_{t \geq s}$ in (2.7). One can easily see that $(V_K^\lambda(t, s))_{t \geq s}$ is p -periodic for every $\lambda \in \mathbb{C}$.

Remark 5.8. If $\lambda \in \mathbb{C}$ and $y \in Y_0$, then $t \mapsto V_K^\lambda(t, s)y$, $t \geq s$, is the unique mild solution of the initial value problem

$$\begin{aligned} \frac{d}{dt}w(t) &= Cw(t) + e^{-\lambda p}\hat{K}(t)w(t), \quad t \geq s, \\ w(s) &= y \in Y_0. \end{aligned} \quad (5.12)$$

Next we derive a generalized *characteristic equation* for (5.1), with $h(t) = 0$ for all t , under the additional condition (K). In the finite dimensional case, that is, $Y = \mathbb{C}^n$, $C = 0$, and $\hat{K}(t)y = k(t)y$, where k is a p -periodic matrix-valued function, this leads to the classical characteristic equation as it can be found, for example, in [17, Theorem 8.3.1]. In the autonomous case, that is, $\hat{K}(t) = \hat{K} \in \mathcal{L}(Y_0)$, a related result is shown in [14, Chapter VI, Proposition 6.7].

THEOREM 5.9. *Assume that condition (K) holds and let $\lambda \in \mathbb{C}$. If $(U_B(t, s))_{t \geq s}$ is the evolution family defined in (5.7) and $(V_K^\lambda(t, s))_{t \geq s}$ is the evolution family defined in (5.11), then*

$$e^{\lambda p} \in \rho(U_B(p, 0)) \quad \text{if and only if } e^{\lambda p} \in \rho(V_K^\lambda(p, 0)). \quad (5.13)$$

Proof. First we show the “only if” part. Let $e^{\lambda p} \in \rho(U_B(p, 0))$. Then for every $\phi \in E$ there exists $\psi_\phi \in E$ such that $\phi = e^{\lambda p}\psi_\phi - U_B(p, 0)\psi_\phi$. By (5.9) we have

$$\phi(-p) = e^{\lambda p}\psi_\phi(-p) - \psi_\phi(0), \quad (5.14)$$

and

$$\begin{aligned} \phi(\xi) &= e^{\lambda p}\psi_\phi(\xi) - S_0(p + \xi)\psi_\phi(0) - \int_0^{p+\xi} S_{-1}(p + \xi - \sigma)\hat{K}(\sigma)\psi_\phi(\sigma - p)d\sigma \\ &= e^{\lambda p}\psi_\phi(\xi) + S_0(p + \xi)\phi(-p) - e^{\lambda p}S_0(p + \xi)\psi_\phi(-p) \\ &\quad - \int_{-p}^{\xi} S_{-1}(\xi - \sigma)\hat{K}(\sigma)\psi_\phi(\sigma)d\sigma \end{aligned} \quad (5.15)$$

for $\xi \in [-p, 0]$. In order to show surjectivity of $e^{\lambda p}Id - V_K^\lambda(p, 0)$ fix $y \in Y_0$ and set

$$\phi(\xi) = S_0(p + \xi)y, \quad \xi \in [-p, 0]. \quad (5.16)$$

Then (5.15) leads to

$$\psi_\phi(\xi) = S_0(p + \xi)\psi_\phi(-p) + \int_{-p}^{\xi} S_{-1}(\xi - \sigma)e^{-\lambda p}\hat{K}(\sigma)\psi_\phi(\sigma)d\sigma, \quad \xi \in [-p, 0]. \quad (5.17)$$

By Remark 5.8, $\xi \mapsto V_K^\lambda(\xi, -p)\psi_\phi(-p)$ is the unique mild solution of $(\lambda P)_{-p, \psi_\phi(-p)}$. Hence $\psi_\phi(\xi) = V_K^\lambda(\xi, -p)\psi_\phi(-p)$ for $\xi \in [-p, 0]$. Since $\phi(-p) = y$ we obtain from (5.14)

$$y = e^{\lambda p}\psi_\phi(-p) - V_K^\lambda(0, -p)\psi_\phi(-p) \quad (5.18)$$

which proves the surjectivity of $e^{\lambda p}Id - V_K^\lambda(p, 0)$.

In order to prove injectivity assume that $V_K^\lambda(0, -p)x = e^{\lambda p}x$ for some $x \in Y_0$. Let $\psi_x(\xi) = V_K^\lambda(\xi, -p)x$, $\xi \in [-p, 0]$. Using (5.11), (5.9), and condition (K), a straightforward computation shows that $U_B(0, -p)\psi_x = e^{\lambda p}\psi_x$. Thus $\psi_x = 0$. In particular, $x = \psi_x(-p) = 0$.

Now we prove the “if” part. Let $\phi \in E$. An application of the generalized form of Banach’s fixed point theorem shows that for every $y \in Y_0$ there is $\psi_y \in E$ such that

$$e^{\lambda p}\psi_y(\xi) = \phi(\xi) + S_0(p + \xi)y + \int_{-p}^{\xi} S_{-1}(\xi - \sigma)\hat{K}(\sigma)\psi_y(\sigma)d\sigma \quad (5.19)$$

for $\xi \in [-p, 0]$. By subtracting (5.11) we obtain

$$\begin{aligned} e^{\lambda p}\psi_y(\xi) - V_K^\lambda(\xi, -p)y \\ = \phi(\xi) + \int_{-p}^{\xi} S_{-1}(\xi - \sigma)e^{-\lambda p}\hat{K}(\sigma)(e^{\lambda p}\psi_y(\sigma) - V_K^\lambda(\sigma, -p)y)d\sigma. \end{aligned} \quad (5.20)$$

For $y_1, y_2 \in Y_0$ this leads to

$$\begin{aligned} e^{\lambda p}\psi_{y_1}(\xi) - V_K^\lambda(\xi, -p)y_1 - e^{\lambda p}\psi_{y_2}(\xi) + V_K^\lambda(\xi, -p)y_2 \\ = \int_{-p}^{\xi} S_{-1}(\xi - \sigma)e^{-\lambda p}\hat{K}(\sigma)(e^{\lambda p}\psi_{y_1}(\sigma) - V_K^\lambda(\sigma, -p)y_1 \\ - e^{\lambda p}\psi_{y_2}(\sigma) + V_K^\lambda(\sigma, -p)y_2)d\sigma. \end{aligned} \quad (5.21)$$

An application of Gronwall’s inequality yields

$$e^{\lambda p}\psi_{y_1}(\xi) - V_K^\lambda(\xi, -p)y_1 = e^{\lambda p}\psi_{y_2}(\xi) - V_K^\lambda(\xi, -p)y_2 \quad \text{for } \xi \in [-p, 0]. \quad (5.22)$$

By the assumption and the y -independence of $e^{\lambda p}\psi_y - V_K^\lambda(\cdot, -p)y$, we can choose $\tilde{y} \in Y_0$ such that

$$e^{\lambda p}\tilde{y} - V_K^\lambda(0, -p)\tilde{y} = \phi(0) + \int_{-p}^0 S_{-1}(-\sigma)e^{-\lambda p}\hat{K}(\sigma)(e^{\lambda p}\psi_{\tilde{y}}(\sigma) - V_K^\lambda(\sigma, -p)\tilde{y})d\sigma. \quad (5.23)$$

Evaluation of (5.20) at $\xi = 0$ then leads to

$$\tilde{y} = \psi_{\phi, \tilde{y}}(0). \quad (5.24)$$

By using (5.9) and (5.19) a direct computation yields $U_B(0, -p)\psi_{\tilde{y}} = e^{\lambda p}\psi_{\tilde{y}} - \phi$, which shows the surjectivity of $e^{\lambda p} - U_B(p, 0)$.

To prove injectivity assume that $e^{\lambda p}\psi - U_B(p, 0)\psi = 0$ for some $\psi \in E$. Then (5.9) leads to

$$e^{\lambda p}\psi(\xi) - S_0(p + \xi)\psi(0) - \int_{-p}^{\xi} S_{-1}(\xi - \sigma)\hat{K}(\sigma)\psi(\sigma)d\sigma = 0 \quad \text{for } \xi \in [-p, 0]. \quad (5.25)$$

Since $\xi \mapsto V_K^\lambda(\xi, -p)\psi(0)$ is the unique mild solution of $(\lambda P)_{-p, \psi(0)}$, we obtain $e^{\lambda p}\psi(\xi) = V_K^\lambda(\xi, -p)\psi(0)$, $\xi \in [-p, 0]$. In particular $e^{\lambda p}\psi(0) = V_K^\lambda(0, -p)\psi(0)$, and the invertibility of $e^{\lambda p}Id - V_K^\lambda(0, -p)$ implies $\psi(0) = 0$. Hence $\psi = e^{-\lambda p}V_K^\lambda(\cdot, -p) \times \psi(0) = 0$. \square

As a concrete example we discuss the retarded differential equation

$$\begin{aligned} \frac{\partial}{\partial t} w(t, x) &= \frac{\partial^2}{\partial x^2} w(t, x) - aw(t, x) - b(t)w(t-1, x) + f(t, x), \quad 0 \leq x \leq 2\pi, \quad t \geq s, \\ w(t, x) &= \varphi(t-s, x), \quad 0 \leq x \leq 2\pi, \quad s-1 \leq t \leq s, \end{aligned} \quad (5.26)$$

with initial value $\varphi \in C([-1, 0] \times [0, 2\pi])$. We assume that $a \in \mathbb{R}$, $b : \mathbb{R} \rightarrow \mathbb{R}$ is 1-periodic and locally integrable, and $f : \mathbb{R} \times [0, 2\pi] \rightarrow \mathbb{R}$ is continuous. It is known (see [11]) that on the Banach space $Y = C[0, 2\pi]$ the operator $(C, D(C))$ given by

$$\begin{aligned} (C\psi)(x) &= \frac{\partial^2}{\partial x^2} \psi(x) - a\psi(x), \quad x \in [0, 2\pi], \\ D(C) &= \{\psi \in C^2([0, 2\pi]) : \psi(0) = \psi(2\pi) = 0\}, \end{aligned} \quad (5.27)$$

is a Hille-Yosida operator. The spectrum of the part C_0 of C in $Y_0 = \overline{D(C)} = \{\psi \in C[0, 2\pi] : \psi(0) = \psi(2\pi) = 0\}$ is the set $\{-n^2 - a : n = 1, 2, 3, \dots\}$. For $E = C([-1, 0], Y_0)$ and $t \in \mathbb{R}$ we define

$$K(t) : E \longrightarrow Y_0 : \varphi \longmapsto -b(t)\varphi(t-1, \cdot). \quad (5.28)$$

Clearly, the operator family $(K(t))_{t \in \mathbb{R}}$ satisfies condition (K) for $p = 1$. The evolution family $(U_B(t, s))_{t \geq s}$ on E given by (5.7) is 1-periodic. Hence we can apply [Theorem 5.9](#) to determine the spectrum of $U_B(1, 0)$. We set $\bar{b} = \int_0^1 b(\tau) d\tau$.

PROPOSITION 5.10. $e^\lambda \in \sigma(U_B(1, 0))$ if and only if there exists $k \in \mathbb{Z}$ and $n \in \{1, 2, 3, \dots\}$ such that

$$\lambda + 2\pi ik = -e^{-\lambda} \bar{b} - n^2 - a. \quad (5.29)$$

Proof. The evolution family $(V_K^\lambda(t, s))_{t \geq s}$ defined in (5.11) is given by

$$V_K^\lambda(t, s) = e^{-\int_s^t e^{-\lambda} b(\tau) d\tau} S_0(t-s), \quad t \geq s, \quad (5.30)$$

where $(S_0(t))_{t \geq 0}$ is the C_0 -semigroup on $Y_0 = \overline{D(C)}$ generated by C_0 . From [Theorem 5.9](#) it follows that $e^\lambda \in \sigma(U_B(1, 0))$ if and only if $e^\lambda \in \sigma(e^{-\int_0^1 e^{-\lambda} b(\tau) d\tau} S(1))$. Since for C_0 the spectral mapping theorem holds we have

$$\sigma(S(1)) = \{e^{-n^2 - a} : n = 1, 2, 3, \dots\}. \quad (5.31)$$

Thus $e^\lambda \in \sigma(U_B(1, 0))$ if and only if

$$e^\lambda = e^{-\int_0^1 e^{-\lambda} b(\tau) d\tau} e^{-n^2 - a} = e^{-e^{-\lambda} \bar{b} - n^2 - a} \quad \text{for some } n \in \{1, 2, 3, \dots\}. \quad (5.32)$$

However, this is the case if and only if

$$\lambda + 2\pi ik = -e^{-\lambda} \bar{b} - n^2 - a \quad \text{for some } n \in \{1, 2, 3, \dots\} \text{ and some } k \in \mathbb{Z}. \quad (5.33)$$

□

While it appears difficult to determine the set of all $\lambda \in \mathbb{C}$ satisfying (5.29), there are results saying for which values of $n^2 + a$ and \bar{b} all solutions $\mu \in \mathbb{C}$ of

$$\mu = -e^{-\mu} \bar{b} - n^2 - a \quad (5.34)$$

have negative real part (cf. [17, page 135]). This is the case if $(n^2 + a, \bar{b})$ belongs to the shaded region shown in Figure 5.1.




FIGURE 5.1.

Since $n^2 \geq 1$, we obtain $\operatorname{Re} \mu < 0$ for each μ satisfying (5.34) for some $n \in \{1, 2, 3, \dots\}$ if (a, \bar{b}) is in the shaded region in Figure 5.2.

For example, if $a = -1$ and $\bar{b} = \pi/3$, then all $\lambda \in \mathbb{C}$ such that $e^\lambda \in \sigma(U_B(1, 0))$ have negative real part. Hence $(U_B(t, s))_{t \geq s}$ is exponentially stable and, in particular, has an exponential dichotomy. Note that in this case the semigroup $(S_0(t))_{t \geq 0}$ does not have an exponential dichotomy. The proof of our next theorem follows from Theorem 5.5 and Theorem 5.6.

THEOREM 5.11. *Assume that (a, \bar{b}) belongs to the shaded region shown in Figure 5.2. Then the following holds.*

- (i) *If the function f is bounded, then there exists exactly one bounded mild solution w of (5.26).*
- (ii) *If $f(\cdot, x) \in C_0(\mathbb{R})$ for every $x \in [0, 2\pi]$, then there exists exactly one mild solution w of (5.26) such that $\lim_{t \rightarrow \pm\infty} w(t, x) = 0$.*
- (iii) *If $f(\cdot, x)$ is almost periodic uniformly for $x \in [0, 2\pi]$, then there exists exactly one mild solution w of (5.26) such that $w(\cdot, x)$ is almost periodic uniformly for $x \in [0, 2\pi]$.*

Proposition 5.10 and **Theorem 5.7** lead to conditions for the existence of a unique periodic mild solution.




FIGURE 5.2.

THEOREM 5.12. *If $f(\cdot, x) \in P_p(\mathbb{R})$ for all $x \in [0, 2\pi]$ and $\bar{b} + a \neq n^2$ for all $n \in \{1, 2, 3, \dots\}$, then there exists exactly one mild solution w of (5.26) such that $w(\cdot, x) \in P_p(\mathbb{R})$.*

Acknowledgement

Most of the work on this paper was done while the authors were visiting the Department of Mathematics at the University of Missouri-Columbia. The authors are most grateful for the warm hospitality they experienced during their visit. Support of the first author by “Friedrich-Ebert-Stiftung” is gratefully acknowledged.

References

- [1] H. Amann, *Gewöhnliche Differentialgleichungen [Ordinary Differential Equations]*, de Gruyter Lehrbuch. [de Gruyter Textbook] Walter de Gruyter & Co., Berlin, 1983 (German). [MR 85f:34001](#).
- [2] B. Amir and L. Maniar, *Asymptotic behaviour of hyperbolic Cauchy problems for Hille-Yosida operators with an application to retarded differential equations*, to appear in *Quaestiones Math.*
- [3] W. Arendt and C. J. K. Batty, *Almost periodic solutions of first- and second-order Cauchy problems*, *J. Differential Equations* **137** (1997), no. 2, 363–383. [MR 98g:34099](#). [Zbl 879.34046](#).
- [4] B. Aulbach and N. Van Minh, *Nonlinear semigroups and the existence and stability of solutions of semilinear nonautonomous evolution equations*, *Abstr. Appl. Anal.* **1** (1996), no. 4, 361–380. [MR 98k:47135](#). [Zbl 934.34051](#).
- [5] A. G. Baskakov, *Semigroups of difference operators in spectral analysis of linear differential operators*, *Funct. Anal. Appl.* **30** (1996), no. 3, 149–157, translation from [Funktsional. Anal. i Prilozhen. **30** (1996), no. 3, 1–11. [MR 98b:47053](#)]. [Zbl 911.47040](#).

- [6] C. J. K. Batty, W. Hutter, and F. Räbiger, *Almost periodicity of mild solutions of inhomogeneous periodic Cauchy problems*, J. Differential Equations **156** (1999), no. 2, 309–327. [CMP 1 705 399](#). [Zbl 934.34049](#).
- [7] C. Chicone and Y. Latushkin, *Evolution Semigroups in Dynamical Systems and Differential Equations*, Mathematical Surveys and Monographs, vol. 70, American Mathematical Society, Providence, R.I., 1999. [CMP 1 707 332](#). [Zbl 990.38381](#).
- [8] P. Clément, O. Diekmann, M. Gyllenberg, H. J. A. M. Heijmans, and H. R. Thieme, *Perturbation theory for dual semigroups. I. The sun-reflexive case*, Math. Ann. **277** (1987), no. 4, 709–725. [MR 88m:47071](#). [Zbl 634.47039](#).
- [9] ———, *Perturbation theory for dual semigroups. II. Time-dependent perturbations in the sun-reflexive case*, Proc. Roy. Soc. Edinburgh Sect. A **109** (1988), no. 1-2, 145–172. [MR 89i:47072](#). [Zbl 661.47015](#).
- [10] G. Da Prato and A. Lunardi, *Solvability on the real line of a class of linear Volterra integro-differential equations of parabolic type*, Ann. Mat. Pura Appl. (4) **150** (1988), 67–117. [MR 89g:45024](#). [Zbl 646.45013](#).
- [11] G. Da Prato and E. Sinestrari, *Differential operators with nondense domain*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **14** (1987), no. 2, 285–344. [MR 89f:47062](#). [Zbl 652.34069](#).
- [12] J. L. Dalec'kii and M. G. Krein, *Stability of Solutions of Differential Equations in Banach Space*, translations of Mathematical Monographs, vol. 43, American Mathematical Society, Providence, R.I., 1974, translated from the Russian by S. Smith. [MR 50#5126](#). [Zbl 286.34094](#).
- [13] W. Desch, W. Schappacher, and K. P. Zhang, *Semilinear evolution equations*, Houston J. Math. **15** (1989), no. 4, 527–552. [MR 92d:47087](#). [Zbl 712.47052](#).
- [14] K.-J. Engel and R. Nagel, *One-parameter Semigroups for Linear Evolution Equations*, Graduate Texts in Mathematics, vol. 194, Springer-Verlag, New York, 2000, with contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt. [MR 2000i:47075](#). [Zbl 990.64050](#).
- [15] W. E. Fitzgibbon, *Stability for abstract nonlinear Volterra equations involving finite delay*, J. Math. Anal. Appl. **60** (1977), no. 2, 429–434. [MR 57#3765](#).
- [16] G. Gühring, F. Räbiger, and W. M. Ruess, *Linearized stability for semilinear non-autonomous evolution equations with applications to retarded differential equations*, Differential Integral Equations **13** (2000), no. 4-6, 503–527. [CMP 1 750 038](#).
- [17] J. K. Hale and S. M. Verduyn Lunel, *Introduction to Functional-differential Equations*, Applied Mathematical Sciences, vol. 99, Springer-Verlag, New York, 1993. [MR 94m:34169](#). [Zbl 787.34002](#).
- [18] D. Henry, *Geometric Theory of Semilinear Parabolic Equations*, Lecture Notes in Mathematics, vol. 840, Springer-Verlag, Berlin, 1981. [MR 83j:35084](#). [Zbl 456.35001](#).
- [19] W. Hutter, *Spectral theory and almost periodicity of mild solutions of non-autonomous Cauchy problems*, Ph.d. thesis, Univ. Tuebingen, Mathematische Fakultaet, Tuebingen, 1997. [Zbl 912.47022](#).
- [20] Y. Katznelson, *An Introduction to Harmonic Analysis*, 2nd corr. ed., Dover Publications Inc., New York, 1976. [MR 54#10976](#). [Zbl 352.43001](#).
- [21] Y. Latushkin and S. Montgomery-Smith, *Evolutionary semigroups and Lyapunov theorems in Banach spaces*, J. Funct. Anal. **127** (1995), no. 1, 173–197. [MR 96k:47072](#). [Zbl 878.47024](#).
- [22] Y. Latushkin, T. Randolph, and R. Schnaubelt, *Exponential dichotomy and mild solutions of nonautonomous equations in Banach spaces*, J. Dynam. Differential Equations **10** (1998), no. 3, 489–510. [MR 99g:34123](#). [Zbl 908.34045](#).
- [23] B. M. Levitan and V. V. Zhikov, *Almost Periodic Functions and Differential Equations*, Cambridge University Press, Cambridge, 1982, translated from the Russian by L. W. Longdon. [MR 84g:34004](#). [Zbl 499.43005](#).

- [24] J. L. Massera and J. J. Schäffer, *Linear Differential Equations and Function Spaces*, Pure and Applied Mathematics, vol. 21, Academic Press, New York, 1966. [MR 35#3197](#). [Zbl 243.34107](#).
- [25] R. Nagel, *Semigroup methods for nonautonomous Cauchy problems*, Evolution Equations (Baton Rouge, LA, 1992) (G. Ferreyra et al., eds.), Lecture Notes in Pure and Appl. Math., vol. 168, Dekker, New York, 1994, pp. 301–316. [MR 96d:47042](#). [Zbl 811.34047](#).
- [26] R. Nagel and E. Sinestrari, *Inhomogeneous Volterra integrodifferential equations for Hille-Yosida operators*, Functional Analysis (Essen, 1991), Lecture Notes in Pure and Appl. Math., vol. 150, Dekker, New York, 1994, pp. 51–70. [MR 94i:34121](#).
- [27] T. Naito and N. Van Minh, *Evolution semigroups and spectral criteria for almost periodic solutions of periodic evolution equations*, J. Differential Equations **152** (1999), no. 2, 358–376. [MR 99m:34131](#). [Zbl 924.34038](#).
- [28] M. E. Parrott, *Linearized stability and irreducibility for a functional-differential equation*, SIAM J. Math. Anal. **23** (1992), no. 3, 649–661. [MR 93f:34134](#). [Zbl 763.34058](#).
- [29] G. Pecelli, *Functional-differential equations: dichotomies, perturbations, and admissibility*, J. Differential Equations **16** (1974), 72–102. [MR 56#9043](#). [Zbl 311.34090](#).
- [30] Vũ Quoc Phóng, *Stability and almost periodicity of trajectories of periodic processes*, J. Differential Equations **115** (1995), no. 2, 402–415. [MR 95k:34062](#). [Zbl 815.34050](#).
- [31] J. Prüss, *Bounded solutions of Volterra equations*, SIAM J. Math. Anal. **19** (1988), no. 1, 133–149. [MR 89a:45027](#). [Zbl 642.45005](#).
- [32] ———, *Evolutionary Integral Equations and Applications*, Monographs in Mathematics, vol. 87, Birkhäuser Verlag, Basel, 1993. [MR 94h:45010](#). [Zbl 784.45006](#).
- [33] A. Rhondi, *Extrapolation methods to solve non-autonomous retarded partial differential equations*, Studia Math. **126** (1998), no. 3, 219–233. [MR 99c:47058](#). [Zbl 907.47041](#).
- [34] W. M. Ruess, *Existence of solutions to partial functional-differential equations with delay*, Theory and Applications of Nonlinear Operators of Accretive and Monotone Type (A. G. Kartsatos, ed.), Lecture Notes in Pure and Appl. Math., vol. 178, Dekker, New York, 1996, pp. 259–288. [MR 97b:34085](#). [Zbl 859.35132](#).
- [35] W. M. Ruess and Vũ Quoc Phóng, *Asymptotically almost periodic solutions of evolution equations in Banach spaces*, J. Differential Equations **122** (1995), no. 2, 282–301. [MR 96i:34143](#). [Zbl 837.34067](#).
- [36] R. J. Sacker and G. R. Sell, *Dichotomies for linear evolutionary equations in Banach spaces*, J. Differential Equations **113** (1994), no. 1, 17–67. [MR 96k:34136](#). [Zbl 815.34049](#).
- [37] J. J. Schäffer, *Linear differential equations with delays: Admissibility and conditional exponential stability. II*, J. Differential Equations **10** (1971), 471–484. [MR 45#686](#).
- [38] R. Schnaubelt, *Sufficient conditions for exponential stability and dichotomy of evolution equations*, Forum Math. **11** (1999), no. 5, 543–566. [CMP 1 705 902](#). [Zbl 936.34038](#).
- [39] H. R. Thieme, *Semiflows generated by Lipschitz perturbations of non-densely defined operators*, Differential Integral Equations **3** (1990), no. 6, 1035–1066. [MR 92e:47121](#). [Zbl 734.34059](#).
- [40] C. C. Travis and G. F. Webb, *Existence and stability for partial functional differential equations*, Trans. Amer. Math. Soc. **200** (1974), 395–418. [MR 52#3690](#). [Zbl 299.35085](#).

GABRIELE GÜHRING: UNIVERSITÄT TÜBINGEN, MATHEMATISCHES INSTITUT, AUF DER MORGENSTELLE 10, 72076 TÜBINGEN, GERMANY

E-mail address: gagu@michelangelo.mathematik.uni-tuebingen.de

FRANK RÄBIGER: UNIVERSITÄT TÜBINGEN, MATHEMATISCHES INSTITUT, AUF DER MORGENSTELLE 10, 72076 TÜBINGEN, GERMANY

E-mail address: fra@michelangelo.mathematik.uni-tuebingen.de

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru