

ON THE EXISTENCE OF A POSITIVE SOLUTION FOR A SEMILINEAR ELLIPTIC EQUATION IN THE UPPER HALF SPACE

CHEN SHAOWEI AND LI YONGQING

Received 14 September 2001

We show the existence of a nontrivial solution to the semilinear elliptic equation $-\Delta u + u = b(x)|u|^{p-2}u$, $u > 0$, $u \in H_0^1(\mathbb{R}_+^N)$ under some suitable conditions.

1. Introduction

In this paper, we study the following problem:

$$-\Delta u + u = b(x)|u|^{p-2}u, \quad u > 0, \quad u \in H_0^1(\mathbb{R}_+^N), \quad (1.1)$$

where $2 < p < 2^*$, $2^* = 2N/(N-2)$ when $N > 2$ and $2^* = +\infty$ when $N = 2$, $\mathbb{R}_+^N = \{(x_1, x_2, \dots, x_N) \mid x_N > 0\}$ is the upper half space of \mathbb{R}^N . We put the following conditions on b :

(B₁) $b \in C(\mathbb{R}_+^N)$, and

$$1 = \inf_{x \in \mathbb{R}_+^N} b(x) = \lim_{|x| \rightarrow +\infty, x \in \mathbb{R}_+^N} b(x); \quad (1.2)$$

(B₂) there exists $\delta \in (0, 1)$, such that

$$\lim_{|x| \rightarrow \infty} (b(x) - 1)e^{(1-\delta)|x|} = +\infty. \quad (1.3)$$

Our main result is the following result.

THEOREM 1.1. *If b satisfies (B₁) and (B₂), then (1.1) has a nontrivial solution.*

Similar problems have been studied extensively. In [3], Esteban and Lions showed that $-\Delta u + u = |u|^{p-2}u$, $u \in H_0^1(\mathbb{R}_+^N)$ does not have nontrivial solution.

However, with the help of the potential $b(x)$ and the comparison technique derived in [2], we proved the existence of a nontrivial solution to problem (1.1).

2. Proof of Theorem 1.1

Let

$$\varphi(u) = \int_{\mathbb{R}_+^N} \left(\frac{1}{2} |\nabla u|^2 + \frac{1}{2} |u|^2 \right) dx - \frac{1}{p} \int_{\mathbb{R}_+^N} b(x) |u|^p dx. \quad (2.1)$$

LEMMA 2.1. Denote $S_p = \inf_{u \in H_0^1(\mathbb{R}_+^N), |u|_p=1} \|u\|^2$. Then

$$\inf_{u \in H_0^1(\mathbb{R}_+^N), |u|_p=1} \|u\|^2 = S_p. \quad (2.2)$$

LEMMA 2.2. If $0 < c < c^* := (1/2 - 1/p)S_p^{p/(p-2)}$, then φ satisfies $(PS)_c$ condition.

One can see [2] or [5] for the proof of Lemmas 2.1 and 2.2.

LEMMA 2.3. The equation $-\Delta u + u = |u|^{p-2}u$ has a unique positive solution w in \mathbb{R}^N and satisfies the following conditions:

- (i) $w > 0$ in \mathbb{R}^N , $w \in C^\infty(\mathbb{R}^N)$.
- (ii) $w(x) = w(|x|)$.
- (iii) $w(r)r^{(N-1)/2} \exp(r) \rightarrow C > 0$ and $w'(r)r^{(N-1)/2} \exp(r) \rightarrow -C$, where $r = |x| \rightarrow +\infty$.

Proof. See [1, 4]. □

Proof of Theorem 1.1. By Lemma 2.2, we need only to prove that the Mountain Pass value of φ ,

$$c = \inf \left\{ \left(\frac{1}{2} - \frac{1}{p} \right) \left(\frac{\|v\|^2}{\left(\int_{\mathbb{R}_+^N} b(x) |v|^p dx \right)^{2/p}} \right)^{p/(p-2)} ; v \in H_0^1(\mathbb{R}_+^N), v \neq \theta \right\}, \quad (2.3)$$

satisfies that $0 < c < c^*$.

Set

$$\psi_n(x) = \begin{cases} 1, & |x| \leq n-1, \\ n-|x|, & n-1 < |x| \leq n, \\ 0, & |x| > n. \end{cases} \quad (2.4)$$

Let $x_n = (0, 0, \dots, 0, n)$, $u_n = \psi_n(\cdot - x_n)w(\cdot - x_n)$. Then $u_n \in H_0^1(\mathbb{R}_+^N)$, $n = 1, 2, \dots$.

Notice that

$$\begin{aligned}
\|u_n\|^2 &= \int_{\mathbb{R}^N} |\nabla u_n|^2 dx + \int_{\mathbb{R}^N} u_n^2 dx \\
&= \int_{\mathbb{R}^N} |\nabla(\psi_n(\cdot-x_n)w(\cdot-x_n))|^2 dx + \int_{\mathbb{R}^N} |\psi_n(\cdot-x_n)w(\cdot-x_n)|^2 dx \\
&= \int_{B(\theta,n)} |\nabla(\psi_n \cdot w)|^2 dx + \int_{B(\theta,n)} |\psi_n \cdot w|^2 dx \\
&= \int_{B(\theta,n-1)} |\nabla \omega|^2 dx + \int_{B(\theta,n-1)} \omega^2 dx \\
&\quad + \int_{B_{n-1,n}} \psi_n^2 \cdot |\nabla \omega|^2 dx + \int_{B_{n-1,n}} |\nabla \psi_n|^2 \cdot \omega^2 dx \\
&\quad + 2 \int_{B_{n-1,n}} \omega \cdot \psi_n \cdot \nabla \omega \cdot \nabla \psi_n dx + \int_{B_{n-1,n}} \psi_n^2 \cdot \omega^2 dx \\
&= \int_{\mathbb{R}^N} |\nabla \omega|^2 dx + \int_{\mathbb{R}^N} \omega^2 dx \\
&\quad + \int_{\mathbb{R}^N \setminus B(\theta,n-1)} |\nabla \omega|^2 dx + \int_{\mathbb{R}^N \setminus B(\theta,n-1)} |\omega|^2 dx \\
&\quad + \int_{B_{n-1,n}} \psi_n^2 \cdot |\nabla \omega|^2 dx + 2 \int_{B_{n-1,n}} \omega \cdot \psi_n \cdot \nabla \omega \cdot \nabla \psi_n dx \\
&\quad + \int_{B_{n-1,n}} |\nabla \psi_n|^2 \cdot \omega^2 dx + \int_{B_{n-1,n}} \psi_n^2 \cdot \omega^2 dx,
\end{aligned} \tag{2.5}$$

where $B_{n-1,n} = B(\theta, n) \setminus B(\theta, n-1)$.

From [Lemma 2.3](#) we know that, when n is big enough,

$$\begin{aligned}
\int_{\mathbb{R}^N \setminus B(\theta,n-1)} |\nabla \omega|^2 dx &= C_1 \int_{n-1}^{+\infty} |\omega'(r)|^2 \cdot r^{N-1} dr \\
&\leq C'_1 \int_{n-1}^{+\infty} e^{-2r} dr = \frac{C'_1}{e^{2(n-1)}}.
\end{aligned} \tag{2.6}$$

Similarly, when n is big enough, we get

$$\int_{\mathbb{R}^N \setminus B(\theta,n-1)} \omega^2 dx \leq \frac{C'_2}{e^{2(n-1)}}. \tag{2.7}$$

So

$$\|u_n\|^2 = \|\omega\|^2 + O\left(\frac{1}{e^{2(n-1)}}\right). \tag{2.8}$$

32 Positive solutions in upper half space

Let $a(x) = b(x) - 1$. Then

$$\begin{aligned}
\int_{\mathbb{R}_+^N} b(x)|u_n|^p dx &= \int_{\mathbb{R}_+^N} |u_n|^p dx + \int_{\mathbb{R}_+^N} a(x)|u_n|^p dx \\
&= \int_{B(\theta, n-1)} |\omega|^p dx + \int_{B(\theta, n-1)} a(x+x_n)|\omega|^p dx \\
&\quad + \int_{B_{n-1, n}} a(x+x_n)|\psi_n(x) \cdot \omega(x)|^p dx \\
&= \int_{\mathbb{R}^N} |\omega|^p dx + \int_{B(\theta, n-1)} a(x+x_n)|\omega|^p dx \\
&\quad + \int_{B_{n-1, n}} a(x+x_n)|\psi_n(x) \cdot \omega(x)|^p dx + \int_{\mathbb{R}^N \setminus B(\theta, n-1)} |\omega|^p dx.
\end{aligned} \tag{2.9}$$

Thus

$$\begin{aligned}
\int_{\mathbb{R}^N \setminus B(\theta, n-1)} |\omega|^p dx &= o\left(\frac{1}{e^{2(n-1)}}\right), \\
\int_{B_{n-1, n}} a(x+x_n)|\psi_n(x) \cdot \omega(x)|^p dx &= o\left(\frac{1}{e^{2(n-1)}}\right).
\end{aligned} \tag{2.10}$$

From (B₂), we see that there exists a positive constant λ , such that

$$a(x) \geq \frac{\lambda}{e^{(1-\delta)|x|}}. \tag{2.11}$$

So there exists a positive constant C_3 , such that

$$\int_{\mathbb{R}_+^N} b(x)|u_n|^p dx \geq \int_{\mathbb{R}^N} |\omega|^p dx + \frac{C_3}{e^{2(1-\delta)(n-1)}}. \tag{2.12}$$

Then

$$\begin{aligned}
&\left(\int_{\mathbb{R}_+^N} b(x)|u_n|^p dx \right)^{2/p} \\
&\geq \left(\int_{\mathbb{R}^N} |\omega|^p dx + \frac{C_3}{e^{2(1-\delta)(n-1)}} \right)^{2/p} \\
&= \left(\int_{\mathbb{R}^N} |\omega|^p dx \right)^{2/p} \left(1 + \frac{C_4}{e^{2(1-\delta)(n-1)}} \right)^{2/p} \\
&= \left(\int_{\mathbb{R}^N} |\omega|^p dx \right)^{2/p} \left(1 + \frac{C_5}{e^{2(1-\delta)(n-1)}} + o\left(\frac{1}{e^{2(1-\delta)(n-1)}}\right) \right).
\end{aligned} \tag{2.13}$$

Thus

$$\frac{\|u_n\|^2}{\left(\int_{\mathbb{R}_+^N} b(x)|u_n|^p dx\right)^{2/p}} = \frac{\|\omega\|^2}{\left(\int_{\mathbb{R}^N} \omega^p dx\right)^{2/p}} - \frac{C_6}{e^{2(1-\delta)(n-1)}} + o\left(\frac{1}{e^{2(1-\delta)(n-1)}}\right). \quad (2.14)$$

So

$$\frac{\|u_n\|^2}{\left(\int_{\mathbb{R}_+^N} b(x)|u_n|^p dx\right)^{2/p}} < \frac{\|\omega\|^2}{\left(\int_{\mathbb{R}^N} \omega^p dx\right)^{2/p}}, \quad (2.15)$$

when n is big enough.

Thus, we come to the conclusion that

$$c < \left(\frac{1}{2} - \frac{1}{p}\right) S_p^{p/(p-2)}. \quad (2.16)$$

□

Acknowledgement

This work was supported in part by the National Natural Science Foundation and the Fujian Province Natural Science Foundation of China.

References

- [1] H. Berestycki and P.-L. Lions, *Nonlinear scalar field equations. I. Existence of a ground state*, Arch. Rational Mech. Anal. **82** (1983), no. 4, 313–345.
- [2] W. Y. Ding and W. M. Ni, *On the existence of positive entire solutions of a semilinear elliptic equation*, Arch. Rational Mech. Anal. **91** (1986), no. 4, 283–308.
- [3] M. J. Esteban and P.-L. Lions, *Existence and nonexistence results for semilinear elliptic problems in unbounded domains*, Proc. Roy. Soc. Edinburgh Sect. A **93** (1982/83), no. 1-2, 1–14.
- [4] B. Gidas, W. M. Ni, and L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), no. 3, 209–243.
- [5] M. Willem, *Minimax Theorems*, Progress in Nonlinear Differential Equations and Their Applications, vol. 24, Birkhäuser, Massachusetts, 1996.

CHEN SHAOWEI: DEPARTMENT OF MATHEMATICS, FUJIAN NORMAL UNIVERSITY, FUZHOU, FUJIAN 35007, CHINA

LI YONGQING: DEPARTMENT OF MATHEMATICS, FUJIAN NORMAL UNIVERSITY, FUZHOU, FUJIAN 35007, CHINA

E-mail address: yqli@pub1.fz.fj.cn

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk