

# ISOLATION AND SIMPLICITY FOR THE FIRST EIGENVALUE OF THE $p$ -LAPLACIAN WITH A NONLINEAR BOUNDARY CONDITION

SANDRA MARTÍNEZ AND JULIO D. ROSSI

Received 21 May 2001

We prove the simplicity and isolation of the first eigenvalue for the problem  $\Delta_p u = |u|^{p-2} u$  in a bounded smooth domain  $\Omega \subset \mathbb{R}^N$ , with a nonlinear boundary condition given by  $|\nabla u|^{p-2} \partial u / \partial \nu = \lambda |u|^{p-2} u$  on the boundary of the domain.

## 1. Introduction

In this paper, we study the first eigenvalue for the following problem:

$$\begin{aligned} \Delta_p u &= |u|^{p-2} u && \text{in } \Omega, \\ |\nabla u|^{p-2} \frac{\partial u}{\partial \nu} &= \lambda |u|^{p-2} u && \text{on } \partial \Omega. \end{aligned} \tag{1.1}$$

Here  $\Omega$  is a bounded domain in  $\mathbb{R}^N$  with smooth boundary,  $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$  is the  $p$ -Laplacian, and  $\partial/\partial \nu$  is the outer normal derivative. In the linear case, that is for  $p = 2$ , this eigenvalue problem is known as the *Steklov* problem (see [3]).

Problems of the form (1.1) appear in a natural way when one considers the Sobolev trace inequality. In fact, the immersion  $W^{1,p}(\Omega) \hookrightarrow L^p(\partial \Omega)$  is compact, hence there exists a constant  $\lambda_1$  such that

$$\lambda_1^{1/p} \|u\|_{L^p(\partial \Omega)} \leq \|u\|_{W^{1,p}(\Omega)}. \tag{1.2}$$

The extremals (functions where the constant is attained) are solutions of (1.1). This Sobolev trace constant  $\lambda_1$  can be characterized as

$$\lambda_1 = \inf_{u \in W^{1,p}(\Omega)} \left\{ \int_{\Omega} |\nabla u|^p + |u|^p dx, \int_{\partial \Omega} |u|^p = 1 \right\}, \tag{1.3}$$

and is the first eigenvalue of (1.1) in the sense that  $\lambda_1 \leq \lambda$  for any other eigenvalue  $\lambda$ .

In [13] it is proved that, there exists a sequence of eigenvalues  $\lambda_n$  of (1.1) such that  $\lambda_n \rightarrow +\infty$  as  $n \rightarrow +\infty$ . This is done using standard variational arguments together with the Sobolev trace immersion that provide the necessary compactness. Indeed, for solutions of (1.1) we can understand critical points of the associated energy functional

$$\mathcal{F}(u) = \frac{1}{p} \int_{\Omega} |\nabla u|^p + \frac{1}{p} \int_{\Omega} |u|^p - \frac{\lambda}{p} \int_{\partial\Omega} |u|^p. \quad (1.4)$$

This functional  $\mathcal{F}$  is well defined and  $C^1$  in  $W^{1,p}(\Omega)$  and the usual min-max techniques can be applied (see [13]). Also see [14] for similar results for the  $p$ -Laplacian with Dirichlet boundary conditions.

We prove the following result.

**THEOREM 1.1.** *The first eigenvalue  $\lambda_1$  is isolated and simple.*

We remark that this theorem says that the extremals of the Sobolev trace inequality are unique up to multiplication by a real number. In the special case of a ball,  $\Omega = B(0, R)$ , our result implies that the first eigenfunction is radial. In fact, if  $u_1(x)$  is an eigenfunction associated to  $\lambda_1$  and  $R(x)$  is any rotation, then  $u_1(R(x))$  is also an eigenfunction, by our result we have that  $u_1(x) = u_1(R(x))$ . We conclude that  $u_1$  must be radial. Also from our results it follows that any other eigenvalue has nonradial eigenfunctions as they have to change sign on the boundary (see Lemma 2.4).

The study of the eigenvalue problem when the nonlinear term is placed in the equation, that is, when one considers a quasilinear problem of the form  $-\Delta_p u = \lambda|u|^{p-2}u$  with Dirichlet boundary conditions, has received considerable attention (cf. [1, 2, 15, 14, 17], etc.).

However, nonlinear boundary conditions have only been considered in recent years. For the Laplace operator with nonlinear boundary conditions (cf. [5, 6, 8, 16, 19]). For elliptic systems with nonlinear boundary conditions (see [11, 12]). For previous work for the  $p$ -Laplacian with nonlinear boundary conditions of different type see [7, 13, 18]. Also, one is led to nonlinear boundary conditions in the study of conformal deformations on Riemannian manifolds with boundary (cf. [4, 9, 10]).

## 2. Proof of the main result

In this section, we prove that the first eigenvalue  $\lambda_1$  is isolated and simple. To clarify the exposition, we will divide the proof in several lemmas.

**LEMMA 2.1.** *Let  $u_1$  be an eigenfunction with eigenvalue  $\lambda_1$ , then  $u_1$  does not change sign on  $\Omega$ . Moreover, if  $u_1$  is  $C^{1,\alpha}$ , it does not vanish on  $\bar{\Omega}$ .*

*Proof.* We have that  $|u_1|$  is also a minimizer of (1.3). By the maximum principle (see [20]) we have that  $|u_1| > 0$  in  $\Omega$ . Assume that  $u_1$  is regular and that there exists  $x_0 \in \partial\Omega$  such that  $u_1(x_0) = 0$ , by the Hopf lemma (see [20]) we have that the normal derivative has strict sign,  $(\partial|u_1|/\partial\nu)(x_0) < 0$ , but the boundary condition imposes  $(\partial|u_1|/\partial\nu)(x_0) = 0$ , a contradiction which proves that  $|u_1| > 0$  in  $\bar{\Omega}$ . The result follows.  $\square$

Now we state an auxiliary lemma,

LEMMA 2.2. (a) Let  $p \geq 2$ , then for all  $\xi_1, \xi_2 \in \mathbb{R}^N$

$$|\xi_2|^p \geq |\xi_1|^p + p|\xi_1|^{p-2}\langle \xi_1, \xi_2 - \xi_1 \rangle + C(p)|\xi_1 - \xi_2|^p. \quad (2.1)$$

(b) Let  $p < 2$ , then for all  $\xi_1, \xi_2 \in \mathbb{R}^N$

$$|\xi_2|^p \geq |\xi_1|^p + p|\xi_1|^{p-2}\langle \xi_1, \xi_2 - \xi_1 \rangle + C(p)\frac{|\xi_1 - \xi_2|^p}{(|\xi_2| + |\xi_1|)^{2-p}}, \quad (2.2)$$

where  $C(p)$  is a constant depending only on  $p$ .

*Proof.* See [17].  $\square$

LEMMA 2.3. The first eigenvalue  $\lambda_1$  is simple. Let  $u, v$  be two eigenfunctions associated with  $\lambda_1$ , then there exists  $c$  such that  $u = cv$ .

*Proof.* By Lemma 2.1, we can assume that  $u, v$  are positive in  $\Omega$ . We perform the following calculations assuming that  $u, v$  are strictly positive in  $\bar{\Omega}$ , to obtain our result we can consider  $u + \varepsilon$  and  $v + \varepsilon$  and let  $\varepsilon \rightarrow 0$  at the end as in [17]. Therefore, we can take  $\eta_1 = (u^p - v^p)/u^{p-1}$  and  $\eta_2 = (v^p - u^p)/v^{p-1}$  as test functions in the weak form of (1.1) satisfied by  $u$  and  $v$ , respectively. We have

$$\begin{aligned} & \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla \left( \frac{u^p - v^p}{u^{p-1}} \right) \\ &= \lambda \int_{\partial\Omega} |u|^{p-2} u \left( \frac{u^p - v^p}{u^{p-1}} \right) - \int_{\Omega} |u|^{p-2} u \left( \frac{u^p - v^p}{u^{p-1}} \right), \\ & \int_{\Omega} |\nabla v|^{p-2} \nabla v \nabla \left( \frac{v^p - u^p}{v^{p-1}} \right) \\ &= \lambda \int_{\partial\Omega} |v|^{p-2} v \left( \frac{v^p - u^p}{v^{p-1}} \right) - \int_{\Omega} |v|^{p-2} v \left( \frac{v^p - u^p}{v^{p-1}} \right). \end{aligned} \quad (2.3)$$

Adding both equations we get

$$0 = \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla \left( \frac{u^p - v^p}{u^{p-1}} \right) + \int_{\Omega} |\nabla v|^{p-2} \nabla v \nabla \left( \frac{v^p - u^p}{v^{p-1}} \right). \quad (2.4)$$

Using

$$\nabla \left( \frac{u^p - v^p}{u^{p-1}} \right) = \nabla u - p \frac{v^{p-1}}{u^{p-1}} \nabla v + (p-1) \frac{v^p}{u^p} \nabla u, \quad (2.5)$$

we obtain that the first term of (2.4) is

$$\begin{aligned} & \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla \left( \frac{u^p - v^p}{u^{p-1}} \right) \\ &= \int_{\Omega} |\nabla u|^p - p \int_{\Omega} \frac{v^{p-1}}{u^{p-1}} |\nabla u|^{p-2} \nabla v \nabla u + \int_{\Omega} (p-1) \frac{v^p}{u^p} |\nabla u|^p \\ &= \int_{\Omega} |\nabla \ln u|^p u^p - p \int_{\Omega} v^p |\nabla \ln u|^{p-2} \langle \nabla \ln u, \nabla \ln v \rangle u v \\ & \quad + \int_{\Omega} (p-1) |\nabla \ln u|^p v^p. \end{aligned} \quad (2.6)$$

We also have an analogous expression for the second term of (2.4). Using both expressions we get that (2.4) becomes

$$\begin{aligned} 0 &= \int_{\Omega} (u^p - v^p) (|\nabla \ln u|^p - |\nabla \ln v|^p) \\ & \quad - p \int_{\Omega} v^p (|\nabla \ln u|^{p-2} \langle \nabla \ln u, \nabla \ln v - \nabla \ln u \rangle) \\ & \quad - p \int_{\Omega} u^p (|\nabla \ln v|^{p-2} \langle \nabla \ln v, \nabla \ln u - \nabla \ln v \rangle). \end{aligned} \quad (2.7)$$

Taking  $\xi_1 = \nabla \ln u$  and  $\xi_2 = \nabla \ln v$  and using Lemma 2.2 we get, for  $p \geq 2$ ,

$$0 \geq \int_{\Omega} C(p) |\nabla \ln u - \nabla \ln v|^p (u^p + v^p). \quad (2.8)$$

Hence,

$$0 = |\nabla \ln u - \nabla \ln v|. \quad (2.9)$$

This implies that  $u = kv$ , as we wanted to prove. For  $p < 2$ , we use the second part of Lemma 2.2 as above.  $\square$

Now we turn our attention to the proof of the isolation of the first eigenvalue, in order to prove this we need the following nodal result.

LEMMA 2.4. *Let  $w$  be an eigenfunction corresponding to  $\lambda \neq \lambda_1$ , then  $w$  changes sign on  $\partial\Omega$ , that is,  $w^+|_{\partial\Omega} \neq 0$  and  $w^-|_{\partial\Omega} \neq 0$ . Moreover, there exists a constant  $C$  such that*

$$|\partial\Omega^+| \geq C\lambda^{-\beta}, \quad |\partial\Omega^-| \geq C\lambda^{-\beta}, \quad (2.10)$$

where  $\partial\Omega^+ = \partial\Omega \cap \{w > 0\}$ ,  $\partial\Omega^- = \partial\Omega \cap \{w < 0\}$ ,  $\beta = (N-1)/(p-1)$  if  $1 < p < N$  and  $\beta = 2$  if  $p \geq N$ . Here  $|A|$  denotes the  $(N-1)$ -dimensional measure of a subset  $A$  of the boundary.

*Proof.* Assume that  $w$  does not change sign in  $\Omega$ , then we can assume that  $w > 0$  in  $\Omega$  using ideas similar to those of [Lemma 2.1](#). Let  $u_1$  be a positive eigenfunction associated to  $\lambda_1$ . Making similar computations as the ones performed in the proof of [Lemma 2.3](#) we arrive at

$$(\lambda_1 - \lambda) \int_{\partial\Omega} (u_1^p - w^p) \geq C \int_{\Omega} |\nabla \ln w - \nabla \ln u_1|^p (u_1^p + w^p) \geq 0. \quad (2.11)$$

Therefore, if we take  $kw$  instead of  $w$  we get that, for every  $k > 0$ , we have

$$\int_{\partial\Omega} (u_1^p - k^p w^p) \leq 0, \quad (2.12)$$

which is a contradiction if we take

$$k^p \left( \int_{\partial\Omega} w^p \right) < \left( \int_{\partial\Omega} u_1^p \right). \quad (2.13)$$

Therefore,  $w$  changes sign in  $\Omega$  and by the maximum principle, [\[20\]](#), also  $w$  changes sign in  $\partial\Omega$ .

We use  $w^-$  as test function in the weak form of [\(1.1\)](#) satisfied by  $w$  to obtain

$$\int_{\Omega} |\nabla w^-|^p + \int_{\Omega} |w^-|^p = \lambda \int_{\partial\Omega \cap \{w < 0\}} |w^-|^p. \quad (2.14)$$

Hence,

$$\|w^-\|_{W^{1,p}(\Omega)}^p \leq \lambda \left( \int_{\partial\Omega} |w^-|^{p\alpha} \right)^{1/\alpha} |\partial\Omega^-|^{1/\beta}. \quad (2.15)$$

If  $1 < p < N$  we choose  $\alpha = (N-1)/(N-p)$  and  $\beta = (N-1)/(p-1)$ . Now we use the trace theorem to get that there exists a constant  $C$  such that

$$\|w^-\|_{L^{p\alpha}(\partial\Omega)}^p \leq C \|w^-\|_{W^{1,p}(\Omega)}^p. \quad (2.16)$$

If  $p \geq N$ , we choose  $\alpha = \beta = 2$  and we argue as before using that  $W^{1,p}(\Omega) \hookrightarrow L^{2p}(\partial\Omega)$ . A similar argument works for  $w^+$ .  $\square$

**LEMMA 2.5.** *Let  $\phi \in W^{1,p}(\Omega)'$ , then there exists a unique weak solution  $u \in W^{1,p}(\Omega)$  of*

$$-\Delta_p u + |u|^{p-2} u = \phi. \quad (2.17)$$

Moreover, the operator  $A_p : \phi \mapsto u$  is continuous.

*Proof.* See [13]. □

With these lemmas we can prove the isolation of  $\lambda_1$ .

LEMMA 2.6. *The first eigenvalue  $\lambda_1$  is isolated, that is, there exists  $a > \lambda_1$  such that  $\lambda_1$  is the unique eigenvalue in  $[0, a]$ .*

*Proof.* From the characterization of  $\lambda_1$ , it is easy to see that  $\lambda_1 \leq \lambda$  for every eigenvalue  $\lambda$ . Assume that  $\lambda_1$  is not isolated, then there exists a sequence  $\lambda_k$  with  $\lambda_k > \lambda_1$ ,  $\lambda_k \searrow \lambda_1$ . Let  $w_k$  be an eigenfunction associated to  $\lambda_k$ , we can assume that  $\|w_k\|_{W^{1,p}(\Omega)} = 1$ . Therefore, we can extract a subsequence (that we still denote by  $w_k$ ) such that  $w_k \rightarrow u_1$  in  $L^p(\partial\Omega)$ . Define  $\phi_k \in (W^{1,p}(\Omega))'$  as

$$\phi_k(u) = \lambda_k \int_{\partial\Omega} |w_k|^{p-2} w_k u \quad (2.18)$$

and  $\phi \in (W^{1,p}(\Omega))'$  by

$$\phi(u) = \lambda_1 \int_{\partial\Omega} |u_1|^{p-2} u_1 u. \quad (2.19)$$

From the  $L^p(\partial\Omega)$  convergence of  $w_k$  to  $u_1$  we get that  $\phi_k$  converges to  $\phi$  in  $(W^{1,p}(\Omega))'$ . Using the continuity of  $A_p$  given by Lemma 2.5 we get that the sequence  $w_k$  converge strongly in  $W^{1,p}(\Omega)$ . Therefore, passing to the limit in the weak form of (1.1) we get that  $u_1$  is an eigenfunction with eigenvalue  $\lambda_1$ . By Lemma 2.1 we can assume that  $u_1 > 0$  on  $\partial\Omega$ . By Egorov's theorem we can find a subset  $A_\varepsilon$  of  $\partial\Omega$  such that  $|A_\varepsilon| < \varepsilon$  and  $w_k \rightarrow u_1 > 0$  uniformly in  $\partial\Omega \setminus A_\varepsilon$ . This contradicts the fact that, by (2.10), we have, for every  $k$

$$|\partial\Omega_k^-| = \partial\Omega \cap \{w_k < 0\} \geq C \lambda_k^{-(N-1)/(p-1)}. \quad (2.20)$$

This completes the proof. □

### Acknowledgments

We want to thank J. Garcia-Azorero and I. Peral for several suggestions and interesting discussions. This work is supported by ANPCyT PICT 00137 and CONICET.

### References

- [1] W. Allegretto and Y. X. Huang, *A Picone's identity for the  $p$ -Laplacian and applications*, Nonlinear Anal. **32** (1998), no. 7, 819–830.
- [2] A. Anane, *Simplicité et isolation de la première valeur propre du  $p$ -laplacien avec poids*, C. R. Acad. Sci. Paris Sér. I Math. **305** (1987), no. 16, 725–728 (French).
- [3] I. Babuška and J. Osborn, *Eigenvalue problems*, Handbook of Numerical Analysis, vol. 2, North-Holland, Amsterdam, 1991, pp. 641–787.
- [4] P. Cherrier, *Problèmes de Neumann non linéaires sur les variétés Riemanniennes*, J. Funct. Anal. **57** (1984), no. 2, 154–206 (French).

- [5] M. Chipot, M. Chlebík, M. Fila, and I. Shafrir, *Existence of positive solutions of a semilinear elliptic equation in  $\mathbb{R}_+^n$  with a nonlinear boundary condition*, J. Math. Anal. Appl. **223** (1998), no. 2, 429–471.
- [6] M. Chipot, I. Shafrir, and M. Fila, *On the solutions to some elliptic equations with nonlinear Neumann boundary conditions*, Adv. Differential Equations **1** (1996), no. 1, 91–110.
- [7] F.-C. Șt. Cîrstea and V. D. Rădulescu, *Existence and non-existence results for a quasilinear problem with nonlinear boundary condition*, J. Math. Anal. Appl. **244** (2000), no. 1, 169–183.
- [8] M. del Pino and C. Flores, *Asymptotic behavior of best constants and extremals for trace embeddings in expanding domains*, Comm. Partial Differential Equations **26** (2001), no. 11–12, 2189–2210.
- [9] J. F. Escobar, *Uniqueness theorems on conformal deformation of metrics, Sobolev inequalities, and an eigenvalue estimate*, Comm. Pure Appl. Math. **43** (1990), no. 7, 857–883.
- [10] ———, *Conformal deformation of a Riemannian metric to a scalar flat metric with constant mean curvature on the boundary*, Ann. of Math. (2) **136** (1992), no. 1, 1–50.
- [11] J. Fernández Bonder, J. P. Pinasco, and J. D. Rossi, *Existence results for Hamiltonian elliptic systems with nonlinear boundary conditions*, Electron. J. Differential Equations **1999** (1999), no. 40, 1–15.
- [12] J. Fernández Bonder and J. D. Rossi, *Existence for an elliptic system with nonlinear boundary conditions via fixed point methods*, Adv. Differential Equations **6** (2001), 1–20.
- [13] ———, *Existence results for the  $p$ -Laplacian with nonlinear boundary conditions*, J. Math. Anal. Appl. **263** (2001), no. 1, 195–223.
- [14] J. García-Azorero and I. Peral, *Existence and nonuniqueness for the  $p$ -Laplacian: nonlinear eigenvalues*, Comm. Partial Differential Equations **12** (1987), 1389–1430.
- [15] ———, *Multiplicity of solutions for elliptic problems with critical exponent or with a nonsymmetric term*, Trans. Amer. Math. Soc. **323** (1991), no. 2, 877–895.
- [16] B. Hu, *Nonexistence of a positive solution of the Laplace equation with a nonlinear boundary condition*, Differential Integral Equations **7** (1994), no. 2, 301–313.
- [17] P. Lindqvist, *On the equation  $\operatorname{div}(|\nabla u|^{p-2}\nabla u) + \lambda|u|^{p-2}u = 0$* , Proc. Amer. Math. Soc. **109** (1990), no. 1, 157–164.
- [18] K. Pflüger, *Existence and multiplicity of solutions to a  $p$ -Laplacian equation with nonlinear boundary condition*, Electron. J. Differential Equations **10** (1998), 1–13.
- [19] S. Terracini, *Symmetry properties of positive solutions to some elliptic equations with nonlinear boundary conditions*, Differential Integral Equations **8** (1995), no. 8, 1911–1922.
- [20] J. L. Vázquez, *A strong maximum principle for some quasilinear elliptic equations*, Appl. Math. Optim. **12** (1984), no. 3, 191–202.

SANDRA MARTÍNEZ: DEPARTAMENTO DE MATEMÁTICA, FCEyN, UBA (1428) BUENOS AIRES, ARGENTINA  
*E-mail address:* [srmartin@bigua.dm.uba.ar](mailto:srmartin@bigua.dm.uba.ar)

JULIO D. ROSSI: DEPARTAMENTO DE MATEMÁTICA, FCEyN, UBA (1428) BUENOS AIRES, ARGENTINA  
*E-mail address:* [jrossi@dm.uba.ar](mailto:jrossi@dm.uba.ar)

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)