

*Research Article*

## Fixed Points of Nonlinear and Asymptotic Contractions in the Modular Space

A. Razani, E. Nabizadeh, M. Beyg Mohamadi, and S. Homaei Pour

Received 24 September 2006; Revised 25 November 2006; Accepted 15 February 2007

Recommended by William Art Kirk

A fixed point theorem for nonlinear contraction in the modular space is proved. Moreover, a fixed point theorem for asymptotic contraction in this space is studied.

Copyright © 2007 A. Razani et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### 1. Introduction

The theory of modular space was initiated by Nakano [1] in connection with the theory of order spaces and was redefined and generalized by Musielak and Orlicz [2]. By defining a norm, particular Banach spaces of functions can be considered. Metric fixed theory for these Banach spaces of functions has been widely studied (see [3]). Another direction is based on considering and abstractly given functional which control the growth of the functions. Even though a metric is not defined, many problems in fixed point theory for nonexpansive mappings can be reformulated in modular spaces.

In this paper, a fixed point theorem for nonlinear contraction in the modular space is proved. Moreover, Kirk's fixed point theorem for asymptotic contraction is presented in this space. In order to do this and for the sake of convenience, some definitions and notations are recalled from [1–6].

*Definition 1.1.* Let  $X$  be an arbitrary vector space over  $K$  ( $= \mathbb{R}$  or  $\mathbb{C}$ ). A functional  $\rho : X \rightarrow [0, +\infty)$  is called modular if

- (1)  $\rho(x) = 0$  if and only if  $x = 0$ ;
- (2)  $\rho(\alpha x) = \rho(x)$  for  $\alpha \in K$  with  $|\alpha| = 1$ , for all  $x, y \in X$ ;
- (3)  $\rho(\alpha x + \beta y) \leq \rho(x) + \rho(y)$  if  $\alpha, \beta \geq 0$ ,  $\alpha + \beta = 1$ , for all  $x, y \in X$ ;

## 2 Abstract and Applied Analysis

*Definition 1.2.* If (3) in Definition 1.1 is replaced by

$$\rho(\alpha x + \beta y) \leq \alpha^s \rho(x) + \beta^s \rho(y), \quad (1.1)$$

for  $\alpha, \beta \geq 0$ ,  $\alpha^s + \beta^s = 1$  with an  $s \in (0, 1]$ , then the modular  $\rho$  is called an  $s$ -convex modular, and if  $s = 1$ ,  $\rho$  is called a convex modular.

*Definition 1.3.* A modular  $\rho$  defines a corresponding modular space, that is, the space  $X_\rho$  given by

$$X_\rho = \{x \in X \mid \rho(\lambda x) \rightarrow 0 \text{ as } \lambda \rightarrow 0\}. \quad (1.2)$$

*Definition 1.4.* Let  $X_\rho$  be a modular space.

- (1) A sequence  $\{x_n\}_n$  in  $X_\rho$  is said to be
  - (a)  $\rho$ -convergent to  $x$  if  $\rho(x_n - x) \rightarrow 0$  as  $n \rightarrow +\infty$ ;
  - (b)  $\rho$ -Cauchy if  $\rho(x_n - x_m) \rightarrow 0$  as  $n, m \rightarrow +\infty$ .
- (2)  $X_\rho$  is  $\rho$ -complete if any  $\rho$ -Cauchy sequence is  $\rho$ -convergent.
- (3) A subset  $B \subset X_\rho$  is said to be  $\rho$ -closed if for any sequence  $\{x_n\}_n \subset B$  with  $x_n \rightarrow x$ , one has  $x \in B$ .  $\overline{B}^\rho$  denotes the closure of  $B$  in the sense of  $\rho$ .
- (4) A subset  $B \subset X_\rho$  is called  $\rho$ -bounded if

$$\delta_\rho(B) = \sup_{x, y \in B} \rho(x - y) < +\infty, \quad (1.3)$$

where  $\delta_\rho(B)$  is called the  $\rho$ -diameter of  $B$ .

- (5) Say that  $\rho$  has Fatou property if

$$\rho(x - y) \leq \liminf \rho(x_n - y_n), \quad (1.4)$$

whenever

$$x_n \xrightarrow{\rho} x, \quad y_n \xrightarrow{\rho} y. \quad (1.5)$$

- (6)  $\rho$  is said to satisfy the  $\Delta_2$ -condition if  $\rho(2x_n) \rightarrow 0$  as  $n \rightarrow +\infty$  whenever  $\rho(x_n) \rightarrow 0$  as  $n \rightarrow +\infty$ .

*Example 1.5.* Let  $(X_\rho, \rho)$  be a modular space, then the function  $d_\rho$  defined on  $X_\rho \times X_\rho$  by

$$d_\rho(x, y) = \begin{cases} 0 & x = y, \\ \rho(x) + \rho(y) & x \neq y, \end{cases} \quad (1.6)$$

is a metric and  $(X_\rho, d_\rho)$  is a metric space.

*Remark 1.6.* Let  $(X_\rho, d_\rho)$  be a metric space which is given in Example 1.5 and let  $\{x_n\}$  be a Cauchy sequence in it. This means that

$$d_\rho(x_n, x_m) \rightarrow 0 \quad \text{as } n, m \rightarrow \infty, \quad (1.7)$$

hence

$$\rho(x_n) + \rho(x_m) \rightarrow 0 \quad \text{as } n, m \rightarrow \infty, \quad (1.8)$$

and this shows that

$$\rho(x_n) \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (1.9)$$

Therefore

$$d_\rho(x_n, 0) \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (1.10)$$

and this proves that  $(X_\rho, d_\rho)$  is a complete metric space. In addition, it implies that all nonconstant sequences for large indices that are convergent must be convergent to zero.

**THEOREM 1.7.** *Suppose that  $(X_\rho, \rho)$  is a modular space and  $T : X_\rho \rightarrow X_\rho$  satisfies the following condition:*

$$\rho(T(x)) + \rho(T(y)) \leq \psi(\rho(x) + \rho(y)) \quad (1.11)$$

for all  $x, y \in X_\rho$ , where  $\psi : \overline{P} \rightarrow [0, \infty)$  is upper semicontinuous from the right on  $\overline{P}$  and for all  $t \in \overline{P} - \{0\}$ ,  $\psi(t) < t$  and

$$P = \{0\} \cup \{\rho(x) + \rho(y) \mid x, y \in X_\rho, x \neq y\}. \quad (1.12)$$

Then 0 is the only fixed point of  $T$ .

*Proof.* We use the metric  $d_\rho$  and note that the closure of  $P$  which is denoted by  $\overline{P}$  is with respect to metric  $d_\rho$ . This metric and the mapping  $T$  satisfy the conditions of [7, Theorem 1], so the proof is complete.  $\square$

## 2. A fixed point of nonlinear contraction

The Banach contraction mapping principle shows the existence and uniqueness of a fixed point in a complete metric space. This has been generalized by many mathematicians such as Arandelović [8], Edelstein [9], Ćirić [10], Rakotch [11], Reich [12], Kirk [13], and so forth. In addition, Boyd and Wong [7] studied mappings which are nonlinear contractions in the metric space. It is necessary to mention that the applications of contraction, generalized contraction principle for self-mappings, and the applications of nonlinear contractions are well known. In this section, an existence fixed point theorem for nonlinear contractions in modular spaces is proved as follows.

**THEOREM 2.1.** *Let  $X_\rho$  be a  $\rho$ -complete modular space, where  $\rho$  satisfies the  $\Delta_2$ -condition. Assume that  $\psi : \mathbb{R}^+ \rightarrow [0, \infty)$  is an increasing and upper semicontinuous function satisfying*

$$\psi(t) < t, \quad \forall t > 0. \quad (2.1)$$

## 4 Abstract and Applied Analysis

Let  $B$  be a  $\rho$ -closed subset of  $X_\rho$  and  $T : B \rightarrow B$  a mapping such that there exist  $c, l \in \mathbb{R}^+$  with  $c > l$ ,

$$\rho(c(Tx - Ty)) \leq \psi(\rho(l(x - y))) \quad (2.2)$$

for all  $x, y \in B$ . Then  $T$  has a fixed point.

*Proof.* Let  $x \in X_\rho$ . At first, we show that the sequence  $\{\rho(c(T^n x - T^{n-1} x))\}$  converges to 0. For  $n \in \mathbb{N}$ , we have

$$\begin{aligned} \rho(c(T^n x - T^{n-1} x)) &\leq \psi(\rho(l(T^{n-1} x - T^{n-2} x))) \\ &< \rho(l(T^{n-1} x - T^{n-2} x)) < \rho(c(T^{n-1} x - T^{n-2} x)). \end{aligned} \quad (2.3)$$

Consequently,  $\{\rho(c(T^n x - T^{n-1} x))\}$  is decreasing and bounded from below ( $\rho(x) \geq 0$ ). Therefore,  $\{\rho(c(T^n x - T^{n-1} x))\}$  converges to  $a$ .

Now, if  $a \neq 0$ ,

$$\begin{aligned} a &= \lim_{n \rightarrow \infty} \rho(c(T^n x - T^{n-1} x)) \leq \lim_{n \rightarrow \infty} \psi(\rho(l(T^{n-1} x - T^{n-2} x))) \\ &\leq \lim_{n \rightarrow \infty} \psi(\rho(c(T^{n-1} x - T^{n-2} x))), \end{aligned} \quad (2.4)$$

then

$$a \leq \psi(a), \quad (2.5)$$

which is a contradiction, so  $a = 0$ .

Now, we show that  $\{T^n x\}$  is a  $\rho$ -Cauchy sequence for  $x \in X_\rho$ . Suppose that  $\{lT^n x\}$  is not a  $\rho$ -Cauchy sequence. Then, there are an  $\epsilon > 0$  and sequences of integers  $\{m_k\}$ ,  $\{n_k\}$ , with  $m_k > n_k \geq k$ , and such that

$$d_k = \rho(l(T^{m_k} x - T^{n_k} x)) \geq \epsilon \quad \text{for } k = 1, 2, \dots \quad (2.6)$$

We can assume that

$$\rho(l(t^{m_k-1} x - t^{n_k} x)) < \epsilon. \quad (2.7)$$

Let  $m_k$  be the smallest number exceeding  $n_k$  for which (2.6) holds, and

$$\Sigma_k = \{m \in \mathbb{N} \mid \exists n_k \in \mathbb{N}; \rho(l(T^m x - T^{n_k} x)) \geq \epsilon, m > n_k \geq k\}. \quad (2.8)$$

Obviously,  $\Sigma_k \neq \emptyset$  and since  $\Sigma_k \subset \mathbb{N}$ , then by Well ordering principle, the minimum element of  $\Sigma_k$  is denoted by  $m_k$ , and clearly (2.7) holds.

Now, let  $\alpha_0 \in \mathbb{R}^+$  be such that  $l/c + 1/\alpha_0 = 1$ , then we have

$$\begin{aligned}
d_k &= \rho(l(T^{m_k}x - T^{n_k}x)) = \rho\left(\frac{lc}{c}(T^{m_k}x - T^{n_k+1}x + T^{n_k+1}x - T^{n_k}x)\right) \\
&\leq \rho(c(T^{m_k}x - T^{n_k+1}x)) + \rho(\alpha_0 l(T^{n_k+1}x - T^{n_k}x)) \\
&\leq \psi(\rho(l(T^{m_k-1}x - T^{n_k}x))) + \rho(\alpha_0 l(T^{n_k+1}x - T^{n_k}x)) \\
&\leq \rho(l(T^{m_k-1}x - T^{n_k}x)) + \rho(\alpha_0 l(T^{n_k+1}x - T^{n_k}x)) \\
&\leq \epsilon + \rho(\alpha_0 l(T^{n_k+1}x - T^{n_k}x)).
\end{aligned} \tag{2.9}$$

If  $k$  tends to infinity, and by  $\Delta_2$ -condition,  $\rho(\alpha_0 l(T^{n_k+1}x - T^{n_k}x)) \rightarrow 0$  (note that  $\alpha_0 l = c(\alpha_0 - 1)$ ). Hence,  $d_k \rightarrow \epsilon$ , as  $k \rightarrow \infty$ . Now,

$$\begin{aligned}
d_k &= \rho(l(T^{m_k}x - T^{n_k}x)) \\
&\leq \rho(c(T^{m_k+1}x - T^{n_k+1}x)) + \rho(2\alpha_0 l(T^{m_k}x - T^{m_k+1}x)) + \rho(2\alpha_0 l(T^{n_k+1}x - T^{n_k}x)) \\
&\leq \psi(\rho(l(T^{m_k}x - T^{n_k}x))) + \rho(2\alpha_0 l(T^{m_k}x - T^{m_k+1}x)) + \rho(2\alpha_0 l(T^{n_k+1}x - T^{n_k}x)).
\end{aligned} \tag{2.10}$$

Thus, as  $k \rightarrow \infty$ , we obtain  $\epsilon \leq \psi(\epsilon)$ , which is a contradiction for  $\epsilon > 0$ . Therefore  $\{lT^n x\}$  is a  $\rho$ -Cauchy sequence, and by  $\Delta_2$ -condition,  $\{T^n x\}$  is a  $\rho$ -Cauchy sequence, and by the fact that  $X_\rho$  is  $\rho$ -complete, there is a  $z \in B$  such that  $\rho(T^n x - z) \rightarrow 0$  as  $n \rightarrow +\infty$ . Now, it is enough to show that  $z$  is a fixed point of  $T$ . Indeed,

$$\begin{aligned}
\rho\left(\frac{c}{2}(Tz - z)\right) &= \rho\left(\frac{c}{2}(Tz - T^{n+1}x) + \frac{c}{2}(T^{n+1}x - z)\right) \\
&\leq \rho(c(Tz - T^{n+1}x)) + \rho(c(T^{n+1}x - z)) \\
&\leq \psi(\rho(l(z - T^n x))) + \rho(c(T^{n+1}x - z)) \\
&\leq \rho(c(z - T^n x)) + \rho(c(T^{n+1}x - z)).
\end{aligned} \tag{2.11}$$

Since  $\rho(c(z - T^n x)) + \rho(c(T^{n+1}x - z)) \rightarrow 0$  as  $n \rightarrow \infty$ , then  $\rho(c/2(Tz - z)) = 0$  and  $Tz = z$ . The proof is complete.  $\square$

The following two corollaries (see [5, 14]) are immediate consequences of Theorem 2.1.

**COROLLARY 2.2.** *Let  $X_\rho$  be a  $\rho$ -complete modular space where  $\rho$  satisfies the  $\Delta_2$ -condition. Let  $B$  be a  $\rho$ -closed subset of  $X_\rho$  and let  $T : B \rightarrow B$  be a mapping such that there exist  $c, k, l \in \mathbb{R}^+$ ,  $c > l$  and  $k \in (0, 1)$ ,*

$$\rho(c(Tx - Ty)) \leq k\rho(l(x - y)), \tag{2.12}$$

for all  $x, y \in B$ . Then  $T$  has a fixed point.

## 6 Abstract and Applied Analysis

**COROLLARY 2.3.** *Let  $X_\rho$  be a  $\rho$ -complete modular space, where  $\rho$  is  $s$ -convex and satisfies the  $\Delta_2$ -condition. Also, assume that  $B \subseteq X_\rho$  is a  $\rho$ -closed subset of  $X_\rho$  and  $T : B \rightarrow B$  is a mapping such that there exist  $c, k, l \in \mathbb{R}^+$  with  $c > \max\{l, kl\}$ ,*

$$\rho(c(Tx - Ty)) \leq k^s \rho(l(x - y)), \quad (2.13)$$

for all  $x, y \in B$ . Then  $T$  has a fixed point.

*Proof.* Consider  $l_0$  to be one constant such that  $c > l_0 > \max\{l, kl\}$ . Then we have

$$\rho(c(Tx - Ty)) \leq k^s \rho(l(x - y)) = k^s \rho\left(\frac{l}{l_0}l_0(x - y)\right) \leq \left(\frac{lk}{l_0}\right)^s \rho(l_0(x - y)). \quad (2.14)$$

Thus we get

$$\rho(c(Tx - Ty)) \leq k_0 \rho(l_0(x - y)), \quad (2.15)$$

where  $c > l_0$  and  $k_0 = (lk/l_0)^s < 1$ . So by using Corollary 2.2, the proof is complete.  $\square$

### 3. A fixed point of asymptotic contraction

The concept of “asymptotic contraction” is suggested by one of the earliest versions of Banach’s principle attributed to Caccioppoli [15] and it has a long history in the nonlinear functional analysis [16]. Many mathematicians (such as Chen [17], Gerhardy [18], Jachymski and Józwik [19], Kirk [20], Suzuki [21], Xu [22], etc.) studied this concept and proved the existence of fixed points. In this section, Kirk’s fixed point theorem for asymptotic contraction is proved in modular spaces. In order to do this, we need a theorem from [14] as follows.

**THEOREM 3.1.** *Let  $X_\rho$  be a  $\rho$ -complete modular space. Let  $\{F_n\}_n$  be a decreasing sequence of nonempty  $\rho$ -closed subsets of  $X_\rho$  with  $\delta_\rho(F_n) \rightarrow 0$  as  $n \rightarrow +\infty$ . Then  $\bigcap_n F_n$  is reduced to one point.*

**Definition 3.2.** A function  $T : X_\rho \rightarrow X_\rho$  is called  $\rho$ -continuous if

$$\rho(x_n - x) \rightarrow 0, \quad \text{then } \rho(T(x_n) - T(x)) \rightarrow 0. \quad (3.1)$$

Now, we state Kirk’s fixed point theorem for asymptotic contraction in modular spaces (see [8]).

**THEOREM 3.3.** *Let  $X_\rho$  be a  $\rho$ -complete modular space. Also, assume that  $\rho$  satisfies the  $\Delta_2$ -condition and the Fatou property. Let  $f : X_\rho \rightarrow X_\rho$  be a  $\rho$ -continuous mapping and there exists a sequence  $\{\varphi_i\}_i$  of continuous functions such that  $\varphi_i : [0, +\infty) \rightarrow [0, +\infty)$  for  $i \in \mathbb{N}$  and there exists  $c > 1$  such that*

$$\rho(c(f^i(x) - f^i(y))) \leq \varphi_i(\rho(x - y)), \quad (3.2)$$

for all  $x, y \in X_\rho$ . Let  $\varphi_i \rightarrow \varphi$  uniformly on the range of  $\rho$ , where  $\varphi : [0, +\infty) \rightarrow [0, +\infty)$  and  $\varphi(r) < r$  for all  $r > 0$  and  $\varphi(0) = 0$ . If there exists an  $x \in X_\rho$  such that the sequence  $\{f^n(x)\}_{n \in \mathbb{N}}$  is  $\rho$ -bounded, then  $f$  has a unique fixed point.

*Proof.* Note that  $\{\varphi_i\}_i$  is continuous for all  $i$  and since  $\{\varphi_i\}_i$  converge uniformly to  $\varphi$ , then  $\varphi$  is continuous.

Now for each  $x, y \in X_\rho$ ,  $x \neq y$ ,

$$\limsup \rho(c(f^n(x) - f^n(y))) \leq \limsup \varphi_n(\rho(x - y)) = \varphi(\rho(x - y)) < \rho(x - y). \quad (3.3)$$

Now, we prove that  $\lim \rho(f^n(x) - f^n(y)) = 0$  for all  $x, y \in X_\rho$ . Otherwise, there exist  $x, y \in X_\rho$  and  $\varepsilon > 0$  such that

$$\limsup \rho(f^n(x) - f^n(y)) = \varepsilon. \quad (3.4)$$

Then there exists  $k$  such that

$$\varphi \rho(f^k(x) - f^k(y)) < \varepsilon. \quad (3.5)$$

Otherwise,  $\varphi \rho(f^k(x) - f^k(y)) \geq \varepsilon$  for all  $k$ . Then by taking  $\limsup$  from both sides of it, continuity of  $\varphi$ , and (3.4), we have  $\varphi(\varepsilon) \geq \varepsilon$ . This is in contradiction with  $\varphi(\varepsilon) < \varepsilon$ .

Therefore, (3.4) and (3.5) state that

$$\begin{aligned} \varepsilon &= \limsup \rho(f^n(x) - f^n(y)) \leq \limsup \rho(c(f^n(x) - f^n(y))) \\ &= \limsup \rho(c(f^n(f^k(x)) - f^n(f^k(y)))) \leq \limsup \varphi_n \rho((f^k(x) - f^k(y))) \\ &= \varphi(\rho(f^k(x) - f^k(y))) < \varepsilon. \end{aligned} \quad (3.6)$$

This is clearly a contradiction. Thus we get

$$\lim_{n \rightarrow \infty} \rho(f^n(x) - f^n(y)) = 0, \quad (3.7)$$

for all  $x, y \in X_\rho$ . Since  $\rho$  satisfies the  $\Delta_2$ -condition, then

$$\lim_{n \rightarrow \infty} \rho(c(f^n(x) - f^n(y))) = 0, \quad (3.8)$$

for all  $x, y \in X_\rho$ . This means that the sequence  $\{f^n(x)\}_n$  for all  $x \in X_\rho$  and all  $n \in \mathbb{N}$  is  $\rho$ -bounded.

Now, we assume that  $a \in X_\rho$  is arbitrary and  $a_n = f^n(a)$  for  $n \in \mathbb{N}$ , and let  $Y = \overline{\{a_n\}}^\rho$ . We can choose  $\alpha \in \mathbb{R}^+$  such that  $1/\alpha + 1/c = 1$ . Consider the sets defined by

$$F_n = \left\{ x \in Y; \rho(L(x - f^k(x))) \leq \frac{1}{n}, k = 1, \dots, n \right\}, \quad (3.9)$$

where  $L = \max\{c, 2\alpha\}$ .

The  $\rho$ -boundedness of  $\{a_n\}$  implies that  $Y$  is  $\rho$ -bounded. By using (3.8), and considering the  $\Delta_2$ -condition of  $\rho$ , we get  $F_n \neq \emptyset$  for all  $n$ , and  $F_n$  is  $\rho$ -closed, since  $f$  is continuous. Indeed, if  $\{x_m\} \subset F_n$  is a sequence such that  $x_m \rightarrow x_0$ , then

$$\rho(L(x_m - f^k(x_m))) < \frac{1}{n}, \quad (3.10)$$

## 8 Abstract and Applied Analysis

for all  $m$  and  $k = 1, 2, \dots, n$ . By the Fatou property of  $\rho$ , and (3.10), we have

$$\rho(L(x_0 - f^k(x_0))) < \liminf_{m \rightarrow \infty} \rho(L(x_m - f^k(x_m))) < \frac{1}{n}. \quad (3.11)$$

Therefore  $x_0 \in F_n$  and this means that  $F_n$  is  $\rho$ -closed.

It is clear that  $F_{n+1} \subseteq F_n$ , for all  $n$ . Now, it is enough to show that  $\delta_\rho(F_n) \rightarrow 0$ , as  $n \rightarrow \infty$ . Suppose that  $\{x_n\}$ ,  $\{y_n\}$  are two arbitrary sequences with  $x_n, y_n \in F_n$ . Consider the subsequences  $\{x_{n_j}\}$ ,  $\{y_{n_j}\}$  such that

$$\lim_{n_j \rightarrow \infty} \rho(x_{n_j} - y_{n_j}) = \limsup \rho(x_n - y_n). \quad (3.12)$$

Then

$$\begin{aligned} \rho(x_{n_j} - y_{n_j}) &= \rho\left(\frac{\alpha}{\alpha}(x_{n_j} - f^{n_j}(x_{n_j})) + \frac{c}{c}(f^{n_j}(x_{n_j}) - f^{n_j}(y_{n_j})) + \frac{\alpha}{\alpha}(f^{n_j}(y_{n_j}) - y_{n_j})\right) \\ &\leq \rho(\alpha(x_{n_j} - f^{n_j}(x_{n_j})) + \alpha(f^{n_j}(y_{n_j}) - y_{n_j})) + \rho(c(f^{n_j}(x_{n_j}) - f^{n_j}(y_{n_j}))) \\ &= \rho\left(\frac{2\alpha}{2}(x_{n_j} - f^{n_j}(x_{n_j})) + \frac{2\alpha}{2}(f^{n_j}(y_{n_j}) - y_{n_j})\right) + \rho(c(f^{n_j}(x_{n_j}) - f^{n_j}(y_{n_j}))) \\ &\leq \rho(2\alpha(x_{n_j} - f^{n_j}(x_{n_j}))) + \rho(2\alpha(f^{n_j}(y_{n_j}) - y_{n_j})) + \varphi_{n_j}(\rho(x_{n_j} - y_{n_j})) \\ &\leq \rho(L(x_{n_j} - f^{n_j}(x_{n_j}))) + \rho(L(f^{n_j}(y_{n_j}) - y_{n_j})) + \varphi_{n_j}(\rho(x_{n_j} - y_{n_j})) \\ &\leq \frac{2}{n_j} + \varphi_{n_j}(\rho(x_{n_j} - y_{n_j})). \end{aligned} \quad (3.13)$$

Taking limit from both sides,

$$\lim_{n_j \rightarrow +\infty} \rho(x_{n_j} - y_{n_j}) \leq \lim_{n_j \rightarrow +\infty} \frac{2}{n_j} + \lim_{n_j \rightarrow +\infty} \varphi_{n_j}(\rho(x_{n_j} - y_{n_j})) = \varphi\left(\lim_{n_j \rightarrow +\infty} (\rho(x_{n_j} - y_{n_j}))\right). \quad (3.14)$$

Thus, we have

$$\limsup \rho(x_n - y_n) \leq \varphi(\limsup \rho(x_n - y_n)). \quad (3.15)$$

On the other hand, we have  $\varphi(\limsup \rho(x_n - y_n)) < \limsup \rho(x_n - y_n)$ . So, we get

$$\limsup \rho(x_n - y_n) = 0. \quad (3.16)$$

Therefore

$$\delta_\rho(F_n) = 0 \quad \text{as } n \rightarrow \infty. \quad (3.17)$$

Consequently,  $\{F_n\}$  satisfies all conditions of Theorem 3.1, and then  $\bigcap_n F_n = \{z\}$ . Since  $z \in F_n$  for all  $n$ , then  $\rho(L(z - f(z))) < 1/n$ , for all  $n$ . Then letting  $n \rightarrow \infty$ , we have  $\rho(L(z - f(z))) = 0$ . Thus  $L(z - f(z)) = 0$ . this means that  $f(z) = z$ , and the proof is complete.  $\square$

## Acknowledgments

The authors would like to thank the anonymous referees whose helpful comments led to many improvements in this paper. Moreover, the first author would like to thank the Institute for Studies in Theoretical Physics and Mathematics (IPM), Tehran, Iran, for supporting this research (Grant no. 85340045).

## References

- [1] H. Nakano, *Modular Semi-Ordered Spaces*, Tokyo, Japan, 1959.
- [2] J. Musielak and W. Orlicz, “On modular spaces,” *Studia Mathematica*, vol. 18, pp. 49–65, 1959.
- [3] T. Dominguez Benavides, M. A. Khamsi, and S. Samadi, “Uniformly Lipschitzian mappings in modular function spaces,” *Nonlinear Analysis*, vol. 46, no. 2, Ser. A: Theory Methods, pp. 267–278, 2001.
- [4] A. Hajji and E. Hanebaly, “Fixed point theorem and its application to perturbed integral equations in modular function spaces,” *Electronic Journal of Differential Equations*, vol. 2005, no. 105, pp. 1–11, 2005.
- [5] E. Hanebaly, “Fixed point theorems in modular space,” November 2005, <http://arxiv.org/abs/math.FA/0511319v1>.
- [6] M. A. Khamsi, “Nonlinear semigroups in modular function spaces,” *Mathematica Japonica*, vol. 37, no. 2, pp. 291–299, 1992.
- [7] D. W. Boyd and J. S. W. Wong, “On nonlinear contractions,” *Proceedings of the American Mathematical Society*, vol. 20, no. 2, pp. 458–464, 1969.
- [8] I. D. Arandelović, “On a fixed point theorem of Kirk,” *Journal of Mathematical Analysis and Applications*, vol. 301, no. 2, pp. 384–385, 2005.
- [9] M. Edelstein, “On fixed and periodic points under contractive mappings,” *Journal of the London Mathematical Society*, vol. 37, no. 1, pp. 74–79, 1962.
- [10] L. B. Ćirić, “A generalization of Banach’s contraction principle,” *Proceedings of the American Mathematical Society*, vol. 45, no. 2, pp. 267–273, 1974.
- [11] E. Rakotch, “A note on contractive mappings,” *Proceedings of the American Mathematical Society*, vol. 13, no. 3, pp. 459–465, 1962.
- [12] S. Reich, “Fixed points of contractive functions,” *Bollettino dell’Unione Matematica Italiana* (4), vol. 5, pp. 26–42, 1972.
- [13] W. A. Kirk, “Contraction mappings and extensions,” in *Handbook of Metric Fixed Point Theory*, W. A. Kirk and B. Sims, Eds., pp. 1–34, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2001.
- [14] A. Ait Taleb and E. Hanebaly, “A fixed point theorem and its application to integral equations in modular function spaces,” *Proceedings of the American Mathematical Society*, vol. 128, no. 2, pp. 419–426, 2000.
- [15] R. Caccioppoli, “Una teorem general sull’esistenza di elementi uniti in una transformazione funzionale,” *Rendiconti dell’Accademia Nazionale dei Lincei*, vol. 11, pp. 794–799, 1930.
- [16] F. E. Browder, “Nonlinear operators and nonlinear equations of evolution in Banach spaces,” in *Nonlinear Functional Analysis (Proc. Sympos. Pure Math., Vol. XVIII, Part 2, Chicago, Ill., 1968)*, pp. 1–308, American Mathematical Society, Providence, RI, USA, 1976.
- [17] Y.-Z. Chen, “Asymptotic fixed points for nonlinear contractions,” *Fixed Point Theory and Applications*, vol. 2005, no. 2, pp. 213–217, 2005.
- [18] P. Gerhardy, “A quantitative version of Kirk’s fixed point theorem for asymptotic contractions,” *Journal of Mathematical Analysis and Applications*, vol. 316, no. 1, pp. 339–345, 2006.
- [19] J. Jachymski and I. Józwik, “On Kirk’s asymptotic contractions,” *Journal of Mathematical Analysis and Applications*, vol. 300, no. 1, pp. 147–159, 2004.

- [20] W. A. Kirk, “Fixed points of asymptotic contractions,” *Journal of Mathematical Analysis and Applications*, vol. 277, no. 2, pp. 645–650, 2003.
- [21] T. Suzuki, “Fixed-point theorem for asymptotic contractions of Meir-Keeler type in complete metric spaces,” *Nonlinear Analysis*, vol. 64, no. 5, pp. 971–978, 2006.
- [22] H.-K. Xu, “Asymptotic and weakly asymptotic contractions,” *Indian Journal of Pure and Applied Mathematics*, vol. 36, no. 3, pp. 145–150, 2005.

A. Razani: Department of Mathematics, Faculty of Science, Imam Khomeini International University, P.O. Box 34194-288, Qazvin, Iran; Institute for Studies in Theoretical Physics and Mathematics (IPM), P.O. Box 19395-5531, Tehran, Iran

*Email addresses:* razani@ipm.ir; razani@ikiu.ac.ir

E. Nabizadeh: Department of Mathematics, Iran University of Science and Technology, Tehran 16844, Iran

*Email address:* elmira\_nabizadeh@yahoo.com

M. Beyg Mohamadi: Department of Mathematics, Iran University of Science and Technology, Tehran 16844, Iran

*Email address:* maryambmohamadi@mathdep.iust.ac.ir

S. Homaei Pour: Department of Mathematics, Iran University of Science and Technology, Tehran 16844, Iran

*Email address:* smn\_homaei@yahoo.com

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

### Guest Editors

**Edson Denis Leonel**, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)