

ON THE EXISTENCE OF A POSITIVE SOLUTION FOR A SEMILINEAR ELLIPTIC EQUATION IN THE UPPER HALF SPACE

CHEN SHAOWEI AND LI YONGQING

Received 14 September 2001

We show the existence of a nontrivial solution to the semilinear elliptic equation $-\Delta u + u = b(x)|u|^{p-2}u$, $u > 0$, $u \in H_0^1(\mathbb{R}_+^N)$ under some suitable conditions.

1. Introduction

In this paper, we study the following problem:

$$-\Delta u + u = b(x)|u|^{p-2}u, \quad u > 0, \quad u \in H_0^1(\mathbb{R}_+^N), \quad (1.1)$$

where $2 < p < 2^*$, $2^* = 2N/(N-2)$ when $N > 2$ and $2^* = +\infty$ when $N = 2$, $\mathbb{R}_+^N = \{(x_1, x_2, \dots, x_N) \mid x_N > 0\}$ is the upper half space of \mathbb{R}^N . We put the following conditions on b :

(B₁) $b \in C(\mathbb{R}_+^N)$, and

$$1 = \inf_{x \in \mathbb{R}_+^N} b(x) = \lim_{|x| \rightarrow +\infty, x \in \mathbb{R}_+^N} b(x); \quad (1.2)$$

(B₂) there exists $\delta \in (0, 1)$, such that

$$\lim_{|x| \rightarrow \infty} (b(x) - 1)e^{(1-\delta)|x|} = +\infty. \quad (1.3)$$

Our main result is the following result.

THEOREM 1.1. *If b satisfies (B₁) and (B₂), then (1.1) has a nontrivial solution.*

Similar problems have been studied extensively. In [3], Esteban and Lions showed that $-\Delta u + u = |u|^{p-2}u$, $u \in H_0^1(\mathbb{R}_+^N)$ does not have nontrivial solution.

However, with the help of the potential $b(x)$ and the comparison technique derived in [2], we proved the existence of a nontrivial solution to problem (1.1).

2. Proof of Theorem 1.1

Let

$$\varphi(u) = \int_{\mathbb{R}_+^N} \left(\frac{1}{2} |\nabla u|^2 + \frac{1}{2} |u|^2 \right) dx - \frac{1}{p} \int_{\mathbb{R}_+^N} b(x) |u|^p dx. \quad (2.1)$$

LEMMA 2.1. Denote $S_p = \inf_{u \in H_0^1(\mathbb{R}_+^N), |u|_p=1} \|u\|^2$. Then

$$\inf_{u \in H_0^1(\mathbb{R}_+^N), |u|_p=1} \|u\|^2 = S_p. \quad (2.2)$$

LEMMA 2.2. If $0 < c < c^* := (1/2 - 1/p)S_p^{p/(p-2)}$, then φ satisfies $(PS)_c$ condition.

One can see [2] or [5] for the proof of Lemmas 2.1 and 2.2.

LEMMA 2.3. The equation $-\Delta u + u = |u|^{p-2}u$ has a unique positive solution w in \mathbb{R}^N and satisfies the following conditions:

- (i) $w > 0$ in \mathbb{R}^N , $w \in C^\infty(\mathbb{R}^N)$.
- (ii) $w(x) = w(|x|)$.
- (iii) $w(r)r^{(N-1)/2} \exp(r) \rightarrow C > 0$ and $w'(r)r^{(N-1)/2} \exp(r) \rightarrow -C$, where $r = |x| \rightarrow +\infty$.

Proof. See [1, 4]. □

Proof of Theorem 1.1. By Lemma 2.2, we need only to prove that the Mountain Pass value of φ ,

$$c = \inf \left\{ \left(\frac{1}{2} - \frac{1}{p} \right) \left(\frac{\|v\|^2}{\left(\int_{\mathbb{R}_+^N} b(x) |v|^p dx \right)^{2/p}} \right)^{p/(p-2)} ; v \in H_0^1(\mathbb{R}_+^N), v \neq \theta \right\}, \quad (2.3)$$

satisfies that $0 < c < c^*$.

Set

$$\psi_n(x) = \begin{cases} 1, & |x| \leq n-1, \\ n-|x|, & n-1 < |x| \leq n, \\ 0, & |x| > n. \end{cases} \quad (2.4)$$

Let $x_n = (0, 0, \dots, 0, n)$, $u_n = \psi_n(\cdot - x_n)w(\cdot - x_n)$. Then $u_n \in H_0^1(\mathbb{R}_+^N)$, $n = 1, 2, \dots$.

Notice that

$$\begin{aligned}
\|u_n\|^2 &= \int_{\mathbb{R}^N} |\nabla u_n|^2 dx + \int_{\mathbb{R}^N} u_n^2 dx \\
&= \int_{\mathbb{R}^N} |\nabla(\psi_n(\cdot-x_n)w(\cdot-x_n))|^2 dx + \int_{\mathbb{R}^N} |\psi_n(\cdot-x_n)w(\cdot-x_n)|^2 dx \\
&= \int_{B(\theta,n)} |\nabla(\psi_n \cdot w)|^2 dx + \int_{B(\theta,n)} |\psi_n \cdot w|^2 dx \\
&= \int_{B(\theta,n-1)} |\nabla \omega|^2 dx + \int_{B(\theta,n-1)} \omega^2 dx \\
&\quad + \int_{B_{n-1,n}} \psi_n^2 \cdot |\nabla \omega|^2 dx + \int_{B_{n-1,n}} |\nabla \psi_n|^2 \cdot \omega^2 dx \\
&\quad + 2 \int_{B_{n-1,n}} \omega \cdot \psi_n \cdot \nabla \omega \cdot \nabla \psi_n dx + \int_{B_{n-1,n}} \psi_n^2 \cdot \omega^2 dx \\
&= \int_{\mathbb{R}^N} |\nabla \omega|^2 dx + \int_{\mathbb{R}^N} \omega^2 dx \\
&\quad + \int_{\mathbb{R}^N \setminus B(\theta,n-1)} |\nabla \omega|^2 dx + \int_{\mathbb{R}^N \setminus B(\theta,n-1)} |\omega|^2 dx \\
&\quad + \int_{B_{n-1,n}} \psi_n^2 \cdot |\nabla \omega|^2 dx + 2 \int_{B_{n-1,n}} \omega \cdot \psi_n \cdot \nabla \omega \cdot \nabla \psi_n dx \\
&\quad + \int_{B_{n-1,n}} |\nabla \psi_n|^2 \cdot \omega^2 dx + \int_{B_{n-1,n}} \psi_n^2 \cdot \omega^2 dx,
\end{aligned} \tag{2.5}$$

where $B_{n-1,n} = B(\theta, n) \setminus B(\theta, n-1)$.

From [Lemma 2.3](#) we know that, when n is big enough,

$$\begin{aligned}
\int_{\mathbb{R}^N \setminus B(\theta,n-1)} |\nabla \omega|^2 dx &= C_1 \int_{n-1}^{+\infty} |\omega'(r)|^2 \cdot r^{N-1} dr \\
&\leq C'_1 \int_{n-1}^{+\infty} e^{-2r} dr = \frac{C'_1}{e^{2(n-1)}}.
\end{aligned} \tag{2.6}$$

Similarly, when n is big enough, we get

$$\int_{\mathbb{R}^N \setminus B(\theta,n-1)} \omega^2 dx \leq \frac{C'_2}{e^{2(n-1)}}. \tag{2.7}$$

So

$$\|u_n\|^2 = \|\omega\|^2 + O\left(\frac{1}{e^{2(n-1)}}\right). \tag{2.8}$$

32 Positive solutions in upper half space

Let $a(x) = b(x) - 1$. Then

$$\begin{aligned}
\int_{\mathbb{R}_+^N} b(x)|u_n|^p dx &= \int_{\mathbb{R}_+^N} |u_n|^p dx + \int_{\mathbb{R}_+^N} a(x)|u_n|^p dx \\
&= \int_{B(\theta, n-1)} |\omega|^p dx + \int_{B(\theta, n-1)} a(x+x_n)|\omega|^p dx \\
&\quad + \int_{B_{n-1, n}} a(x+x_n)|\psi_n(x) \cdot \omega(x)|^p dx \\
&= \int_{\mathbb{R}^N} |\omega|^p dx + \int_{B(\theta, n-1)} a(x+x_n)|\omega|^p dx \\
&\quad + \int_{B_{n-1, n}} a(x+x_n)|\psi_n(x) \cdot \omega(x)|^p dx + \int_{\mathbb{R}^N \setminus B(\theta, n-1)} |\omega|^p dx.
\end{aligned} \tag{2.9}$$

Thus

$$\begin{aligned}
\int_{\mathbb{R}^N \setminus B(\theta, n-1)} |\omega|^p dx &= o\left(\frac{1}{e^{2(n-1)}}\right), \\
\int_{B_{n-1, n}} a(x+x_n)|\psi_n(x) \cdot \omega(x)|^p dx &= o\left(\frac{1}{e^{2(n-1)}}\right).
\end{aligned} \tag{2.10}$$

From (B₂), we see that there exists a positive constant λ , such that

$$a(x) \geq \frac{\lambda}{e^{(1-\delta)|x|}}. \tag{2.11}$$

So there exists a positive constant C_3 , such that

$$\int_{\mathbb{R}_+^N} b(x)|u_n|^p dx \geq \int_{\mathbb{R}^N} |\omega|^p dx + \frac{C_3}{e^{2(1-\delta)(n-1)}}. \tag{2.12}$$

Then

$$\begin{aligned}
&\left(\int_{\mathbb{R}_+^N} b(x)|u_n|^p dx\right)^{2/p} \\
&\geq \left(\int_{\mathbb{R}^N} |\omega|^p dx + \frac{C_3}{e^{2(1-\delta)(n-1)}}\right)^{2/p} \\
&= \left(\int_{\mathbb{R}^N} |\omega|^p dx\right)^{2/p} \left(1 + \frac{C_4}{e^{2(1-\delta)(n-1)}}\right)^{2/p} \\
&= \left(\int_{\mathbb{R}^N} |\omega|^p dx\right)^{2/p} \left(1 + \frac{C_5}{e^{2(1-\delta)(n-1)}} + o\left(\frac{1}{e^{2(1-\delta)(n-1)}}\right)\right).
\end{aligned} \tag{2.13}$$

Thus

$$\frac{\|u_n\|^2}{\left(\int_{\mathbb{R}_+^N} b(x)|u_n|^p dx\right)^{2/p}} = \frac{\|\omega\|^2}{\left(\int_{\mathbb{R}^N} \omega^p dx\right)^{2/p}} - \frac{C_6}{e^{2(1-\delta)(n-1)}} + o\left(\frac{1}{e^{2(1-\delta)(n-1)}}\right). \quad (2.14)$$

So

$$\frac{\|u_n\|^2}{\left(\int_{\mathbb{R}_+^N} b(x)|u_n|^p dx\right)^{2/p}} < \frac{\|\omega\|^2}{\left(\int_{\mathbb{R}^N} \omega^p dx\right)^{2/p}}, \quad (2.15)$$

when n is big enough.

Thus, we come to the conclusion that

$$c < \left(\frac{1}{2} - \frac{1}{p}\right) S_p^{p/(p-2)}. \quad (2.16)$$

□

Acknowledgement

This work was supported in part by the National Natural Science Foundation and the Fujian Province Natural Science Foundation of China.

References

- [1] H. Berestycki and P.-L. Lions, *Nonlinear scalar field equations. I. Existence of a ground state*, Arch. Rational Mech. Anal. **82** (1983), no. 4, 313–345.
- [2] W. Y. Ding and W. M. Ni, *On the existence of positive entire solutions of a semilinear elliptic equation*, Arch. Rational Mech. Anal. **91** (1986), no. 4, 283–308.
- [3] M. J. Esteban and P.-L. Lions, *Existence and nonexistence results for semilinear elliptic problems in unbounded domains*, Proc. Roy. Soc. Edinburgh Sect. A **93** (1982/83), no. 1-2, 1–14.
- [4] B. Gidas, W. M. Ni, and L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), no. 3, 209–243.
- [5] M. Willem, *Minimax Theorems*, Progress in Nonlinear Differential Equations and Their Applications, vol. 24, Birkhäuser, Massachusetts, 1996.

CHEN SHAOWEI: DEPARTMENT OF MATHEMATICS, FUJIAN NORMAL UNIVERSITY, FUZHOU, FUJIAN 35007, CHINA

LI YONGQING: DEPARTMENT OF MATHEMATICS, FUJIAN NORMAL UNIVERSITY, FUZHOU, FUJIAN 35007, CHINA

E-mail address: yqli@pub1.fz.fj.cn

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru