

ON THE DIFFERENCE EQUATION $x_{n+1} = ax_n - bx_n/(cx_n - dx_{n-1})$

E. M. ELABBASY, H. EL-METWALLY, AND E. M. ELSAYED

Received 14 June 2006; Revised 3 September 2006; Accepted 26 September 2006

We investigate some qualitative behavior of the solutions of the difference equation $x_{n+1} = ax_n - bx_n/(cx_n - dx_{n-1})$, $n = 0, 1, \dots$, where the initial conditions x_{-1}, x_0 are arbitrary real numbers and a, b, c, d are positive constants.

Copyright © 2006 E. M. Elabbasy et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In this paper we deal with some properties of the solutions of the difference equation

$$x_{n+1} = ax_n - \frac{bx_n}{cx_n - dx_{n-1}}, \quad n = 0, 1, \dots, \quad (1.1)$$

where the initial conditions x_{-1}, x_0 are arbitrary real numbers and a, b, c, d are positive constants.

Recently, there has been a lot of interest in studying the global attractivity, boundedness character, and the periodic nature of nonlinear difference equations. For some results in this area, see, for example, [1–13], we recall some notations and results which will be useful in our investigation.

Let I be some interval of real numbers and the function f has continuous partial derivatives on I^{k+1} , where $I^{k+1} = I \times I \times \dots \times I$ ($k+1$ – times). Then, for initial conditions $x_{-k}, x_{-k+1}, \dots, x_0 \in I$, it is easy to see that the difference equation

$$x_{n+1} = f(x_n, x_{n-1}, \dots, x_{n-k}), \quad n = 0, 1, \dots, \quad (1.2)$$

has a unique solution $\{x_n\}_{n=-k}^{\infty}$.

2 On the difference equation $x_{n+1} = ax_n - bx_n/(cx_n - dx_{n-1})$

A point $\bar{x} \in I$ is called an equilibrium point of (1.2) if

$$\bar{x} = f(\bar{x}, \bar{x}, \dots, \bar{x}). \quad (1.3)$$

That is, $x_n = \bar{x}$ for $n \geq 0$ is a solution of (1.2), or equivalently, \bar{x} is a fixed point of f .

Definition 1.1 (stability). (i) The equilibrium point \bar{x} of (1.2) is locally stable if for every $\epsilon > 0$, there exists $\delta > 0$ such that for all $x_{-k}, x_{-k+1}, \dots, x_{-1}, x_0 \in I$, with

$$\begin{aligned} |x_{-k} - \bar{x}| + |x_{-k+1} - \bar{x}| + \dots + |x_0 - \bar{x}| &< \delta, \\ |x_n - \bar{x}| &< \epsilon \quad \forall n \geq -k. \end{aligned} \quad (1.4)$$

(ii) The equilibrium point \bar{x} of (1.2) is locally asymptotically stable if \bar{x} is locally stable solution of (1.2) and there exists $\gamma > 0$ such that for all $x_{-k}, x_{-k+1}, \dots, x_{-1}, x_0 \in I$, with

$$\begin{aligned} |x_{-k} - \bar{x}| + |x_{-k+1} - \bar{x}| + \dots + |x_0 - \bar{x}| &< \gamma, \\ \lim_{n \rightarrow \infty} x_n &= \bar{x}. \end{aligned} \quad (1.5)$$

(iii) The equilibrium point \bar{x} of (1.2) is global attractor if for all $x_{-k}, x_{-k+1}, \dots, x_{-1}, x_0 \in I$,

$$\lim_{n \rightarrow \infty} x_n = \bar{x}. \quad (1.6)$$

(iv) The equilibrium point \bar{x} of (1.2) is globally asymptotically stable if \bar{x} is locally stable, and \bar{x} is also a global attractor of (1.2).

(v) The equilibrium point \bar{x} of (1.2) is unstable if \bar{x} is not locally stable.

The linearized equation of (1.2) about the equilibrium \bar{x} is the linear difference equation

$$y_{n+1} = \sum_{i=0}^k \frac{\partial f(\bar{x}, \bar{x}, \dots, \bar{x})}{\partial x_{n-i}} y_{n-i}. \quad (1.7)$$

Now assume that the characteristic equation associated with (1.7) is

$$p(\lambda) = p_0 \lambda^k + p_1 \lambda^{k-1} + \dots + p_{k-1} \lambda + p_k = 0, \quad (1.8)$$

where $p_i = \partial f(\bar{x}, \bar{x}, \dots, \bar{x}) / \partial x_{n-i}$.

THEOREM 1.2 [9]. *Assume that $p_i \in \mathbb{R}$, $i = 1, 2, \dots$, and $k \in \{0, 1, 2, \dots\}$. Then*

$$\sum_{i=1}^k |p_i| < 1 \quad (1.9)$$

is a sufficient condition for the asymptotic stability of the difference equation

$$y_{n+k} + p_1 y_{n+k-1} + \dots + p_k y_n = 0, \quad n = 0, 1, \dots \quad (1.10)$$

COROLLARY 1.3 [9]. *Assume that f is a C^1 function and let \bar{x} be an equilibrium of (1.2). Then the following statements are true.*

- (a) *If all roots of the polynomial equation (1.8) lie in the open unite disk $|\lambda| < 1$, then the equilibrium \bar{x} of (1.2) is asymptotically stable.*
- (b) *If at least one root of (1.8) has absolute value greater than one, then the equilibrium \bar{x} of (1.2) is unstable.*

Remark 1.4. The condition (1.9) implies that all the roots of the polynomial equation (1.8) lie in the open unite disk $|\lambda| < 1$.

Consider the following equation:

$$x_{n+1} = f(x_n, x_{n-1}). \quad (1.11)$$

The following theorem will be useful for the proof of our main results in this paper.

THEOREM 1.5 [10]. *Let $[a, b]$ be an interval of real numbers and assume that*

$$f : [a, b]^2 \longrightarrow [a, b] \quad (1.12)$$

is a continuous function satisfying the following properties.

- (a) *$f(x, y)$ is nondecreasing in $x \in [a, b]$ for each $y \in [a, b]$, and is nonincreasing in $y \in [a, b]$ for each $x \in [a, b]$.*
- (b) *If $(m, M) \in [a, b] \times [a, b]$ is a solution of the system*

$$m = f(m, M), \quad M = f(M, m), \quad (1.13)$$

then

$$m = M. \quad (1.14)$$

Then (1.11) has a unique equilibrium $\bar{x} \in [a, b]$ and every solution of (1.11) converges to \bar{x} .

2. Periodic solutions

In this section we study the existence of periodic solutions of (1.1). The following theorem states the necessary and sufficient conditions that this equation has periodic solutions.

THEOREM 2.1. *Equation (1.1) has positive prime period-two solutions if and only if*

$$(c + d)(a + 1) > 4d, \quad ac \neq d, c > d. \quad (2.1)$$

Proof. First suppose that there exists a prime period-two solution

$$\dots, p, q, p, q, \dots \quad (2.2)$$

of (1.1). We will prove that condition (2.1) holds.

4 On the difference equation $x_{n+1} = ax_n - bx_n/(cx_n - dx_{n-1})$

We see from (1.1) that

$$\begin{aligned} p &= aq - \frac{bq}{cq - dp}, \\ q &= ap - \frac{bp}{cp - dq}. \end{aligned} \tag{2.3}$$

Then

$$cpq - dp^2 = acq^2 - adpq - bq, \tag{2.4}$$

$$cpq - dq^2 = acp^2 - adpq - bp. \tag{2.5}$$

Subtracting (2.5) from (2.4) gives

$$d(q^2 - p^2) = ac(q^2 - p^2) - b(q - p). \tag{2.6}$$

Since $p \neq q$, it follows that

$$p + q = \frac{b}{ac - d}. \tag{2.7}$$

Again, adding (2.4) and (2.5) yields

$$2cpq - d(p^2 + q^2) = ac(p^2 + q^2) - 2adpq - b(p + q). \tag{2.8}$$

It follows by (2.7), (2.8), and the relation

$$p^2 + q^2 = (p + q)^2 - 2pq \quad \forall p, q \in \mathbb{R}, \tag{2.9}$$

that

$$pq = \frac{b^2 d}{(ac - d)^2 (c + d)(a + 1)}. \tag{2.10}$$

Now it is clear from (2.7) and (2.10) that p and q are the two positive distinct roots of the quadratic equation

$$(ac - d)t^2 - bt + \frac{b^2 d}{(ac - d)(c + d)(a + 1)} = 0 \tag{2.11}$$

and so

$$b^2 > \frac{4b^2 d}{(c + d)(a + 1)}. \tag{2.12}$$

Therefore, inequality (2.1) holds.

Second, suppose that inequality (2.1) is true. We will show that (1.1) has a prime period-two solution.

Assume that

$$\begin{aligned} p &= \frac{b+\alpha}{2(ac-d)}, \\ q &= \frac{b-\alpha}{2(ac-d)}, \end{aligned} \quad (2.13)$$

where $\alpha = \sqrt{b^2 - 4b^2d/((c+d)(a+1))}$.

From inequality (2.1) it follows that α is a real positive number, therefore, p and q are distinct positive real numbers.

Set

$$x_{-1} = p, \quad x_0 = q. \quad (2.14)$$

We show that $x_1 = x_{-1} = p$ and $x_2 = x_0 = q$.

It follows from (1.1) that

$$\begin{aligned} x_1 &= aq - \frac{bq}{cq - dp} = \frac{acq^2 - adpq - bq}{cq - dp} \\ &= \frac{ac[(b-\alpha)/(2(ac-d))]^2 - ad[b^2d/((ac-d)^2(c+d)(a+1))] - b[(b-\alpha)/(2(ac-d))]}{c[(b-\alpha)/(2(ac-d))] - d[(b+\alpha)/(2(ac-d))]} \end{aligned} \quad (2.15)$$

Multiplying the denominator and numerator by $4(ac-d)^2$ gives

$$x_1 = \frac{2b^2d - (4ab^2cd + 4ab^2d^2)/((c+d)(a+1)) - 2bd\alpha}{2(ac-d)\{cb - bd - (c+d)\alpha\}\{(c+d)(a+1)\}}. \quad (2.16)$$

Multiplying the denominator and numerator by $\{cb - bd + (c+d)\alpha\}\{(c+d)(a+1)\}$ we get

$$x_1 = \frac{[(4b^3d^3 + 4b^3cd^2 - 4ab^3c^2d - 4ab^3cd^2) + (4b^2cd^2 + 4b^2d^3 - 4ab^2c^2d - 4ab^2cd^2)\alpha]}{2(ac-d)\{4b^2cd^2 + 4b^2d^3 - 4ab^2c^2d - 4ab^2cd^2\}}. \quad (2.17)$$

Dividing the denominator and numerator by $\{4b^2cd^2 + 4b^2d^3 - 4ab^2c^2d - 4ab^2cd^2\}$ gives

$$x_1 = \frac{b+\alpha}{2(ac-d)} = p. \quad (2.18)$$

Similarly as before one can easily show that

$$x_2 = q. \quad (2.19)$$

Then it follows by induction that

$$x_{2n} = q, \quad x_{2n+1} = p \quad \forall n \geq -1. \quad (2.20)$$

6 On the difference equation $x_{n+1} = ax_n - bx_n/(cx_n - dx_{n-1})$

Thus (1.1) has the positive prime period two solution

$$\dots, p, q, p, q, \dots, \quad (2.21)$$

where p and q are the distinct roots of the quadratic equation (2.11) and the proof is complete. \square

3. Local stability of the equilibrium point

In this section we study the local stability character of the solutions of (1.1).

The equilibrium points of (1.1) are given by the relation

$$\bar{x} = a\bar{x} - \frac{b\bar{x}}{c\bar{x} - d\bar{x}}. \quad (3.1)$$

If $(c - d)(a - 1) > 0$, then the only positive equilibrium point of (1.1) is given by

$$\bar{x} = \frac{b}{(c - d)(a - 1)}. \quad (3.2)$$

Let $f : (0, \infty)^2 \rightarrow (0, \infty)$ be a function defined by

$$f(u, v) = au - \frac{bu}{cu - dv}. \quad (3.3)$$

Therefore,

$$\begin{aligned} \frac{\partial f(u, v)}{\partial u} &= a + \frac{bdv}{(cu - dv)^2}, \\ \frac{\partial f(u, v)}{\partial v} &= -\frac{bdu}{(cu - dv)^2}. \end{aligned} \quad (3.4)$$

Then we see that

$$\begin{aligned} \frac{\partial f(\bar{x}, \bar{x})}{\partial u} &= a + \frac{d(a - 1)}{(c - d)} = p_0, \\ \frac{\partial f(\bar{x}, \bar{x})}{\partial v} &= -\frac{d(a - 1)}{(c - d)} = p_1. \end{aligned} \quad (3.5)$$

Then the linearized equation of (1.1) about \bar{x} is

$$y_{n+1} - p_0 y_{n-1} - p_1 y_n = 0. \quad (3.6)$$

THEOREM 3.1. *Assume that*

$$|ac - d| + |ad - d| < |c - d|. \quad (3.7)$$

Then the equilibrium point of (1.1) is locally asymptotically stable.

Proof. Suppose that

$$|ac - d| + |ad - d| < |c - d|, \quad (3.8)$$

then

$$\left| a + \frac{d(a-1)}{(c-d)} \right| + \left| - \frac{d(a-1)}{(c-d)} \right| < 1. \quad (3.9)$$

Thus

$$|p_1| + |p_0| < 1. \quad (3.10)$$

It is followed by Theorem 1.2 that (3.6) is asymptotically stable. The proof is complete. \square

4. Global attractor of the equilibrium point of (1.1)

In this section we investigate the global attractivity character of solutions of (1.1).

THEOREM 4.1. *The equilibrium point \bar{x} of (1.1) is a global attractor if $c \neq d$.*

Proof. We can easily see that the function $f(u, v)$ which is defined by (3.3) is increasing in u and decreasing in v .

Suppose that (m, M) is a solution of the system

$$m = f(m, M), \quad M = f(M, m). \quad (4.1)$$

Then it results

$$\frac{1}{cm - dM} = \frac{1}{cM - dm}, \quad (4.2)$$

that is, $M = m$. It follows by Theorem 1.5 that \bar{x} is a global attractor of (1.1) and then the proof is complete. \square

5. Special case of (1.1)

In this section we study the following special case of (1.1):

$$x_{n+1} = x_n - \frac{x_n}{x_n - x_{n-1}}, \quad (5.1)$$

where the initial conditions x_{-1}, x_0 are arbitrary real numbers with $x_{-1}, x_0 \in \mathbb{R}/\{0\}$, and $x_{-1} \neq x_0$.

8 On the difference equation $x_{n+1} = ax_n - bx_n/(cx_n - dx_{n-1})$

5.1. The solution form of (5.1). In this section we give a specific form of the solutions of (5.1).

THEOREM 5.1. Let $\{x_n\}_{n=-1}^{\infty}$ be the solution of (5.1) satisfying $x_{-1} = k$, $x_0 = h$ with $k \neq h$, $k, h \in \mathbb{R}/\{0\}$. Then for $n = 0, 1, \dots$,

$$\begin{aligned} x_{2n-1} &= k + n \left(h - k - (n-1) - \frac{h}{h-k} \right), \\ x_{2n} &= h + n \left(h - k - n - \frac{h}{h-k} \right). \end{aligned} \quad (5.2)$$

Proof. For $n = 0$ the result holds. Now suppose that $n > 0$ and that our assumption holds for $n - 1$. That is,

$$\begin{aligned} x_{2n-3} &= k + (n-1) \left(h - k - (n-2) - \frac{h}{h-k} \right), \\ x_{2n-2} &= h + (n-1) \left(h - k - (n-1) - \frac{h}{h-k} \right). \end{aligned} \quad (5.3)$$

Now, it follows from (5.1) that

$$\begin{aligned} x_{2n-1} &= x_{2n-2} - \frac{x_{2n-2}}{x_{2n-2} - x_{2n-3}} \\ &= h + (n-1) \left(h - k - (n-1) - \frac{h}{h-k} \right) \\ &\quad - \frac{h + (n-1)(h - k - (n-1) - h/(h-k))}{(h + (n-1)(h - k - (n-1) - h/(h-k))) - (k + (n-1)(h - k - (n-2) - h/(h-k)))} \\ &= h + (n-1) \left(h - k - (n-1) - \frac{h}{h-k} \right) - \frac{h + (n-1)(h - k - (n-1) - h/(h-k))}{h - k - (n-1)}. \end{aligned} \quad (5.4)$$

Multiplying the denominator and numerator by $(h - k)$ we get

$$\begin{aligned} x_{2n-1} &= k + (n-1) \left(h - k - (n-1) - \frac{h}{h-k} \right) - \frac{(h + (n-1)(h - k))}{(h - k)} + (h - k) \\ &= k + (n-1) \left(h - k - (n-1) - \frac{h}{h-k} \right) + (h - k) - (n-1) - \frac{h}{(h-k)}, \end{aligned} \quad (5.5)$$

then we have

$$x_{2n-1} = k + n \left(h - k - (n-1) - \frac{h}{h-k} \right). \quad (5.6)$$

Also, we get from (5.1)

$$\begin{aligned} x_{2n} &= x_{2n-1} - \frac{x_{2n-1}}{x_{2n-1} - x_{2n-2}} \\ &= k + n \left(h - k - (n-1) - \frac{h}{h-k} \right) + \frac{k+n(h-k-(n-1)-h/(h-k))}{(n-1)+h/(h-k)}. \end{aligned} \quad (5.7)$$

Multiplying the denominator and numerator by $(h-k)$ we get

$$\begin{aligned} x_{2n} &= k + n \left(h - k - (n-1) - \frac{h}{h-k} \right) \\ &\quad + \frac{(k(h-k)+n(h-k)^2)-(n(n-1)(h-k)+nh)}{(n-1)(h-k)+h}. \end{aligned} \quad (5.8)$$

Thus we obtain

$$x_{2n} = h + n \left(h - k - n - \frac{h}{h-k} \right). \quad (5.9)$$

Hence, the proof is complete. \square

Remark 5.2. It is easy to see that every solution of (5.1) is unbounded.

Acknowledgment

The authors would like to thank the referees for their valuable comments.

References

- [1] E. Camouzis, R. DeVault, and G. Papaschinopoulos, *On the recursive sequence*, Advances in Difference Equations **2005** (2005), no. 1, 31–40.
- [2] C. Çinar, *On the difference equation $x_{n+1} = x_{n-1}/-1 + x_n x_{n-1}$* , Applied Mathematics and Computation **158** (2004), no. 3, 813–816.
- [3] ———, *On the positive solutions of the difference equation $x_{n+1} = ax_{n-1}/1 + bx_n x_{n-1}$* , Applied Mathematics and Computation **156** (2004), no. 2, 587–590.
- [4] ———, *On the positive solutions of the difference equation $x_{n+1} = x_{n-1}/1 + x_n x_{n-1}$* , Applied Mathematics and Computation **150** (2004), no. 1, 21–24.
- [5] E. M. Elabbasy, H. El-Metwally, and E. M. Elsayed, *On the periodic nature of some max-type difference equations*, International Journal of Mathematics and Mathematical Sciences **2005** (2005), no. 14, 2227–2239.
- [6] ———, *On the Difference Equation $x_{n+1} = \alpha x_{n-k}/\beta + \gamma \prod_{i=0}^k x_{n-i}$* , to appear in Journal of Concrete and Applicable Mathematics.
- [7] H. El-Metwally, E. A. Grove, and G. Ladas, *A global convergence result with applications to periodic solutions*, Journal of Mathematical Analysis and Applications **245** (2000), no. 1, 161–170.
- [8] H. El-Metwally, E. A. Grove, G. Ladas, and H. D. Voulov, *On the global attractivity and the periodic character of some difference equations*, Journal of Difference Equations and Applications **7** (2001), no. 6, 837–850.
- [9] V. L. Kocić and G. Ladas, *Global Behavior of Nonlinear Difference Equations of Higher Order with Applications*, Mathematics and Its Applications, vol. 256, Kluwer Academic, Dordrecht, 1993.
- [10] M. R. S. Kulenovic and G. Ladas, *Dynamics of Second Order Rational Difference Equations with Open Problems and Conjectures*, Chapman & Hall/CRC, Florida, 2001.

10 On the difference equation $x_{n+1} = ax_n - bx_n/(cx_n - dx_{n-1})$

- [11] M. Migda, A. Musielak, and E. Schmeidel, *On a class of fourth-order nonlinear difference equations*, Advances in Difference Equations **2004** (2004), no. 1, 23–36.
- [12] Ch. G. Philos and I. K. Purnaras, *An asymptotic result for some delay difference equations with continuous variable*, Advances in Difference Equations **2004** (2004), no. 1, 1–10.
- [13] T. Sun, H. Xi, and L. Hong, *On the system of rational difference equations $x_{n+1} = f(x_n, y_{n-k})$, $y_{n+1} = f(y_n, x_{n-k})$* , Advances in Difference Equations **2006** (2006), Article ID 16949, 7 pages.

E. M. Elabbasy: Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt

E-mail address: emelabbasy@mans.edu.eg

H. El-Metwally: Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt

E-mail address: helmetwally@mans.edu.eg

E. M. Elsayed: Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt

E-mail address: emelsayed@mans.edu.eg

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk