

Research Article

On a k -Order System of Lyness-Type Difference Equations

G. Papaschinopoulos, C. J. Schinas, and G. Stefanidou

Received 17 January 2007; Revised 24 April 2007; Accepted 14 June 2007

Recommended by John R. Graef

We consider the following system of Lyness-type difference equations: $x_1(n+1) = (a_k x_k(n) + b_k)/x_{k-1}(n-1)$, $x_2(n+1) = (a_1 x_1(n) + b_1)/x_k(n-1)$, $x_i(n+1) = (a_{i-1} x_{i-1}(n) + b_{i-1})/x_{i-2}(n-1)$, $i = 3, 4, \dots, k$, where $a_i, b_i, i = 1, 2, \dots, k$, are positive constants, $k \geq 3$ is an integer, and the initial values are positive real numbers. We study the existence of invariants, the boundedness, the persistence, and the periodicity of the positive solutions of this system.

Copyright © 2007 G. Papaschinopoulos et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Difference equations and systems of difference equations have many applications in biology, economy, and other sciences. So there exist many papers concerning systems of difference equations (see [1–10] and the references cited therein).

In [11], Kocić and Ladas investigated the existence of invariants, the boundedness, the persistence, the periodicity, and the oscillation of the positive solutions of the Lyness difference equation

$$x_{n+1} = \frac{x_n + A}{x_{n-1}}, \quad n = 0, 1, \dots, \quad (1.1)$$

where A is a positive constant and the initial conditions x_{-1}, y_{-1}, x_0, y_0 are positive real numbers.

2 Advances in Difference Equations

In [6–8], the authors studied the behavior of the positive solutions of the system of two Lyness difference equations

$$x_{n+1} = \frac{b y_n + c}{x_{n-1}}, \quad y_{n+1} = \frac{d x_n + e}{y_{n-1}}, \quad n = 0, 1, \dots, \quad (1.2)$$

where b, c, d, e are positive constants and the initial conditions x_{-1}, y_{-1}, x_0, y_0 are positive numbers.

Now in this paper, we consider the system of difference equations:

$$\begin{aligned} x_1(n+1) &= \frac{a_k x_k(n) + b_k}{x_{k-1}(n-1)}, \\ x_2(n+1) &= \frac{a_1 x_1(n) + b_1}{x_k(n-1)}, \\ x_i(n+1) &= \frac{a_{i-1} x_{i-1}(n) + b_{i-1}}{x_{i-2}(n-1)}, \quad i = 3, 4, \dots, k, \end{aligned} \quad (1.3)$$

where $a_i, b_i, i = 1, 2, \dots, k$, are positive constant numbers, $k \geq 3$ is an integer, and the initial values $x_i(-1), x_i(0)$, $i = 1, 2, \dots, k$, are positive real numbers. For simplicity, system (1.3) can be written as follows:

$$x_i(n+1) = \frac{a_{i-1} x_{i-1}(n) + b_{i-1}}{x_{i-2}(n-1)}, \quad i = 1, 2, \dots, k, \quad (1.4)$$

where

$$a_0 = a_k, \quad b_0 = b_k, \quad x_j(n) = x_{k+j}(n), \quad j = -1, 0, n = -1, 0, \dots. \quad (1.5)$$

We study the existence of invariants, the boundedness, the persistence, and the periodicity of the positive solutions of the system (1.3).

2. Boundedness and persistence

In this section, we study the boundedness and the persistence of the positive solutions of (1.3). For this goal, we show the following proposition in which we find conditions so that system (1.3) has an invariant.

PROPOSITION 2.1. *Let $k \geq 3$ and*

$$\begin{aligned} \lambda_{k+i} &= \lambda_i, \quad i \in \{-2, -1, 0, 1, 2, 3, 4\}, \\ a_{k+i} &= a_i, \quad i \in \{-3, -2, -1, 0, 1\}, \\ b_{k+i} &= b_i, \quad i \in \{-2, -1, 0\}. \end{aligned} \quad (2.1)$$

Assume that the system of $2k$ equations, with k unknowns $\lambda_1, \lambda_2, \dots, \lambda_k$ of the form

$$\begin{aligned} \lambda_{i+2} b_{i-1} + \lambda_{i+3} a_i a_{i-1} &= \lambda_{i-2} b_{i-3} + \lambda_{i-3} a_{i-4} a_{i-3}, \quad i \in \{1, 2, \dots, k\}, \\ \lambda_{i+4} a_{i+1} b_i &= \lambda_{i-1} a_{i-2} b_{i-1}, \quad i \in \{1, 2, \dots, k\}, \end{aligned} \quad (2.2)$$

has a nontrivial solution $\lambda_1, \lambda_2, \dots, \lambda_k$. Then system (1.3) has an invariant of the form

$$\begin{aligned}
 I_n = & \sum_{i=1}^k \lambda_{i+2} x_i(n) + \sum_{i=1}^k \lambda_{i+2} x_i(n-1) \\
 & + \sum_{i=1}^k (\lambda_i b_{i-1} + \lambda_{i-1} a_{i-1} a_{i-2}) \frac{1}{x_i(n)} + \sum_{i=1}^k (\lambda_i b_{i-1} + \lambda_{i-1} a_{i-1} a_{i-2}) \frac{1}{x_i(n-1)} \\
 & + \sum_{i=1}^k \lambda_{i-1} a_{i-2} b_{i-1} \frac{1}{x_i(n) x_{i-1}(n-1)} + \sum_{i=1}^k \lambda_i a_{i-1} \frac{x_{i-1}(n-1)}{x_i(n)} \\
 & + \sum_{i=1}^k \lambda_{i+3} a_i \frac{x_i(n)}{x_{i-1}(n-1)}. \tag{2.3}
 \end{aligned}$$

Proof. From (1.5), (1.4), (2.1), (2.2), and (2.3), we have

$$\begin{aligned}
 I_{n+1} = & \sum_{i=1}^k \lambda_{i+2} a_{i-1} \frac{x_{i-1}(n)}{x_{i-2}(n-1)} + \sum_{i=1}^k \lambda_{i+2} b_{i-1} \frac{1}{x_{i-2}(n-1)} + \sum_{i=1}^k \lambda_{i+2} x_i(n) \\
 & + \sum_{i=1}^k \left(\lambda_i b_{i-1} + \lambda_{i-1} a_{i-1} a_{i-2} + \lambda_{i-1} a_{i-2} b_{i-1} \frac{1}{x_{i-1}(n)} + \lambda_i a_{i-1} x_{i-1}(n) \right) \\
 & \times \frac{x_{i-2}(n-1)}{a_{i-1} x_{i-1}(n) + b_{i-1}} \\
 & + \sum_{i=1}^k (\lambda_i b_{i-1} + \lambda_{i-1} a_{i-1} a_{i-2}) \frac{1}{x_i(n)} + \sum_{i=1}^k \lambda_{i+3} a_i a_{i-1} \frac{1}{x_{i-2}(n-1)} \\
 & + \sum_{i=1}^k \lambda_{i+3} a_i b_{i-1} \frac{1}{x_{i-1}(n) x_{i-2}(n-1)} \tag{2.4} \\
 = & \sum_{i=1}^k \lambda_{i+2} x_i(n) + \sum_{i=1}^k \lambda_i x_{i-2}(n-1) \\
 & + \sum_{i=1}^k (\lambda_i b_{i-1} + \lambda_{i-1} a_{i-1} a_{i-2}) \frac{1}{x_i(n)} + \sum_{i=1}^k (\lambda_{i+2} b_{i-1} + \lambda_{i+3} a_i a_{i-1}) \frac{1}{x_{i-2}(n-1)} \\
 & + \sum_{i=1}^k \lambda_{i+3} a_i b_{i-1} \frac{1}{x_{i-1}(n) x_{i-2}(n-1)} + \sum_{i=1}^k \lambda_{i-1} a_{i-2} \frac{x_{i-2}(n-1)}{x_{i-1}(n)} \\
 & + \sum_{i=1}^k \lambda_{i+2} a_{i-1} \frac{x_{i-1}(n)}{x_{i-2}(n-1)} = I_n.
 \end{aligned}$$

This completes the proof of the proposition. \square

4 Advances in Difference Equations

COROLLARY 2.2. *Let $k = 3$. Then system (1.3) for $k = 3$ has the following invariant:*

$$\begin{aligned}
I_n = & b_1x_1(n) + b_2x_2(n) + b_3x_3(n) + b_1x_1(n-1) + b_2x_2(n-1) \\
& + b_3x_3(n-1) + (b_2b_3 + b_1a_2a_3)\frac{1}{x_1(n)} + (b_3b_1 + b_2a_3a_1)\frac{1}{x_2(n)} \\
& + (b_1b_2 + b_3a_1a_2)\frac{1}{x_3(n)} + (b_2b_3 + b_1a_2a_3)\frac{1}{x_1(n-1)} \\
& + (b_3b_1 + b_2a_3a_1)\frac{1}{x_2(n-1)} + (b_1b_2 + b_3a_1a_2)\frac{1}{x_3(n-1)} \quad (2.5) \\
& + b_1a_2b_3\frac{1}{x_1(n)x_3(n-1)} + b_2a_3b_1\frac{1}{x_2(n)x_1(n-1)} \\
& + b_3a_1b_2\frac{1}{x_3(n)x_2(n-1)} + b_2a_3\frac{x_3(n-1)}{x_1(n)} + b_3a_1\frac{x_1(n-1)}{x_2(n)} \\
& + b_1a_2\frac{x_2(n-1)}{x_3(n)} + b_2a_1\frac{x_1(n)}{x_3(n-1)} + b_3a_2\frac{x_2(n)}{x_1(n-1)} + b_1a_3\frac{x_3(n)}{x_2(n-1)}.
\end{aligned}$$

Proof. From (2.1) and (2.2), we get $\lambda_2b_2 = \lambda_1b_3$, $\lambda_3b_3 = \lambda_2b_1$, $\lambda_1b_1 = \lambda_3b_2$. We set $\lambda_1 = b_2$, $\lambda_2 = b_3$, $\lambda_3 = b_1$. Then from (2.3), the proof follows immediately. \square

COROLLARY 2.3. *Let $k = 4$. Suppose that*

$$b_1 = b_2 = b_3 = b_4 = b. \quad (2.6)$$

Then system (1.3) for $k = 4$ has an invariant of the form

$$\begin{aligned}
I_n = & a_1x_1(n) + a_2x_2(n) + a_3x_3(n) + a_4x_4(n) + a_1x_1(n-1) \\
& + a_2x_2(n-1) + a_3x_3(n-1) + a_4x_4(n-1) + (a_3b + a_4a_2a_3)\frac{1}{x_1(n)} \\
& + (a_4b + a_4a_3a_1)\frac{1}{x_2(n)} + (a_1b + a_4a_1a_2)\frac{1}{x_3(n)} \\
& + (a_2b + a_3a_1a_2)\frac{1}{x_4(n)} + (a_3b + a_4a_2a_3)\frac{1}{x_1(n-1)} + (a_4b + a_4a_3a_1)\frac{1}{x_2(n-1)} \\
& + (a_1b + a_4a_1a_2)\frac{1}{x_3(n-1)} + (a_2b + a_3a_1a_2)\frac{1}{x_4(n-1)} \\
& + a_3a_2b\frac{1}{x_1(n)x_4(n-1)} + a_3a_4b\frac{1}{x_2(n)x_1(n-1)}
\end{aligned}$$

$$\begin{aligned}
& + a_1 a_4 b \frac{1}{x_3(n)x_2(n-1)} + a_1 a_2 b \frac{1}{x_4(n)x_3(n-1)} \\
& + a_1 a_4 \frac{x_1(n-1)}{x_2(n)} + a_1 a_2 \frac{x_2(n-1)}{x_3(n)} + a_3 a_2 \frac{x_3(n-1)}{x_4(n)} + a_3 a_4 \frac{x_4(n-1)}{x_1(n)} \\
& + a_1 a_4 \frac{x_4(n)}{x_3(n-1)} + a_3 a_4 \frac{x_3(n)}{x_2(n-1)} + a_3 a_2 \frac{x_2(n)}{x_1(n-1)} + a_1 a_2 \frac{x_1(n)}{x_4(n-1)}.
\end{aligned} \tag{2.7}$$

Proof. From (2.1), (2.2), and (2.6), we obtain

$$\begin{aligned}
\lambda_2 b + \lambda_3 a_4 a_3 &= \lambda_2 b + \lambda_1 a_4 a_1, \\
\lambda_1 b + \lambda_2 a_2 a_3 &= \lambda_1 b + \lambda_4 a_4 a_3, \\
\lambda_3 b + \lambda_4 a_4 a_1 &= \lambda_3 b + \lambda_2 a_2 a_1, \\
\lambda_4 b + \lambda_1 a_2 a_1 &= \lambda_4 b + \lambda_3 a_2 a_3, \\
\lambda_1 a_4 b &= \lambda_2 a_3 b, \\
\lambda_2 a_1 b &= \lambda_3 a_4 b, \\
\lambda_3 a_2 b &= \lambda_4 a_1 b, \\
\lambda_4 a_3 b &= \lambda_1 a_2 b.
\end{aligned} \tag{2.8}$$

We set in (2.8) $\lambda_1 = a_3$, $\lambda_2 = a_4$, $\lambda_3 = a_1$, $\lambda_4 = a_2$. Then from (2.3), the proof follows immediately. \square

COROLLARY 2.4. Consider system (1.3), where $k = 5$. Suppose that

$$\begin{aligned}
a_4 a_5 &= b_2, \\
a_3 a_2 &= b_5, \\
a_5 a_1 &= b_3, \\
a_4 a_3 &= b_1, \\
a_1 a_2 &= b_4.
\end{aligned} \tag{2.9}$$

Then system (1.3), with $k = 5$, has an invariant of the form

$$\begin{aligned}
I_n &= \lambda_3 x_1(n) + \lambda_4 x_2(n) + \lambda_5 x_3(n) + \lambda_1 x_4(n) + \lambda_2 x_5(n) \\
& + \lambda_3 x_1(n-1) + \lambda_4 x_2(n-1) + \lambda_5 x_3(n-1) \\
& + \lambda_1 x_4(n-1) + \lambda_2 x_5(n-1) \\
& + (\lambda_1 a_2 a_3 + \lambda_5 a_4 a_5) \frac{1}{x_1(n)} + (\lambda_2 a_3 a_4 + \lambda_1 a_5 a_1) \frac{1}{x_2(n)}
\end{aligned}$$

6 Advances in Difference Equations

$$\begin{aligned}
& + (\lambda_3 a_4 a_5 + \lambda_2 a_1 a_2) \frac{1}{x_3(n)} + (\lambda_4 a_1 a_5 + \lambda_3 a_2 a_3) \frac{1}{x_4(n)} \\
& + (\lambda_5 a_1 a_2 + \lambda_4 a_3 a_4) \frac{1}{x_5(n)} + (\lambda_1 a_2 a_3 + \lambda_5 a_4 a_5) \frac{1}{x_1(n-1)} \\
& + (\lambda_2 a_3 a_4 + \lambda_1 a_5 a_1) \frac{1}{x_2(n-1)} + (\lambda_3 a_4 a_5 + \lambda_2 a_1 a_2) \frac{1}{x_3(n-1)} \\
& + (\lambda_4 a_1 a_5 + \lambda_3 a_2 a_3) \frac{1}{x_4(n-1)} + (\lambda_5 a_1 a_2 + \lambda_4 a_3 a_4) \frac{1}{x_5(n-1)} \\
& + \lambda_5 a_4 a_3 a_2 \frac{1}{x_1(n)x_5(n-1)} + \lambda_1 a_5 a_4 a_3 \frac{1}{x_2(n)x_1(n-1)} \\
& + \lambda_2 a_1 a_4 a_5 \frac{1}{x_3(n)x_2(n-1)} + \lambda_3 a_2 a_5 a_1 \frac{1}{x_4(n)x_3(n-1)} \\
& + \lambda_4 a_3 a_1 a_2 \frac{1}{x_5(n)x_4(n-1)} \\
& + \lambda_1 a_5 \frac{x_5(n-1)}{x_1(n)} + \lambda_2 a_1 \frac{x_1(n-1)}{x_2(n)} + \lambda_3 a_2 \frac{x_2(n-1)}{x_3(n)} \\
& + \lambda_4 a_3 \frac{x_3(n-1)}{x_4(n)} + \lambda_5 a_4 \frac{x_4(n-1)}{x_5(n)} \\
& + \lambda_4 a_1 \frac{x_1(n)}{x_5(n-1)} + \lambda_5 a_2 \frac{x_2(n)}{x_1(n-1)} + \lambda_1 a_3 \frac{x_3(n)}{x_2(n-1)} \\
& + \lambda_2 a_4 \frac{x_4(n)}{x_3(n-1)} + \lambda_3 a_5 \frac{x_5(n)}{x_4(n-1)}, \\
\end{aligned} \tag{2.10}$$

where λ_i , $i = 1, 2, 3, 4, 5$, are real numbers.

Proof. Using (2.1), (2.2), and (2.9), we get

$$\begin{aligned}
\lambda_1 a_1 a_5 + \lambda_2 a_4 a_3 &= \lambda_2 a_3 a_4 + \lambda_1 a_5 a_1, \\
\lambda_2 a_2 a_1 + \lambda_3 a_4 a_5 &= \lambda_3 a_4 a_5 + \lambda_2 a_1 a_2, \\
\lambda_5 a_4 a_5 + \lambda_1 a_3 a_2 &= \lambda_1 a_2 a_3 + \lambda_5 a_4 a_5, \\
\lambda_3 a_3 a_2 + \lambda_4 a_1 a_5 &= \lambda_4 a_1 a_5 + \lambda_3 a_2 a_3, \\
\lambda_4 a_4 a_3 + \lambda_5 a_2 a_1 &= \lambda_5 a_1 a_2 + \lambda_4 a_3 a_4, \\
\lambda_1 a_3 a_4 a_5 &= \lambda_1 a_3 a_4 a_5, \\
\lambda_2 a_4 a_5 a_1 &= \lambda_2 a_4 a_5 a_1, \\
\lambda_3 a_5 a_1 a_2 &= \lambda_3 a_5 a_1 a_2, \\
\lambda_4 a_1 a_2 a_3 &= \lambda_4 a_1 a_2 a_3, \\
\lambda_5 a_2 a_3 a_4 &= \lambda_5 a_2 a_3 a_4,
\end{aligned} \tag{2.11}$$

which are satisfied for any real numbers λ_i , $i = 1, 2, 3, 4, 5$. Then from (2.3), the corollary is proved. \square

3. Periodicity

We study the periodicity of the positive solutions of (1.3) by investigating three cases: $k = 3$, $k = 4$, and $k \in \{5, 6, \dots\}$. For the first case, we show the following proposition.

PROPOSITION 3.1. *Consider system (1.3) for $k = 3$. If*

$$\begin{aligned} a_1 &= a_2 = a_3 = a, \\ b_1 &= b_2 = b_3 = b, \\ a^2 &= b, \end{aligned} \tag{3.1}$$

then every positive solution of system (1.3) is periodic of period 15.

Proof. We have

$$\begin{aligned} x_1(n+5) &= \frac{ax_3(n+4) + a^2}{x_2(n+3)} = \frac{a((ax_2(n+3) + a^2)/x_1(n+2)) + a^2}{x_2(n+3)} \\ &= \frac{a^2x_2(n+3) + a^3 + a^2x_1(n+2)}{x_1(n+2)x_2(n+3)} \\ &= \frac{a^2((ax_1(n+2) + a^2)/x_3(n+1)) + a^3 + a^2x_1(n+2)}{x_1(n+2)((ax_1(n+2) + a^2)/x_3(n+1))} \\ &= \frac{a^3x_1(n+2) + a^4 + a^3x_3(n+1) + a^2x_1(n+2)x_3(n+1)}{x_1(n+2)[ax_1(n+2) + a^2]} \\ &= \frac{[ax_1(n+2) + a^2][ax_3(n+1) + a^2]}{x_1(n+2)[ax_1(n+2) + a^2]} \\ &= \frac{ax_3(n+1) + a^2}{x_1(n+2)} = x_2(n). \end{aligned} \tag{3.2}$$

Working in a similar way, we can prove that

$$\begin{aligned} x_2(n+5) &= x_3(n), \\ x_3(n+5) &= x_1(n). \end{aligned} \tag{3.3}$$

Thus,

$$x_1(n+15) = x_2(n+10) = x_3(n+5) = x_1(n). \tag{3.4}$$

Similarly,

$$\begin{aligned} x_2(n+15) &= x_2(n), \\ x_3(n+15) &= x_3(n), \end{aligned} \tag{3.5}$$

and the proof of the proposition is complete. \square

8 Advances in Difference Equations

In the sequel, we prove the following proposition which concerns the case $k = 4$.

PROPOSITION 3.2. *Consider system (1.3) for $k = 4$. If*

$$\begin{aligned} a_1 &= a_2 = a_3 = a_4 = a, \\ b_1 &= b_2 = b_3 = b_4 = b, \\ a^2 &= b, \end{aligned} \tag{3.6}$$

then every positive solution of system (1.3) is periodic of period 20.

Proof. We have

$$\begin{aligned} x_1(n+5) &= \frac{ax_4(n+4) + a^2}{x_3(n+3)} = \frac{a((ax_3(n+3) + a^2)/x_2(n+2)) + a^2}{x_3(n+3)} \\ &= \frac{a^2x_3(n+3) + a^3 + a^2x_2(n+2)}{x_2(n+2)x_3(n+3)} \\ &= \frac{a^2((ax_2(n+2) + a^2)/x_1(n+1)) + a^3 + a^2x_2(n+2)}{x_2(n+2)((ax_2(n+2) + a^2)/x_1(n+1))} \\ &= \frac{a^3x_2(n+2) + a^4 + a^3x_1(n+1) + a^2x_1(n+1)x_2(n+2)}{x_2(n+2)[ax_2(n+2) + a^2]} \\ &= \frac{[ax_2(n+2) + a^2][ax_1(n+1) + a^2]}{x_2(n+2)[ax_2(n+2) + a^2]} \\ &= \frac{ax_1(n+1) + a^2}{x_2(n+2)} = x_4(n). \end{aligned} \tag{3.7}$$

Arguing as above, we can show that

$$\begin{aligned} x_2(n+5) &= x_1(n), \\ x_3(n+5) &= x_2(n), \\ x_4(n+5) &= x_3(n). \end{aligned} \tag{3.8}$$

So,

$$x_1(n+20) = x_4(n+15) = x_3(n+10) = x_2(n+5) = x_1(n). \tag{3.9}$$

Similarly,

$$\begin{aligned} x_2(n+20) &= x_2(n), \\ x_3(n+20) &= x_3(n), \\ x_4(n+20) &= x_4(n), \end{aligned} \tag{3.10}$$

and the proof of the proposition is complete. \square

Finally, we study the case $k \in \{5, 6, \dots\}$. To this end, we have at first to prove the following lemma.

LEMMA 3.3. *Let $k \geq 5$. If*

$$a_1 = a_2 = \dots = a_k = a, \quad b_1 = b_2 = \dots = b_k = b, \quad a^2 = b \quad (3.11)$$

then

$$\begin{aligned} x_i(n+5) &= x_{k-5+i}(n), \quad i \in \{1, 2, \dots, 5\}, \\ x_i(n+5) &= x_{i-5}(n), \quad i \in \{6, 7, \dots, k\}. \end{aligned} \quad (3.12)$$

Proof. From (1.3), we have

$$\begin{aligned} x_1(n+5) &= \frac{ax_k(n+4) + a^2}{x_{k-1}(n+3)} = \frac{a((ax_{k-1}(n+3) + a^2)/x_{k-2}(n+2)) + a^2}{x_{k-1}(n+3)} \\ &= \frac{a^2x_{k-1}(n+3) + a^3 + a^2x_{k-2}(n+2)}{x_{k-2}(n+2)x_{k-1}(n+3)} \\ &= \frac{a^2((ax_{k-2}(n+2) + a^2)/x_{k-3}(n+1)) + a^3 + a^2x_{k-2}(n+2)}{x_{k-2}(n+2)((ax_{k-2}(n+2) + a^2)/x_{k-3}(n+1))} \\ &= \frac{a^3x_{k-2}(n+2) + a^4 + a^3x_{k-3}(n+1) + a^2x_{k-2}(n+2)x_{k-3}(n+1)}{x_{k-2}(n+2)(ax_{k-2}(n+2) + a^2)} \\ &= \frac{(ax_{k-2}(n+2) + a^2)(ax_{k-3}(n+1) + a^2)}{x_{k-2}(n+2)(ax_{k-2}(n+2) + a^2)}. \end{aligned} \quad (3.13)$$

Then since, from (1.3),

$$x_{k-4}(n) = \frac{ax_{k-3}(n+1) + a^2}{x_{k-2}(n+2)}, \quad (3.14)$$

it follows that

$$x_1(n+5) = x_{k-4}(n). \quad (3.15)$$

Similarly, we can prove that

$$x_i(n+5) = x_{k-5+i}(n), \quad i \in \{2, 3, \dots, 5\}. \quad (3.16)$$

Let $i \in \{6, 7, \dots, k\}$. Then

$$\begin{aligned}
x_i(n+5) &= \frac{ax_{i-1}(n+4) + a^2}{x_{i-2}(n+3)} = \frac{a((ax_{i-2}(n+3) + a^2)/x_{i-3}(n+2)) + a^2}{x_{i-2}(n+3)} \\
&= \frac{a^2x_{i-2}(n+3) + a^3 + a^2x_{i-3}(n+2)}{x_{i-2}(n+3)x_{i-3}(n+2)} \\
&= \frac{a^2((ax_{i-3}(n+2) + a^2)/x_{i-4}(n+1)) + a^3 + a^2x_{i-3}(n+2)}{x_{i-3}(n+2)((ax_{i-3}(n+2) + a^2)/x_{i-4}(n+1))} \\
&= \frac{a^3x_{i-3}(n+2) + a^4 + a^3x_{i-4}(n+1) + a^2x_{i-3}(n+2)x_{i-4}(n+1)}{x_{i-3}(n+2)(ax_{i-3}(n+2) + a^2)} \\
&= \frac{(ax_{i-3}(n+2) + a^2)(ax_{i-4}(n+1) + a^2)}{x_{i-3}(n+2)(ax_{i-3}(n+2) + a^2)}.
\end{aligned} \tag{3.17}$$

Then since, from (1.3),

$$x_{i-5}(n) = \frac{ax_{i-4}(n+1) + a^2}{x_{i-3}(n+2)}, \tag{3.18}$$

it follows that

$$x_i(n+5) = x_{i-5}(n), \quad i \in \{6, 7, \dots, k\}. \tag{3.19}$$

Now we can show the following proposition. \square

PROPOSITION 3.4. *Consider system (1.3), where $k \geq 5$. Assume that relations (3.11) hold. Then the following statements are true.*

- (i) *Every positive solution of system (1.3) is periodic of period k if $k = 5r$, $r = 1, 2, \dots$.*
- (ii) *Every positive solution of system (1.3) is periodic of period $5k$ if $k \neq 5r$, $r = 1, 2, \dots$.*

Proof. Consider an arbitrary solution $(x_1(n), \dots, x_k(n))$ of (1.3).

- (i) Suppose that $k = 5r$, $r = 1, 2, \dots$. Then from (3.12), we have

$$\begin{aligned}
x_i(n+5) &= x_{5r-5+i}(n), \quad i \in \{1, 2, \dots, 5\}, \\
x_i(n+5) &= x_{i-5}(n), \quad i \in \{6, 7, \dots, 5r\}.
\end{aligned} \tag{3.20}$$

We claim that for $i = 1, 2, \dots, 5$,

$$x_i(n+5s) = x_{5r-5s+i}(n), \quad s = 1, 2, \dots, r. \tag{3.21}$$

From (3.20), it is obvious that (3.21) is true for $s = 1$. Suppose that for $i = 1, 2, \dots, 5$, relation (3.21) is true for $s = 1, 2, \dots, r-1$. Then since $6 \leq 5r-5s+i \leq 5r$, from (3.20) and (3.21), we get for $i = 1, 2, \dots, 5$,

$$x_i(n+5+5s) = x_{5r-5s+i}(n+5) = x_{5r-5(s+1)+i}(n), \tag{3.22}$$

and so (3.21) is true. Then from (3.21) for $s = r$, we have

$$x_i(n + 5r) = x_i(n), \quad i = 1, 2, \dots, 5. \quad (3.23)$$

Therefore the sequences $x_i(n)$, $i = 1, 2, \dots, 5$ are periodic of period 5. Then from (3.20), all the sequences $x_i(n)$, $i = 1, 2, \dots, k$, are periodic of period k .

(ii) Suppose that $k \neq 5r$, $r = 1, 2, \dots$. Let $k = 5r + 1$, $r = 1, 2, \dots$. Then from (3.12), we have

$$\begin{aligned} x_i(n + 5) &= x_{5r-4+i}(n), \quad i \in \{1, 2, \dots, 5\}, \\ x_i(n + 5) &= x_{i-5}(n), \quad i \in \{6, 7, \dots, 5r + 1\}. \end{aligned} \quad (3.24)$$

Applying (3.24) and using the same argument to show (3.21), we can prove that for $i = 1, 2, \dots, 5$

$$x_i(n + 5s) = x_{5r-5s+i+1}(n), \quad s = 1, 2, \dots, r. \quad (3.25)$$

So from (3.24) and (3.25) for $i = 1$, $s = r$, we get

$$\begin{aligned} x_1(n + 25r + 5) &= x_2(n + 20r + 5) = x_3(n + 15r + 5) = x_4(n + 10r + 5), \\ x_5(n + 5r + 5) &= x_6(n + 5) = x_1(n). \end{aligned} \quad (3.26)$$

Therefore $x_1(n)$ is a periodic sequence of period $5(5r + 1) = 5k$. Hence by (3.24), all the sequences $x_i(n)$, $i = 1, 2, \dots, k$, are periodic of period $5k$.

Let $k = 5r + 2$. Then from (3.12), we have

$$\begin{aligned} x_i(n + 5) &= x_{5r-3+i}(n), \quad i \in \{1, 2, \dots, 5\}, \\ x_i(n + 5) &= x_{i-5}(n), \quad i \in \{6, 7, \dots, 5r + 2\}. \end{aligned} \quad (3.27)$$

Then from (3.27) and using the same argument to prove (3.21), we can prove that for $i = 1, 2, \dots, 5$,

$$x_i(n + 5s) = x_{5r-5s+i+2}(n), \quad s = 1, 2, \dots, r. \quad (3.28)$$

Then from (3.27) and (3.28) for $i = 1$, $s = r$, we get

$$\begin{aligned} x_1(n + 25r + 10) &= x_3(n + 20r + 10) = x_5(n + 15r + 10) \\ &= x_7(n + 10r + 10) = x_2(n + 10r + 5) = x_4(n + 5r + 5) \\ &= x_6(n + 5) = x_1(n), \end{aligned} \quad (3.29)$$

which implies that $x_1(n)$ is a periodic sequence of period $5k$. Then by relations (3.27), we can prove that the sequences $x_i(n)$, $i = 2, 3, \dots, k$, are periodic of period $5k$.

Let $k = 5r + 3$. Then from (3.12), we have

$$\begin{aligned} x_i(n + 5) &= x_{5r-2+i}(n), \quad i \in \{1, 2, \dots, 5\}, \\ x_i(n + 5) &= x_{i-5}(n), \quad i \in \{6, 7, \dots, 5r + 3\}. \end{aligned} \quad (3.30)$$

Then from (3.30) and using the same argument to show (3.21), we can prove that for $i = 1, 2, \dots, 5$,

$$x_i(n + 5s) = x_{5r-5s+i+3}(n), \quad s = 1, 2, \dots, r. \quad (3.31)$$

Then from (3.30) and (3.31) for $i = 1, s = r$, we get

$$\begin{aligned} x_1(n + 25r + 15) &= x_4(n + 20r + 15) = x_7(n + 15r + 15) \\ &= x_2(n + 15r + 10) = x_5(n + 10r + 10) \\ &= x_8(n + 5r + 10) = x_3(n + 5r + 5) = x_6(n + 5) = x_1(n), \end{aligned} \quad (3.32)$$

which implies that $x_1(n)$ is a periodic sequence of period $5(5r + 3) = 5k$. Then from relations (3.30), we can prove that the sequences $x_i(n)$, $i = 1, 2, \dots, k$, are periodic of period $5k$.

Let $k = 5r + 4$. Then from (3.12), we have

$$\begin{aligned} x_i(n + 5) &= x_{5r-1+i}(n), \quad i \in \{1, 2, \dots, 5\}, \\ x_i(n + 5) &= x_{i-5}(n), \quad i \in \{6, 7, \dots, 5r + 4\}. \end{aligned} \quad (3.33)$$

Then from (3.33) and using the same argument to show (3.21), we can prove that for $i = 1, 2, \dots, 5$,

$$x_i(n + 5s) = x_{5r-5s+i+4}(n), \quad s = 1, 2, \dots, r. \quad (3.34)$$

Then from (3.33) and (3.34) for $i = 1, s = r$, we get

$$\begin{aligned} x_1(n + 25r + 20) &= x_5(n + 20r + 20) = x_9(n + 15r + 20) \\ &= x_4(n + 15r + 15) = x_8(n + 10r + 15) = x_3(n + 10r + 10) \\ &= x_7(n + 5r + 10) = x_2(n + 5r + 5) = x_6(n + 5) = x_1(n), \end{aligned} \quad (3.35)$$

which implies that $x_1(n)$ is a periodic sequence of period $5(5r + 4) = 5k$. Then from relations (3.33), we can prove that the sequences $x_i(n)$, $i = 1, 2, \dots, k$, are periodic of period $5k$. This completes the proof of the proposition. \square

Acknowledgment

The authors would like to thank the referee for their helpful suggestions.

References

- [1] D. Clark and M. R. S. Kulenović, “A coupled system of rational difference equations,” *Computers & Mathematics with Applications*, vol. 43, no. 6-7, pp. 849–867, 2002.
- [2] D. Clark, M. R. S. Kulenović, and J. F. Selgrade, “Global asymptotic behavior of a two-dimensional difference equation modelling competition,” *Nonlinear Analysis: Theory, Methods, & Applications*, vol. 52, no. 7, pp. 1765–1776, 2003.
- [3] M. R. S. Kulenović and O. Merino, *Discrete Dynamical Systems and Difference Equations with Mathematica*, Chapman & Hall/CRC, Boca Raton, Fla, USA, 2002.

- [4] M. R. S. Kulenović and Z. Nurkanović, “Global behavior of a three-dimensional linear fractional system of difference equations,” *Journal of Mathematical Analysis and Applications*, vol. 310, no. 2, pp. 673–689, 2005.
- [5] M. R. S. Kulenović and M. Nurkanović, “Asymptotic behavior of a competitive system of linear fractional difference equations,” *Advances in Difference Equations*, vol. 2006, Article ID 19756, 13 pages, 2006.
- [6] G. Papaschinopoulos and C. J. Schinas, “Invariants for systems of two nonlinear difference equations,” *Differential Equations and Dynamical Systems*, vol. 7, no. 2, pp. 181–196, 1999.
- [7] G. Papaschinopoulos and C. J. Schinas, “On the behavior of the solutions of a system of two nonlinear difference equations,” *Communications on Applied Nonlinear Analysis*, vol. 5, no. 2, pp. 47–59, 1998.
- [8] C. J. Schinas, “Invariants for difference equations and systems of difference equations of rational form,” *Journal of Mathematical Analysis and Applications*, vol. 216, no. 1, pp. 164–179, 1997.
- [9] T. Sun, H. Xi, and L. Hong, “On the system of rational difference equations $x_{n+1} = f(x_n, y_{n-k}), y_{n+1} = f(y_n, x_{n-k})$,” *Advances in Difference Equations*, vol. 2006, Article ID 16949, 7 pages, 2006.
- [10] T. Sun and H. Xi, “On the system of rational difference equations $x_{n+1} = f(y_{n-q}, x_{n-s}), y_{n+1} = g(x_{n-t}, y_{n-p})$,” *Advances in Difference Equations*, vol. 2006, Article ID 51520, 8 pages, 2006.
- [11] V. L. Kocić and G. Ladas, *Global Behavior of Nonlinear Difference Equations of Higher Order with Applications*, vol. 256 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1993.

G. Papaschinopoulos: School of Engineering, Democritus University of Thrace,
67100 Xanthi, Greece

Email address: gpapas@ee.duth.gr

C. J. Schinas: School of Engineering, Democritus University of Thrace,
67100 Xanthi, Greece

Email address: cschinas@ee.duth.gr

G. Stefanidou: School of Engineering, Democritus University of Thrace,
67100 Xanthi, Greece

Email address: gstefmath@yahoo.gr

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru