

*Research Article*

## Existence of Solutions for Second-Order Nonlinear Impulsive Differential Equations with Periodic Boundary Value Conditions

Chuanzhi Bai and Dandan Yang

Received 12 February 2007; Revised 19 March 2007; Accepted 13 April 2007

Recommended by Kanishka Perera

We are concerned with the nonlinear second-order impulsive periodic boundary value problem  $u''(t) = f(t, u(t), u'(t))$ ,  $t \in [0, T] \setminus \{t_1\}$ ,  $u(t_1^+) = u(t_1^-) + I(u(t_1))$ ,  $u'(t_1^+) = u'(t_1^-) + J(u(t_1))$ ,  $u(0) = u(T)$ ,  $u'(0) = u'(T)$ , new criteria are established based on Schaefer's fixed-point theorem.

Copyright © 2007 C. Bai and D. Yang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### 1. Introduction

Impulsive differential equations, which arise in physics, population dynamics, economics, and so forth, are important mathematical tools for providing a better understanding of many real-world models, we refer to [1–5] and the references therein. About the applications of the theory of impulsive differential equations to different areas, for example, see [6–15]. Boundary value problems (BVPs) for impulsive differential equations and impulsive difference equations [16–20] have received special attention from many authors in recent years.

Recently, Chen et al. in [21] study the following first-order impulsive nonlinear periodic boundary value problem:

$$\begin{aligned} x'(t) &= f(t, x), \quad t \in [0, N], t \neq t_1, \\ x(t_1^+) &= x(t_1^-) + I_1(x(t_1)), \\ x(0) &= x(T), \end{aligned} \tag{1.1}$$

where  $N > 0$ ,  $t_1 \in (0, N)$ ,  $t_1$  is fixed,  $f : [0, N] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is continuous on  $(t, u) \in ([0, N] \setminus \{t_1\}) \times \mathbb{R}^n$ , and the impulse at  $t = t_1$  is given by a continuous function  $I_1 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ . They

## 2 Boundary Value Problems

investigate the existence of solutions to the problem by means of differential inequalities and Schaefer fixed point theorem. Their results complement and extend those of [22, 23] in the sense that they allow superlinear growth of the nonlinearity of  $\|f(t, p)\|$  in  $\|p\|$ .

Inspired by [21, 24, 25], in this paper, we investigate the following second-order impulsive nonlinear differential equations with periodic boundary value conditions problem:

$$\begin{aligned} u''(t) &= f(t, u(t), u'(t)), \quad t \in [0, T], t \neq t_1, \\ u(t_1^+) &= u(t_1^-) + I(u(t_1)), \\ u'(t_1^+) &= u'(t_1^-) + J(u(t_1)), \\ u(0) &= u(T), \quad u'(0) = u'(T), \end{aligned} \tag{1.2}$$

where  $T > 0$ ,  $t_1 \in (0, T)$ ,  $t_1$  is fixed,  $f : [0, T] \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is continuous on  $(t, x, y) \in ([0, T] \setminus \{t_1\}) \times \mathbb{R}^n \times \mathbb{R}^n$ , and the impulse is given at  $t_1$  by two continuous functions  $I, J : \mathbb{R}^n \rightarrow \mathbb{R}^n$ , the notations  $u(t_1^-) := \lim_{t \rightarrow t_1^-} u(t)$ ,  $u(t_1^+) := \lim_{t \rightarrow t_1^+} u(t)$ ,  $u'(t_1^+) = \lim_{t \rightarrow t_1^+} u'(t)$ , and  $u'(t_1^-) = \lim_{t \rightarrow t_1^-} u'(t)$ .

We note that we could consider impulsive BVPs with an arbitrary finite number of impulses. However, for clarity and brevity, we restrict our attention to BVPs with one impulse. In addition, the difference between the theory of one or an arbitrary finite number of impulses is quite minimal.

Our results extend those of [25] from the nonimpulsive case to the impulsive case. Our approach using differential inequalities is based on ideas in [24, 25]. Moreover, our new results complement and extend those of [26–28] in the sense that we allow superlinear growth of  $\|f(t, p, q)\|$  in  $\|p\|$  and  $\|q\|$ .

The main purpose is to establish the existence of solutions for the impulsive BVP (1.2) by employing the well-known Schaefer fixed point theorem.

**LEMMA 1.1** (see [29] (Schaefer)). *Let  $E$  be a normed linear space with  $H : E \rightarrow E$  be a compact operator. If the set*

$$S := \{x \in E \mid x = \lambda Hx, \text{ for some } \lambda \in (0, 1)\} \tag{1.3}$$

*is bounded, then  $H$  has at least one fixed point.*

The paper is formulated as follows. In Section 2, some definitions and lemmas are given. In Section 3, we establish new existence theorems for (1.2). In Section 4, an illustrative example is given to demonstrate the effectiveness of the obtained results.

## 2. Preliminaries

First, we briefly introduce some appropriate concepts connected with impulsive differential equations. Most of the following notations can be found in [30].

Assume that

$$f(t_1^+, x, y) := \lim_{t \rightarrow t_1^+} f(t, x, y), \quad f(t_1^-, x, y) := \lim_{t \rightarrow t_1^-} f(t, x, y) \tag{2.1}$$

both exist with  $f(t_1^-, x, y) = f(t_1, x, y)$ . We introduce and denote the Banach space  $\text{PC}([0, T], \mathbb{R}^n)$  by

$$\begin{aligned} \text{PC}([0, T]; \mathbb{R}^n) = \{u \in C([0, T] \setminus \{t_1\}, \mathbb{R}^n), u \text{ is left continuous at } t = t_1, \\ \text{the right-hand limit } u(t_1^+) \text{ exists}\} \end{aligned} \quad (2.2)$$

with the norm

$$\|u\|_{\text{PC}} = \sup_{t \in [0, T]} \|u(t)\|, \quad (2.3)$$

where  $\|\cdot\|$  is the usual Euclidean norm.

We define and denote the Banach space  $\text{PC}^1([0, T]; \mathbb{R}^n)$  by

$$\begin{aligned} \text{PC}^1([0, T]; \mathbb{R}^n) = \{u \in C^1([0, T] \setminus \{t_1\}, \mathbb{R}^n), u \text{ is left continuous at } t = t_1, \\ \text{the right-hand limit } u(t_1^+) \text{ exists, and the limits } u'(t_1^+), u'(t_1^-) \text{ exist}\} \end{aligned} \quad (2.4)$$

with the norm

$$\|u\|_{\text{PC}^1} = \max \{\|u\|_{\text{PC}}, \|u'\|_{\text{PC}}\}. \quad (2.5)$$

A solution to the impulsive BVP (1.2) is a function  $u \in \text{PC}^1([0, T], \mathbb{R}^n) \cap C^2([0, T] \setminus \{t_1\}, \mathbb{R}^n)$  that satisfies (1.2) for each  $t \in [0, T]$ .

Consider the following impulsive BVP with  $p \geq 0, q > 0$ :

$$\begin{aligned} u''(t) - pu'(t) - qu(t) + \sigma(t) = 0, \quad t \in [0, T], t \neq t_1, \\ u(t_1^+) = u(t_1^-) + I(u(t_1)), \\ u'(t_1^+) = u'(t_1^-) + J(u(t_1)), \\ u(0) = u(T), \quad u'(0) = u'(T), \end{aligned} \quad (2.6)$$

where  $\sigma \in \text{PC}([0, T], \mathbb{R}^n)$  is given,  $I, J : \mathbb{R}^n \rightarrow \mathbb{R}^n$  are continuous.

For convenience, we set

$$r_1 := \frac{p + \sqrt{p^2 + 4q}}{2} > 0, \quad r_2 := \frac{p - \sqrt{p^2 + 4q}}{2} < 0. \quad (2.7)$$

LEMMA 2.1.  $u \in \text{PC}^1([0, T], \mathbb{R}^n) \cap C^2([0, T] \setminus \{t_1\}, \mathbb{R}^n)$  is a solution of (2.6) if and only if  $u \in \text{PC}^1([0, T], \mathbb{R}^n)$  is a solution of the following linear impulsive integral equation:

$$u(t) = \int_0^T G(t, s) \sigma(s) ds + G(t, t_1) (-J(u(t_1))) + W(t, t_1) I(u(t_1)), \quad (2.8)$$

## 4 Boundary Value Problems

where

$$G(t, s) = \frac{1}{r_1 - r_2} \begin{cases} \frac{e^{r_1(t-s)}}{e^{r_1 T} - 1} + \frac{e^{r_2(t-s)}}{1 - e^{r_2 T}}, & 0 \leq s < t \leq T, \\ \frac{e^{r_1(T+t-s)}}{e^{r_1 T} - 1} + \frac{e^{r_2(T+t-s)}}{1 - e^{r_2 T}}, & 0 \leq t \leq s \leq T, \end{cases} \quad (2.9)$$

$$W(t, s) = \frac{1}{r_1 - r_2} \begin{cases} \frac{r_2 e^{r_1(t-s)}}{e^{r_1 T} - 1} + \frac{r_1 e^{r_2(t-s)}}{1 - e^{r_2 T}}, & 0 \leq s < t \leq T, \\ \frac{r_2 e^{r_1(T+t-s)}}{e^{r_1 T} - 1} + \frac{r_1 e^{r_2(T+t-s)}}{1 - e^{r_2 T}}, & 0 \leq t \leq s \leq T. \end{cases} \quad (2.10)$$

*Proof.* If  $u \in \text{PC}^1([0, T]; \mathbb{R}^n) \cap C^2([0, T] \setminus \{t_1\}, \mathbb{R}^n)$  is a solution of (2.6), setting

$$v(t) = u'(t) - r_2 u(t), \quad (2.11)$$

then by the first equation of (2.6), we have

$$v'(t) - r_1 v(t) = -\sigma(t), \quad t \neq t_1. \quad (2.12)$$

Multiplying (2.12) by  $e^{-r_1 t}$  and integrating on  $[0, t_1]$  and  $(t_1, T]$ , respectively, we get

$$\begin{aligned} e^{-r_1 t_1} v(t_1^-) - v(0) &= - \int_0^{t_1} \sigma(s) e^{-r_1 s} ds, \quad 0 \leq t < t_1, \\ e^{-r_1 t} v(t) - e^{-r_1 t_1} v(t_1^+) &= - \int_{t_1}^T \sigma(s) e^{-r_1 s} ds, \quad t_1 < t \leq T, \end{aligned} \quad (2.13)$$

then, we have by the second equation and third equation of (2.6) that

$$v(t) = e^{r_1 t} \left[ v(0) - \int_0^t e^{-r_1 s} \sigma(s) ds + I^* \right], \quad t \in [0, T], \quad (2.14)$$

where

$$v(0) = u'(0) - r_2 u(0), \quad I^* = (J(u(t_1)) - r_2 I(u(t_1))) e^{-r_1 t_1}. \quad (2.15)$$

Integrating (2.11), we have

$$u(t) = e^{r_2 t} \left[ u(0) + \int_0^t v(s) e^{-r_2 s} ds + I(u(t_1)) e^{-r_2 t_1} \right], \quad t \in [0, T]. \quad (2.16)$$

By some calculation, we get

$$\begin{aligned} & \int_0^t v(s) e^{-r_2 s} ds \\ &= \frac{1}{r_1 - r_2} \left[ v(0) (e^{(r_1 - r_2)t} - 1), - \int_0^t (e^{(r_1 - r_2)t} - e^{(r_1 - r_2)s}) \sigma(s) e^{-r_1 s} ds + I^* (e^{(r_1 - r_2)t} - e^{(r_1 - r_2)t_1}) \right]. \end{aligned} \quad (2.17)$$

Substituting (2.17) into (2.16), we have

$$\begin{aligned}
u(t) = & \frac{1}{r_1 - r_2} \left[ (u'(0) - r_2 u(0)) e^{r_1 t} + (r_1 u(0) - u'(0)) e^{r_2 t} \right. \\
& + \int_0^t (e^{r_2(t-s)} - e^{r_1(t-s)}) \sigma(s) ds \\
& + (J(u(t_1)) - r_2 I(u(t_1))) e^{r_1(t-t_1)} \\
& \left. - (J(u(t_1)) - r_1 I(u(t_1))) e^{r_2(t-t_1)} \right], \quad t \in [0, T].
\end{aligned} \tag{2.18}$$

By the fourth equation (boundary condition) of (2.6), we have

$$r_1 u(0) - u'(0) = \frac{1}{1 - e^{r_2 T}} \left[ \int_0^T e^{r_2(T-s)} \sigma(s) ds - (J(u(t_1)) - r_1 I(u(t_1))) e^{r_2(T-t_1)} \right], \tag{2.19}$$

$$u'(0) - r_2 u(0) = \frac{1}{e^{r_1 T} - 1} \left[ \int_0^T e^{r_1(T-s)} \sigma(s) ds - (J(u(t_1)) - r_2 I(u(t_1))) e^{r_1(T-t_1)} \right], \tag{2.20}$$

substituting (2.19) and (2.20) into (2.18), we get (2.8).

Conversely, if  $u$  is a solution to (2.8), then direct differentiation of (2.8) gives  $u''(t) = -\sigma(t) + pu'(t) + qu(t)$ ,  $t \neq t_1$ . Moreover, we have  $u(t_1^+) = u(t_1^-) + I(u(t_1))$ ,  $u'(t_1^+) = u'(t_1^-) + J(u(t_1))$ ,  $u(0) = u(T)$ , and  $u'(0) = u'(T)$ .

Note that the linear part of the periodic BVP (1.2) is not necessarily invertible, that is, we may be unable to equivalently rewrite (1.2) in the integral form. However, if we use Lemma 2.1, then impulsive BVP (1.2) may be equivalently reformulated as the impulsive integral equation.

We now introduce a mapping  $A : \text{PC}^1([0, T]; \mathbb{R}^n) \rightarrow \text{PC}([0, T]; \mathbb{R}^n)$  defined by

$$\begin{aligned}
Au(t) = & \int_0^T G(t, s) [-f(s, u(s), u'(s)) + pu'(s) + qu(s)] ds \\
& + G(t, t_1) (-J(u(t_1))) + W(t, t_1) I(u(t_1)), \quad t \in [0, T].
\end{aligned} \tag{2.21}$$

In view of Lemma 2.1, we easily know that  $u$  is a fixed point of operator  $A$  if and only if  $u$  is a solution to the impulsive boundary value problem (1.2).

It is easy to check that

$$0 \leq G(t, s) \leq G(s, s) = \frac{e^{r_1 T} - e^{r_2 T}}{(r_1 - r_2)(e^{r_1 T} - 1)(1 - e^{r_2 T})} := G_1. \tag{2.22}$$

## 6 Boundary Value Problems

By  $p \geq 0$  and  $q > 0$ , we have  $r_1 \geq -r_2 > 0$ . Thus we obtain that

$$\begin{aligned}
|W(t,s)| &\leq \frac{1}{r_1 - r_2} \begin{cases} \frac{-r_2 e^{r_1(t-s)}}{e^{r_1 T} - 1} + \frac{r_1 e^{r_2(t-s)}}{1 - e^{r_2 T}}, & 0 \leq s < t \leq T, \\ \frac{-r_2 e^{r_1(T+t-s)}}{e^{r_1 T} - 1} + \frac{r_1 e^{r_2(T+t-s)}}{1 - e^{r_2 T}}, & 0 \leq t \leq s \leq T, \end{cases} \\
&\leq \frac{r_1}{r_1 - r_2} \begin{cases} \frac{e^{r_1(t-s)}}{e^{r_1 T} - 1} + \frac{e^{r_2(t-s)}}{1 - e^{r_2 T}}, & 0 \leq s < t \leq T, \\ \frac{e^{r_1(T+t-s)}}{e^{r_1 T} - 1} + \frac{e^{r_2(T+t-s)}}{1 - e^{r_2 T}}, & 0 \leq t \leq s \leq T, \end{cases} \\
&= r_1 G(t,s) \leq r_1 G_1.
\end{aligned} \tag{2.23}$$

Since

$$\begin{aligned}
G_t(t,s) &:= \frac{\partial}{\partial t} G(t,s) = \frac{1}{r_1 - r_2} \begin{cases} \frac{r_1 e^{r_1(t-s)}}{e^{r_1 T} - 1} + \frac{r_2 e^{r_2(t-s)}}{1 - e^{r_2 T}}, & 0 \leq s < t \leq T, \\ \frac{r_1 e^{r_1(T+t-s)}}{e^{r_1 T} - 1} + \frac{r_2 e^{r_2(T+t-s)}}{1 - e^{r_2 T}}, & 0 \leq t \leq s \leq T, \end{cases} \\
W_t(t,s) &:= \frac{\partial}{\partial t} W(t,s) = \frac{1}{r_1 - r_2} \begin{cases} \frac{r_1 r_2 e^{r_1(t-s)}}{e^{r_1 T} - 1} + \frac{r_2 r_1 e^{r_2(t-s)}}{1 - e^{r_2 T}}, & 0 \leq s < t \leq T, \\ \frac{r_1 r_2 e^{r_1(T+t-s)}}{e^{r_1 T} - 1} + \frac{r_1 r_2 e^{r_2(T+t-s)}}{1 - e^{r_2 T}}, & 0 \leq t \leq s \leq T, \end{cases}
\end{aligned} \tag{2.24}$$

we easily get that

$$\begin{aligned}
|G_t(t,s)| &\leq \frac{1}{r_1 - r_2} \begin{cases} \frac{r_1 e^{r_1(t-s)}}{e^{r_1 T} - 1} + \frac{-r_2 e^{r_2(t-s)}}{1 - e^{r_2 T}}, & 0 \leq s < t \leq T, \\ \frac{r_1 e^{r_1(T+t-s)}}{e^{r_1 T} - 1} + \frac{-r_2 e^{r_2(T+t-s)}}{1 - e^{r_2 T}}, & 0 \leq t \leq s \leq T, \end{cases} \\
&\leq r_1 G(t,s) \leq r_1 G_1, \\
|W_t(t,s)| &\leq \frac{1}{r_1 - r_2} \begin{cases} \frac{-r_2 r_1 e^{r_1(t-s)}}{e^{r_1 T} - 1} + \frac{-r_2 r_1 e^{r_2(t-s)}}{1 - e^{r_2 T}}, & 0 \leq s < t \leq T, \\ \frac{-r_2 r_1 e^{r_1(T+t-s)}}{e^{r_1 T} - 1} + \frac{-r_2 r_1 e^{r_2(T+t-s)}}{1 - e^{r_2 T}}, & 0 \leq t \leq s \leq T, \end{cases} \\
&\leq r_1^2 G(t,s) \leq r_1^2 G_1.
\end{aligned} \tag{2.25}$$

Let

$$H := \max \{r_1 G_1, r_1^2 G_1\}. \tag{2.26}$$

So

$$|G_t(t, s)| \leq H, \quad |W_t(t, s)| \leq H. \quad (2.27)$$

□

LEMMA 2.2. Let  $f : [0, T] \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  and  $I, J : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be continuous. Then  $A : \text{PC}^1([0, T]; \mathbb{R}^n) \rightarrow \text{PC}^1([0, T]; \mathbb{R}^n)$  is a compact map.

*Proof.* This is similar to that of [31, Lemma 3.2]. Define two operators  $B, F$  as follows:

$$\begin{aligned} Bu(t) &= \int_0^T G(t, s) [-f(s, u(s), u'(s)) + pu'(s) + qu(s)] ds, \quad t \in [0, T], \\ Fu(t) &= G(t, t_1)(-J(u(t_1))) + W(t, t_1)I(u(t_1)), \quad t \in [0, T]. \end{aligned} \quad (2.28)$$

From the continuity of  $f$ , it is easy to see that  $B$  is compact. Since  $I, J$  are continuous, we have that  $F$  is compact. Thus  $A = B + F$  is a compact map. □

### 3. Main results

THEOREM 3.1. Suppose that  $f : [0, T] \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  and  $I, J : \mathbb{R}^n \rightarrow \mathbb{R}^n$  are continuous. If there exist nonnegative constants  $\alpha, \beta, \gamma, L_1, L_2, N$ , and  $M$  such that for each  $\lambda \in (0, 1)$ ,

$$\begin{aligned} &||f(t, x, y) - py - qx|| \leq 2\alpha[\langle x + y, f(t, x, y) \rangle + \|y\|^2] + M, \\ &(t, x, y) \in ([0, T] \setminus \{t_1\}) \times \mathbb{R}^n \times \mathbb{R}^n, \text{ where } \langle \cdot \rangle \text{ is the Euclidean inner product,} \end{aligned} \quad (3.1)$$

$$||I(x)|| \leq \beta\|x\| + L_1, \quad ||J(x)|| \leq \gamma\|x\| + L_2, \quad \forall x \in \mathbb{R}^n, \quad (3.2)$$

$$\beta + \gamma < \frac{1}{H}, \quad (3.3)$$

where  $H$  is as in (2.26), then BVP (1.2) has at least one solution.

*Proof.* From Lemma 2.2, we know that  $A$  is a compact map. In order to show that  $A$  has at least one fixed point, we apply Lemma 1.1 (Schaefer's theorem) by showing that all potential solutions to

$$u = \lambda Au, \quad \lambda \in (0, 1), \quad (3.4)$$

are bounded a priori, with the bound being independent of  $\lambda$ . Let  $u$  be a solution to (3.4), then

$$\begin{aligned} u''(t) - pu'(t) - qu(t) &= \lambda[f(t, u(t), u'(t)) - pu'(t) - qu(t)], \quad t \in [0, T], \\ u(t_1^+) &= u(t_1^-) + \lambda I(u(t_1)), \\ u'(t_1^+) &= u'(t_1^-) + \lambda J(u(t_1)), \\ u(0) &= u(T), \quad u'(0) = u'(T). \end{aligned} \quad (3.5)$$

## 8 Boundary Value Problems

By (3.1)–(3.3), (2.22) and (2.23), we obtain

$$\begin{aligned}
& \|u(t)\| = \lambda \|Au(t)\| \\
& = \left\| \int_0^T G(t,s) \lambda (f(s, u(s), u'(s)) - pu'(s) - qu(s)) ds \right. \\
& \quad \left. + \lambda G(t, t_1) (-J(u(t_1))) + \lambda W(t, t_1) I(u(t_1)) \right\| \\
& \leq G_1 \int_0^T \lambda \|f(s, u(s), u'(s)) - pu'(s) - qu(s)\| ds \\
& \quad + \lambda G_1 (\|J(u(t_1))\| + \|I(u(t_1))\|) \\
& \leq G_1 \left[ \int_0^T (2\alpha (\langle u(s) + u'(s), \lambda f(s, u(s), u'(s)) \rangle + \|u'\|^2) + M) ds \right. \\
& \quad \left. + \beta \|u(t_1)\| + L_1 + \gamma \|u(t_1)\| + L_2 \right] \quad (3.6) \\
& = G_1 \left[ \int_0^T \left( 2\alpha \left( \langle u(s) + u'(s), \lambda f(s, u(s), u'(s)) \rangle + (1-\lambda)pu'(s) \right. \right. \right. \\
& \quad \left. \left. \left. + (1-\lambda)qu(s) \rangle + \|u'(s)\|^2 \right) + M \right) ds \right. \\
& \quad \left. - \int_0^T 2\alpha \langle u(s) + u'(s), (1-\lambda)pu'(s) + (1-\lambda)qu(s) \rangle ds \right. \\
& \quad \left. + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right].
\end{aligned}$$

Since

$$\begin{aligned}
& - \int_0^T \langle u(s) + u'(s), (1-\lambda)pu'(s) + (1-\lambda)qu(s) \rangle ds \\
& = -(1-\lambda)q \int_0^T \|u(s)\|^2 ds - (1-\lambda)p \|u'(s)\|^2 ds + (1-\lambda)(p+q) \int_0^T \langle u(s), u'(s) \rangle ds \\
& \leq (1-\lambda)(p+q) \int_0^T \langle u(s), u'(s) \rangle ds = \frac{1}{2} (1-\lambda)(p+q) \int_0^T \frac{d}{ds} (\|u(s)\|^2) \\
& = \frac{1}{2} (1-\lambda)(p+q) (\|u(T)\|^2 - \|u(0)\|^2) = 0, \quad (3.7)
\end{aligned}$$

we have by (3.6) and (3.7) that

$$\begin{aligned}
& \|u(t)\| = \lambda \|Au(t)\| \\
& \leq G_1 \left[ \int_0^T \left( 2\alpha \left( \langle u(s) + u'(s), \lambda f(s, u(s), u'(s)) + (1-\lambda)p u'(s) + (1-\lambda)q u(s) \rangle \right. \right. \right. \\
& \quad \left. \left. \left. + \|u'(s)\|^2 \right) + M \right) ds + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\
& = G_1 \left[ \int_0^T \left( 2\alpha \langle u(s) + u'(s), u''(s) \rangle + \langle u(s) + u'(s), u'(s) \rangle \right. \right. \\
& \quad \left. \left. - \langle u(s), u'(s) \rangle + M \right) ds + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\
& = G_1 \left[ \int_0^T (2\alpha \langle u(s) + u'(s), u''(s) + u'(s) \rangle + M) ds + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\
& = G_1 \left[ \int_0^T \left( \alpha \frac{d}{ds} (\|u(s) + u'(s)\|^2) + M \right) ds + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\
& = G_1 \left[ \alpha \left( \|u(T) + u'(T)\|^2 - \|u(0) + u'(0)\|^2 \right) + TM + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\
& = G_1 [TM + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2].
\end{aligned} \tag{3.8}$$

Thus, taking the supremum and rearranging, we have

$$\sup_{t \in [0, T]} \|u(t)\| \leq \frac{G_1(TM + L_1 + L_2)}{1 - G_1(\beta + \gamma)}. \tag{3.9}$$

A similar calculation yields an estimate on  $u'$ : differentiating both sides of the integration equation (3.4) and taking norms yields, by (2.27), for each  $t \in [0, T]$  that

$$\sup_{t \in [0, T]} \|u'(t)\| \leq \frac{H(TM + L_1 + L_2)}{1 - H(\beta + \gamma)}, \tag{3.10}$$

where  $H$  is as in (2.26). By (3.9) and (3.10), we conclude that

$$\|u\|_{PC^1} = \max \left\{ \frac{G_1(TM + L_1 + L_2)}{1 - G_1(\beta + \gamma)}, \frac{H(TM + L_1 + L_2)}{1 - H(\beta + \gamma)} \right\} = \frac{H(TM + L_1 + L_2)}{1 - H(\beta + \gamma)}. \tag{3.11}$$

As a result, we obtain the desired bound. We see that the bound on all possible solutions to (3.4) is independent of  $\lambda$ . Applying Scheafer fixed point theorem,  $A$  has at least one fixed point, which means that (1.2) has at least one solution. We complete the proof.  $\square$

Theorem 3.1 may be suitably modified to include an alternate class of  $f$  as follows.

## 10 Boundary Value Problems

**THEOREM 3.2.** Suppose that  $f : [0, T] \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  and  $I, J : \mathbb{R}^n \rightarrow \mathbb{R}^n$  are continuous. Let the conditions of Theorem 3.1 hold with (3.1) replaced by

$$\|f(t, x, y) - py - qx\| \leq 2\alpha \langle y, f(t, x, y) \rangle + M, \quad (t, x, y) \in ([0, T] \setminus \{t_1\}) \times \mathbb{R}^n \times \mathbb{R}^n. \quad (3.12)$$

Then the impulsive BVP (1.2) has at least one solution.

The proof of Theorem 3.2 is similar to that of Theorem 3.1. It is enough to notice that (3.6) in Theorem 3.1 reduces to

$$\begin{aligned} \|u(t)\| &= \lambda \|Au(t)\| \\ &\leq G_1 \int_0^T \lambda \|f(s, u(s), u'(s)) - pu'(s) - qu(s)\| ds + \lambda G_1 (\|J(u(t_1))\| + \|I(u(t_1))\|) \\ &\leq G_1 \left[ \int_0^T (2\alpha \langle u'(s), \lambda f(s, u(s), u'(s)) \rangle + M) ds \quad (\text{use (3.12)}) \right. \\ &\quad \left. + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\ &\leq G_1 \left[ \int_0^T (2\alpha \langle u'(s), \lambda f(s, u(s), u'(s)) + (1 - \lambda)pu'(s) \rangle + M) ds \right. \\ &\quad \left. + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\ &= G_1 \left[ \int_0^T (2\alpha \langle u'(s), \lambda f(s, u(s), u'(s)) + (1 - \lambda)pu'(s) + (1 - \lambda)qu(s) \rangle + M) ds \right. \\ &\quad \left. - (1 - \lambda)q \int_0^T 2\alpha \langle u'(s), u(s) \rangle ds + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\ &= G_1 \left[ \int_0^T (2\alpha \langle u'(s), u''(s) \rangle + M) ds + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\ &= G_1 \left[ \int_0^T \left( \alpha \frac{d}{ds} (\|u'(s)\|^2) + M \right) ds + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\ &= G_1 \left[ \alpha (\|u'(T)\|^2 - \|u'(0)\|^2) + TM + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2 \right] \\ &= G_1 [TM + (\beta + \gamma) \|u(t_1)\| + L_1 + L_2]. \end{aligned} \quad (3.13)$$

**Remark 3.3.** If  $f$  does not depend on  $u'$ , let the conditions of Theorem 3.1 hold with (3.1) replaced by

$$\|f(t, x) - qx\| \leq 2\alpha \langle x, f(t, x) \rangle + M, \quad (t, x) \in ([0, T] \setminus \{t_1\}) \times \mathbb{R}^n \times \mathbb{R}^n. \quad (3.14)$$

Then the impulsive BVP (1.2) has at least one solution.

#### 4. An example

In this section, we consider an example to illustrate the effectiveness of our new theorems. For brevity, we restrict our attention to scalar-valued impulsive BVPs, although we note that it is not difficult to construct a vector-valued  $f$  such that the conditions of Theorems 3.1 and 3.2 are satisfied.

*Example 4.1.* Consider the scalar impulsive BVP given by

$$\begin{aligned} u''(t) &= (u(t) + u'(t))^3 + u(t) + (u'(t))^2 + u'(t) + t, \quad t \in [0,1] \setminus \{t_1\}, \\ u(t_1^+) &= u(t_1^-) + \frac{u(t_1)}{5}, \quad u'(t_1^+) = u'(t_1^-) + \frac{u(t_1)}{7}, \\ u(0) &= u(1), \quad u'(0) = u'(1), \end{aligned} \tag{4.1}$$

we claim that the above impulsive BVP has at least one solution.

*Proof.* Let  $T = 1$ ,  $f(t, x, y) = (x + y)^5 + x + y^2 + y + t$ , and  $p = q = 1$ . Then  $r_1 = (\sqrt{5} + 1)/2$  and  $r_2 = (1 - \sqrt{5})/2$ . Obviously, (3.2) holds with  $\beta = 1/5$ ,  $\gamma = 1/7$ , and  $L_1 = L_2 = 0$ . We get  $1/H = 0.3534$  ( $H$  is as in (2.26)). Thus, (3.3) in Theorem 3.1 holds. Moreover, we see that

$$|f(t, x, y) - x - y| \leq |x + y|^5 + y^2 + 1, \quad \forall (t, x, y) \in [0, 1] \times \mathbb{R}^2, \tag{4.2}$$

and for  $\alpha = 1/2$  and  $M = 2$ ,

$$\begin{aligned} 2\alpha[(x + y)f(t, x, y) + y^2] + M &= (x + y)^6 + (x + y)^2 + (x + y)t + y^2 + 2 \\ &\geq (x + y)^6 + (x + y)^2 - |x + y| + y^2 + 2 \geq |x + y|^5 + y^2 + 1, \quad \forall (t, x, y) \in [0, 1] \times \mathbb{R}^2. \end{aligned} \tag{4.3}$$

Thus (3.1) holds. Therefore, by Theorem 3.1, BVP (4.1) has at least one solution.  $\square$

#### Acknowledgments

The authors are very grateful to the referees for careful reading of the original manuscript and for valuable suggestions on improving this paper. This project is supported by the Natural Science Foundation of Jiangsu Education Office (06KJB110010) and Jiangsu Planned Projects for Postdoctoral Research Funds.

#### References

- [1] M. Benchohra, J. Henderson, and S. Ntouyas, *Impulsive Differential Equations and Inclusions*, vol. 2 of *Contemporary Mathematics and Its Applications*, Hindawi, New York, NY, USA, 2006.
- [2] X. Liu, Ed., “Advances in Impulsive Differential Equations,” *Dynamics of Continuous, Discrete & Impulsive Systems. Series A. Mathematical Analysis*, vol. 9, no. 3, pp. 313–462, 2002.
- [3] Y. V. Rogovchenko, “Impulsive evolution systems: main results and new trends,” *Dynamics of Continuous, Discrete & Impulsive Systems. Series A. Mathematical Analysis*, vol. 3, no. 1, pp. 57–88, 1997.

## 12 Boundary Value Problems

- [4] A. M. Samoilenko and N. A. Perestyuk, *Impulsive Differential Equations*, vol. 14 of *World Scientific Series on Nonlinear Science. Series A: Monographs and Treatises*, World Scientific, River Edge, NJ, USA, 1995.
- [5] S. T. Zavalishchin and A. N. Sesekin, *Dynamic Impulse Systems. Theory and Applications*, vol. 394 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997.
- [6] M. Choisy, J. F. Guégan, and P. Rohani, “Dynamics of infectious diseases and pulse vaccination: teasing apart the embedded resonance effects,” *Physica D: Nonlinear Phenomena*, vol. 22, no. 1, pp. 26–35, 2006.
- [7] A. d’Onofrio, “On pulse vaccination strategy in the SIR epidemic model with vertical transmission,” *Applied Mathematics Letters*, vol. 18, no. 7, pp. 729–732, 2005.
- [8] S. Gao, L. Chen, J. J. Nieto, and A. Torres, “Analysis of a delayed epidemic model with pulse vaccination and saturation incidence,” *Vaccine*, vol. 24, no. 35–36, pp. 6037–6045, 2006.
- [9] Z. He and X. Zhang, “Monotone iterative technique for first order impulsive difference equations with periodic boundary conditions,” *Applied Mathematics and Computation*, vol. 156, no. 3, pp. 605–620, 2004.
- [10] W.-T. Li and H.-F. Huo, “Global attractivity of positive periodic solutions for an impulsive delay periodic model of respiratory dynamics,” *Journal of Computational and Applied Mathematics*, vol. 174, no. 2, pp. 227–238, 2005.
- [11] S. Tang and L. Chen, “Density-dependent birth rate, birth pulses and their population dynamic consequences,” *Journal of Mathematical Biology*, vol. 44, no. 2, pp. 185–199, 2002.
- [12] W. Wang, H. Wang, and Z. Li, “The dynamic complexity of a three-species Beddington-type food chain with impulsive control strategy,” *Chaos, Solitons & Fractals*, vol. 32, no. 5, pp. 1772–1785, 2007.
- [13] J. Yan, A. Zhao, and J. J. Nieto, “Existence and global attractivity of positive periodic solution of periodic single-species impulsive Lotka–Volterra systems,” *Mathematical and Computer Modelling*, vol. 40, no. 5–6, pp. 509–518, 2004.
- [14] W. Zhang and M. Fan, “Periodicity in a generalized ecological competition system governed by impulsive differential equations with delays,” *Mathematical and Computer Modelling*, vol. 39, no. 4–5, pp. 479–493, 2004.
- [15] X. Zhang, Z. Shuai, and K. Wang, “Optimal impulsive harvesting policy for single population,” *Nonlinear Analysis: Real World Applications*, vol. 4, no. 4, pp. 639–651, 2003.
- [16] R. P. Agarwal and D. O’Regan, “Multiple nonnegative solutions for second order impulsive differential equations,” *Applied Mathematics and Computation*, vol. 114, no. 1, pp. 51–59, 2000.
- [17] L. Chen and J. Sun, “Nonlinear boundary value problem of first order impulsive functional differential equations,” *Journal of Mathematical Analysis and Applications*, vol. 318, no. 2, pp. 726–741, 2006.
- [18] W. Ding, M. Han, and J. Mi, “Periodic boundary value problem for the second-order impulsive functional differential equations,” *Computers & Mathematics with Applications*, vol. 50, no. 3–4, pp. 491–507, 2005.
- [19] J. J. Nieto and R. Rodríguez-López, “Periodic boundary value problem for non-Lipschitzian impulsive functional differential equations,” *Journal of Mathematical Analysis and Applications*, vol. 318, no. 2, pp. 593–610, 2006.
- [20] I. Rachůnková and M. Tvrdý, “Non-ordered lower and upper functions in second order impulsive periodic problems,” *Dynamics of Continuous, Discrete & Impulsive Systems. Series A. Mathematical Analysis*, vol. 12, no. 3–4, pp. 397–415, 2005.
- [21] J. Chen, C. C. Tisdell, and R. Yuan, “On the solvability of periodic boundary value problems with impulse,” *Journal of Mathematical Analysis and Applications*, vol. 331, no. 2, pp. 902–912, 2007.

- [22] J. Li, J. J. Nieto, and J. Shen, “Impulsive periodic boundary value problems of first-order differential equations,” *Journal of Mathematical Analysis and Applications*, vol. 325, no. 1, pp. 226–236, 2007.
- [23] J. J. Nieto, “Periodic boundary value problems for first-order impulsive ordinary differential equations,” *Nonlinear Analysis*, vol. 51, no. 7, pp. 1223–1232, 2002.
- [24] C. Bai, “Existence of solutions for second order nonlinear functional differential equations with periodic boundary value conditions,” *International Journal of Pure and Applied Mathematics*, vol. 16, no. 4, pp. 451–462, 2004.
- [25] M. Rudd and C. C. Tisdell, “On the solvability of two-point, second-order boundary value problems,” *Applied Mathematics Letters*, vol. 20, no. 7, pp. 824–828, 2007.
- [26] Y. Dong, “Sublinear impulse effects and solvability of boundary value problems for differential equations with impulses,” *Journal of Mathematical Analysis and Applications*, vol. 264, no. 1, pp. 32–48, 2001.
- [27] Y. Liu, “Further results on periodic boundary value problems for nonlinear first order impulsive functional differential equations,” *Journal of Mathematical Analysis and Applications*, vol. 327, no. 1, pp. 435–452, 2007.
- [28] D. Qian and X. Li, “Periodic solutions for ordinary differential equations with sublinear impulsive effects,” *Journal of Mathematical Analysis and Applications*, vol. 303, no. 1, pp. 288–303, 2005.
- [29] N. G. Lloyd, *Degree Theory*, Cambridge Tracts in Mathematics, no. 73, Cambridge University Press, Cambridge, UK, 1978.
- [30] V. Lakshmikantham, D. D. Bainov, and P. S. Simeonov, *Theory of Impulsive Differential Equations*, vol. 6 of *Series in Modern Applied Mathematics*, World Scientific, Teaneck, NJ, USA, 1989.
- [31] J. J. Nieto, “Basic theory for nonresonance impulsive periodic problems of first order,” *Journal of Mathematical Analysis and Applications*, vol. 205, no. 2, pp. 423–433, 1997.

Chuanzhi Bai: Department of Mathematics, Huaiyin Teachers College, Huaian, Jiangsu 223300, China

Email address: czbai8@sohu.com

Dandan Yang: Department of Mathematics, Yangzhou University, Yangzhou 225002, China

Email address: yangdandan2600@sina.com

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)