

GENERALIZED QUASILINEARIZATION METHOD AND HIGHER ORDER OF CONVERGENCE FOR SECOND-ORDER BOUNDARY VALUE PROBLEMS

TANYA G. MELTON AND A. S. VATSALA

Received 24 March 2005; Revised 13 September 2005; Accepted 19 September 2005

The method of generalized quasilinearization for second-order boundary value problems has been extended when the forcing function is the sum of 2-hyperconvex and 2-hyperconcave functions. We develop two sequences under suitable conditions which converge to the unique solution of the boundary value problem. Furthermore, the convergence is of order 3. Finally, we provide numerical examples to show the application of the generalized quasilinearization method developed here for second-order boundary value problems.

Copyright © 2006 T. G. Melton and A. S. Vatsala. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The method of quasilinearization [1, 2] combined with the technique of upper and lower solutions is an effective and fruitful technique for solving a wide variety of nonlinear problems. It has been referred to as a generalized quasilinearization method. See [9] for details. The method is extremely useful in scientific computations due to its accelerated rate of convergence as in [10, 11].

In [4, 13], the authors have obtained a higher order of convergence (an order more than 2) for initial value problems. They have considered situations when the forcing function is either hyperconvex or hyperconcave. In [12], we have obtained the results of higher order of convergence for first order initial value problems when the forcing function is the sum of hyperconvex and hyperconcave functions with natural and coupled lower and upper solutions. In this paper we extend the result to the second-order boundary value problems when the forcing function is a sum of 2-hyperconvex and 2-hyperconcave functions. We have proved the existence of the unique solution of the nonlinear problem using natural upper and lower solutions. We demonstrate the iterates converge cubically to the unique solution of the nonlinear problem. We merely state the result related to coupled

2 GQ method for second-order BV problem

lower and upper solutions without proof due to monotony. Finally, we present two numerical applications of our theoretical results developed in our main result. We note that the monotone iterates may not converge linearly or quadratically in general. See [4, 8] for examples. However in our result we have provided sufficient conditions for cubic convergence. For real world applications see [5].

For this purpose, consider the following second-order boundary value problem (BVP for short):

$$-u'' = f(t, u) + g(t, u), \quad Bu(\mu) = b_\mu, \quad \mu = 0, 1, \quad t \in J \equiv [0, 1], \quad (1.1)$$

where $Bu(\mu) = \tau_\mu u(\mu) + (-1)^{\mu+1} \gamma_\mu u'(\mu) = b_\mu$, $\tau_0, \tau_1 \geq 0$, $\tau_0 + \tau_1 > 0$, $\gamma_0, \gamma_1 > 0$, $b_\mu \in R$ and $f, g \in C[J \times R, R]$.

Here we provide the definition of natural lower and upper solutions of (1.1). One can define coupled lower and upper solutions of the other types in the same manner. See for [14, 15] details.

Definition 1.1. The functions $\alpha_0, \beta_0 \in C^2[J, R]$ are said to be natural lower and upper solutions if

$$\begin{aligned} -\alpha_0'' &\leq f(t, \alpha_0) + g(t, \alpha_0), & B\alpha_0(\mu) &\leq b_\mu \quad \text{on } J, \\ -\beta_0'' &\geq f(t, \beta_0) + g(t, \beta_0), & B\beta_0(\mu) &\geq b_\mu \quad \text{on } J. \end{aligned} \quad (1.2)$$

In order to facilitate later explanations, we will need the following definition.

Definition 1.2. A function $h : A \rightarrow B$, $A, B \subset R$ is called m -hyperconvex, $m \geq 0$, if $h \in C^{m+1}[A, B]$ and $d^{m+1}h/du^{m+1} \geq 0$ for $u \in A$; h is called m -hyperconcave if the inequality is reversed.

In this paper, we use the maximum norm of u over J , that is,

$$\|u\| = \max_{t \in J} |u|. \quad (1.3)$$

Also throughout this paper we use the notation

$$\frac{\partial^k f(t, u)}{\partial u^k} = f^{(k)}(t, u) \quad (1.4)$$

for any function $f(t, u)$ and for $k = 0, 1, 2$.

In view of natural upper and lower solutions of (1.1), we will develop results when f is 2-hyperconvex and g is 2-hyperconcave. Furthermore, we show that these iterates converge uniformly and monotonically to the unique solution of (1.1), and the convergence is of order 3.

2. Preliminaries

In this section, we recall some well known theorems and corollaries which we need in our main results relative to the BVP

$$-u'' = f(t, u, u'), \quad Bu(\mu) = b_\mu, \quad \mu = 0, 1, \quad t \in J \equiv [0, 1], \quad (2.1)$$

where $Bu(\mu) = \tau_\mu u(\mu) + (-1)^{\mu+1} \nu_\mu u'(\mu) = b_\mu$, $\tau_0, \tau_1 \geq 0$, $\tau_0 + \tau_1 > 0$, $\nu_0, \nu_1 > 0$, $b_\mu \in R$ and $f \in C[J \times R \times R, R]$. For details see [3, 6, 7].

THEOREM 2.1. *Assume that*

- (i) $\alpha_0, \beta_0 \in C^2[J, R]$ *are lower and upper solutions of (2.1).*
- (ii) $f_u, f_{u'}$ *exist, continuous, $f_u < 0$ and $f_u \neq 0$ on $\Omega = [(t, u, \bar{u}) : t \in [0, 1], \beta_0 \leq u \leq \alpha_0]$ and $\bar{u} = \alpha'_0(t) = \beta'_0(t)$.*

Then we have $\alpha_0(t) \leq \beta_0(t)$ on J .

Next we present a special case of the above theorem which is known as the maximum principle, when u' term is missing.

COROLLARY 2.2. *Let $q, r \in C[I, R]$ with $r(t) \geq 0$ on J . Suppose further that $p \in C^2[I, R]$ and*

$$-p'' \leq -rp, \quad Bp(\mu) \leq 0. \quad (2.2)$$

Then $p(t) \leq 0$ on J . If the inequalities are reversed, then $p(t) \geq 0$ on J .

The next corollary is a special case of [9, Theorem 3.1.3].

COROLLARY 2.3. *Assume that α_0, β_0 are lower and upper solutions of (1.1) respectively such that $\alpha_0(t) \leq \beta_0(t)$ on J . Then there exists a solution u for the BVP (1.1) such that $\alpha_0(t) \leq u(t) \leq \beta_0(t)$ on J .*

3. Main results

In this section, we consider the BVP

$$-u'' = f(t, u) + g(t, u), \quad Bu(\mu) = b_\mu, \quad \mu = 0, 1, \quad t \in J \equiv [0, 1], \quad (3.1)$$

where $Bu(\mu) = \tau_\mu u(\mu) + (-1)^{\mu+1} \nu_\mu u'(\mu) = b_\mu$, $\tau_0, \tau_1 \geq 0$, $\tau_0 + \tau_1 > 0$, $\nu_0, \nu_1 > 0$, $b_\mu \in R$, $f, g \in C[\Omega, R]$, $\Omega = [(t, u) : \alpha_0(t) \leq u(t) \leq \beta_0(t), t \in J]$, and $\alpha_0, \beta_0 \in C^2[J, R]$ with $\alpha_0(t) \leq \beta_0(t)$ on J .

Here, we state the inequalities satisfied by $f(t, u)$ and $g(t, u)$ when $f(t, u)$ is 2-hyperconvex in u and $g(t, u)$ is 2-hyperconcave in u . We need these inequalities for our first main result.

Suppose that $f(t, u)$ is 2-hyperconvex in u , then we have the following inequalities,

$$f(t, \eta) \geq \sum_{i=0}^2 \frac{f^{(i)}(t, \xi)(\eta - \xi)^i}{i!}, \quad \eta \geq \xi, \quad (3.2)$$

$$f(t, \eta) \leq \sum_{i=0}^2 \frac{f^{(i)}(t, \xi)(\eta - \xi)^i}{i!}, \quad \eta \leq \xi. \quad (3.3)$$

4 GQ method for second-order BV problem

Similarly, when $g(t, u)$ is 2-hyperconcave in u , we have the following inequalities:

$$g(t, \eta) \geq \sum_{i=0}^1 \frac{g^{(i)}(t, \xi)(\eta - \xi)^i}{i!} + \frac{g^{(2)}(t, \eta)(\eta - \xi)^2}{(2)!}, \quad \eta \geq \xi, \quad (3.4)$$

$$g(t, \eta) \leq \sum_{i=0}^1 \frac{g^{(i)}(t, \xi)(\eta - \xi)^i}{i!} + \frac{g^{(2)}(t, \eta)(\eta - \xi)^2}{(2)!}, \quad \eta \leq \xi. \quad (3.5)$$

Based on these inequalities, relative to the natural upper and lower solutions, we develop two monotone sequences which converge uniformly and monotonically to the unique solution of (3.1) and the order of convergence is 3.

THEOREM 3.1. *Assume that*

- (i) $\alpha_0, \beta_0 \in C^2[J, R]$ are lower and upper solutions with $\alpha_0(t) \leq \beta_0(t)$ on J .
- (ii) $f, g \in C^3[\Omega, R]$ such that $f(t, u)$ is 2-hyperconvex in u on J [i.e., $f^{(3)}(t, u) \geq 0$ for $(t, u) \in \Omega$], $g(t, u)$ is 2-hyperconcave in u on J [i.e., $g^{(3)}(t, u) \leq 0$ for $(t, u) \in \Omega$], $f(t, u)$ is nondecreasing, $g(t, u)$ is nonincreasing and $f_u + g_u < 0$ on Ω .

Then there exist monotone sequences $\{\alpha_n(t)\}$ and $\{\beta_n(t)\}$, $n \geq 0$ which converge uniformly and monotonically to the unique solution of (3.1) and the convergence is of order 3.

Proof. The assumptions $f^{(3)}(t, u) \geq 0$, $g^{(3)}(t, u) \leq 0$ yield the inequalities (3.2), (3.3), (3.4), and (3.5) whenever $\alpha_0 \leq \eta$, $\xi \leq \beta_0$. Let us first consider the following BVPs:

$$\begin{aligned} -w'' &= \tilde{F}(t, \alpha, \beta; w) \\ &= \sum_{i=0}^2 \frac{f^{(i)}(t, \alpha)(w - \alpha)^i}{i!} + \sum_{i=0}^1 \frac{g^{(i)}(t, \alpha)(w - \alpha)^i}{i!} + \frac{g^{(2)}(t, \beta)(w - \alpha)^2}{2!}, \\ Bw(\mu) &= b_\mu \quad \text{on } J; \end{aligned} \quad (3.6)$$

$$\begin{aligned} -v'' &= \tilde{G}(t, \alpha, \beta; v) \\ &= \sum_{i=0}^2 \frac{f^{(i)}(t, \beta)(v - \beta)^i}{i!} + \sum_{i=0}^1 \frac{g^{(i)}(t, \beta)(v - \beta)^i}{i!} + \frac{g^{(2)}(t, \alpha)(v - \beta)^2}{2!}, \\ Bv(\mu) &= b_\mu \quad \text{on } J. \end{aligned} \quad (3.7)$$

We develop the sequences $\{\alpha_n(t)\}$ and $\{\beta_n(t)\}$ using the above BVPs (3.6) and (3.7) respectively. Initially, we prove (α_0, β_0) are lower and upper solutions of (3.6) and (3.7) respectively. To begin, we will consider natural lower and upper solutions of the equation (3.1):

$$\begin{aligned} -\alpha_0'' &\leq f(t, \alpha_0) + g(t, \alpha_0), & B\alpha_0(\mu) &\leq b_\mu, \\ -\beta_0'' &\geq f(t, \beta_0) + g(t, \beta_0), & B\beta_0(\mu) &\geq b_\mu, \end{aligned} \quad (3.8)$$

where $\alpha_0(t) \leq \beta_0(t)$. The inequalities (3.2) and (3.4), and (3.8) imply

$$\begin{aligned}
 -\alpha_0'' &\leq f(t, \alpha_0) + g(t, \alpha_0) \\
 &= \tilde{F}(t, \alpha_0, \beta_0; \alpha_0), \quad B\alpha_0(\mu) \leq b_\mu, \\
 -\beta_0'' &\geq f(t, \beta_0) + g(t, \beta_0) \\
 &\geq \sum_{i=0}^2 \frac{f^{(i)}(t, \alpha_0)(\beta_0 - \alpha_0)^i}{i!} + \sum_{i=0}^1 \frac{g^{(i)}(t, \alpha_0)(\beta_0 - \alpha_0)^i}{i!} + \frac{g^{(2)}(t, \beta_0)(\beta_0 - \alpha_0)^2}{2!} \\
 &= \tilde{F}(t, \alpha_0, \beta_0; \beta_0), \quad B\beta_0(\mu) \geq b_\mu.
 \end{aligned} \tag{3.9}$$

We can apply Corollary 2.3 together with (3.9) conclude that there exists a solution $\alpha_1(t)$ of (3.6) with $\alpha = \alpha_0$ and $\beta = \beta_0$ such that $\alpha_0 \leq \alpha_1 \leq \beta_0$ on J .

Using the inequalities (3.3), (3.5), and (3.8) on the same lines, we can get

$$-\beta_0'' \geq f(t, \beta_0) + g(t, \beta_0) = \tilde{G}(t, \alpha_0, \beta_0; \beta_0), \quad B\beta_0(\mu) \geq b_\mu, \tag{3.10}$$

$$\begin{aligned}
 -\alpha_0'' &\leq f(t, \alpha_0) + g(t, \alpha_0) \leq \sum_{i=0}^2 \frac{f^{(i)}(t, \beta_0)(\alpha_0 - \beta_0)^i}{i!} \\
 &\quad + \sum_{i=0}^1 \frac{g^{(i)}(t, \beta_0)(\alpha_0 - \beta_0)^i}{i!} + \frac{g^{(2)}(t, \alpha_0)(\alpha_0 - \beta_0)^2}{2!} \\
 &= \tilde{G}(t, \alpha_0, \beta_0; \alpha_0), \quad B\alpha_0(\mu) \leq b_\mu.
 \end{aligned} \tag{3.11}$$

Hence α_0, β_0 are lower and upper solutions of (3.7) with $\alpha_0 \leq \beta_0$. Applying Corollary 2.3, we obtain that there exists a solution $\beta_1(t)$ of (3.7) with $\alpha = \alpha_0$ and $\beta = \beta_0$ such that $\alpha_0 \leq \beta_1 \leq \beta_0$ on J .

Now we will prove that α_1 is a unique solution of (3.6). For this purpose we need to prove that $\partial \tilde{F}(t, \alpha_0, \beta_0; \alpha_1) / \partial \alpha_1 < 0$. Since $f(t, u)$ is 2-hyperconvex in u and $g(t, u)$ is 2-hyperconcave in u on J with $f_u + g_u < 0$ on Ω , we get

$$\begin{aligned}
 \frac{\partial \tilde{F}(t, \alpha_0, \beta_0; \alpha_1)}{\partial \alpha_1} &= f^{(1)}(t, \alpha_1) + g^{(1)}(t, \alpha_1) - \frac{f^{(3)}(t, \xi_1)(\alpha_1 - \alpha_0)^2}{(2)!} \\
 &\quad + g^{(3)}(t, \eta_1)(\alpha_1 - \alpha_0)(\beta_0 - \xi_2) \\
 &\leq f^{(1)}(t, \alpha_1) + g^{(1)}(t, \alpha_1) < 0,
 \end{aligned} \tag{3.12}$$

where $\alpha_0 \leq \xi_1, \xi_2 \leq \alpha_1$ and $\xi_2 \leq \eta_1 \leq \beta_0$. Hence by the special case of Theorem 2.1 with u' -term missing, we can conclude that α_1 is the unique solution of (3.6). Similarly we can prove that β_1 is the unique solution of (3.7).

6 GQ method for second-order BV problem

Using the nonincreasing property of $g^{(2)}(t, u)$, (3.2), (3.3), (3.4), (3.5) with $\alpha_0 \leq \alpha_1 \leq \beta_0$, $\alpha_0 \leq \beta_1 \leq \beta_0$ we have

$$\begin{aligned}
 -\alpha_1'' &= \tilde{F}(t, \alpha_0, \beta_0; \alpha_1) \\
 &= \sum_{i=0}^2 \frac{f^{(i)}(t, \alpha_0)(\alpha_1 - \alpha_0)^i}{i!} \\
 &\quad + \sum_{i=0}^1 \frac{g^{(i)}(t, \alpha_0)(\alpha_1 - \alpha_0)^i}{i!} + \frac{g^{(2)}(t, \beta_0)(\alpha_1 - \alpha_0)^2}{2!} \\
 &\leq f(t, \alpha_1) + g(t, \alpha_1), \quad B\alpha_1(\mu) \leq b_\mu;
 \end{aligned} \tag{3.13}$$

$$\begin{aligned}
 -\beta_1'' &= \tilde{G}(t, \alpha_0, \beta_0; \beta_1) \\
 &= \sum_{i=0}^2 \frac{f^{(i)}(t, \beta_0)(\beta_1 - \beta_0)^i}{i!} \\
 &\quad + \sum_{i=0}^1 \frac{g^{(i)}(t, \beta_0)(\beta_1 - \beta_0)^i}{i!} + \frac{g^{(2)}(t, \alpha_0)(\beta_1 - \beta_0)^2}{2!} \\
 &\geq f(t, \beta_1) + g(t, \beta_1), \quad B\beta_1(\mu) \geq b_\mu.
 \end{aligned} \tag{3.14}$$

Since α_1, β_1 are lower and upper solutions of (3.1), we can apply the special case of Theorem 2.1 to obtain $\alpha_1 \leq \beta_1$ on J . Thus we have $\alpha_0 \leq \alpha_1 \leq \beta_1 \leq \beta_0$ on J .

Assume now that α_n and β_n are solutions of BVPs (3.6) and (3.7), respectively, with $\alpha = \alpha_{n-1}$ and $\beta = \beta_{n-1}$ such that $\alpha_{n-1} \leq \alpha_n \leq \beta_n \leq \beta_{n-1}$ on J and

$$\begin{aligned}
 -\alpha_n'' &\leq f(t, \alpha_n) + g(t, \alpha_n), \quad B\alpha_n(\mu) \leq b_\mu, \\
 -\beta_n'' &\geq f(t, \beta_n) + g(t, \beta_n), \quad B\beta_n(\mu) \geq b_\mu,
 \end{aligned} \tag{3.15}$$

Certainly this is true for $n = 1$.

We need to show that $\alpha_n \leq \alpha_{n+1} \leq \beta_{n+1} \leq \beta_n$ on J , where α_{n+1} and β_{n+1} are solutions of BVPs (3.6) and (3.7), respectively, with $\alpha = \alpha_n$ and $\beta = \beta_n$.

The inequalities (3.2) and (3.4), and (3.15) imply

$$\begin{aligned}
 -\alpha_n'' &\leq f(t, \alpha_n) + g(t, \alpha_n) \\
 &= \tilde{F}(t, \alpha_n, \beta_n; \alpha_n), \quad B\alpha_n(\mu) \leq b_\mu, \\
 -\beta_n'' &\geq f(t, \beta_n) + g(t, \beta_n) \\
 &\geq \sum_{i=0}^2 \frac{f^{(i)}(t, \alpha_n)(\beta_n - \alpha_n)^i}{i!} \\
 &\quad + \sum_{i=0}^1 \frac{g^{(i)}(t, \alpha_n)(\beta_n - \alpha_n)^i}{i!} + \frac{g^{(2)}(t, \beta_n)(\beta_n - \alpha_n)^2}{2!} \\
 &= \tilde{F}(t, \alpha_n, \beta_n; \beta_n), \quad B\beta_n(\mu) \geq b_\mu.
 \end{aligned} \tag{3.16}$$

This proves that α_n, β_n are lower and upper solutions of (3.6) with $\alpha = \alpha_n$ and $\beta = \beta_n$. Hence using (3.16) and Corollary 2.3 we can conclude that there exists a solution $\alpha_{n+1}(t)$ of (3.6) with $\alpha = \alpha_n$ and $\beta = \beta_n$ such that $\alpha_n \leq \alpha_{n+1} \leq \beta_n$ on J .

The inequalities (3.3) and (3.5), and (3.15) imply

$$-\beta_n'' \geq f(t, \beta_n) + g(t, \beta_n) = \tilde{G}(t, \alpha_n, \beta_n; \beta_n), \quad B\beta_n(\mu) \geq b_\mu, \quad (3.17)$$

$$\begin{aligned} -\alpha_n'' &\leq f(t, \alpha_n) + g(t, \alpha_n) \\ &\leq \sum_{i=0}^2 \frac{f^{(i)}(t, \beta_n)(\alpha_n - \beta_n)^i}{i!} \\ &\quad + \sum_{i=0}^1 \frac{g^{(i)}(t, \beta_n)(\alpha_n - \beta_n)^i}{i!} + \frac{g^{(2)}(t, \alpha_n)(\alpha_n - \beta_n)^2}{2!} \\ &= \tilde{G}(t, \alpha_n, \beta_n; \alpha_n), \quad B\alpha_n(\mu) \leq b_\mu. \end{aligned} \quad (3.18)$$

Hence α_n, β_n are lower and upper solutions of (3.7) with $\alpha = \alpha_n$ and $\beta = \beta_n$. Applying Corollary 2.3 we can show that there exists a solution $\beta_{n+1}(t)$ of (3.7) with $\alpha = \alpha_n$ and $\beta = \beta_n$ such that $\alpha_n \leq \beta_{n+1} \leq \beta_n$ on J . In view of assumptions on f and g , $\alpha_{n+1}, \beta_{n+1}$ are unique by the special case of Theorem 2.1.

Furthermore, by (3.2), (3.3), (3.4), (3.5) with $\alpha_n \leq \alpha_{n+1} \leq \beta_n$, $\alpha_n \leq \beta_{n+1} \leq \beta_n$, and $g^{(2)}(t, u)$ nonincreasing u , we get

$$\begin{aligned} -\alpha_{n+1}'' &= \tilde{F}(t, \alpha_n, \beta_n; \alpha_{n+1}) \\ &= \sum_{i=0}^2 \frac{f^{(i)}(t, \alpha_n)(\alpha_{n+1} - \alpha_n)^i}{i!} \\ &\quad + \sum_{i=0}^1 \frac{g^{(i)}(t, \alpha_n)(\alpha_{n+1} - \alpha_n)^i}{i!} + \frac{g^{(2)}(t, \beta_n)(\alpha_{n+1} - \alpha_n)^2}{2!} \\ &\leq f(t, \alpha_{n+1}) + g(t, \alpha_{n+1}), \quad B\alpha_{n+1}(\mu) \leq b_\mu; \\ -\beta_{n+1}'' &= \tilde{G}(t, \alpha_n, \beta_n; \beta_{n+1}) \\ &= \sum_{i=0}^2 \frac{f^{(i)}(t, \beta_n)(\beta_{n+1} - \beta_n)^i}{i!} \\ &\quad + \sum_{i=0}^1 \frac{g^{(i)}(t, \beta_n)(\beta_{n+1} - \beta_n)^i}{i!} + \frac{g^{(2)}(t, \alpha_n)(\beta_{n+1} - \beta_n)^2}{2!} \\ &\geq f(t, \beta_{n+1}) + g(t, \beta_{n+1}), \quad B\beta_{n+1}(\mu) \geq b_\mu. \end{aligned} \quad (3.19)$$

Since $\alpha_{n+1}, \beta_{n+1}$ are lower and upper solutions of (3.1) we can apply the special case of Theorem 2.1 and get $\alpha_{n+1} \leq \beta_{n+1}$ on J . This proves $\alpha_n \leq \alpha_{n+1} \leq \beta_{n+1} \leq \beta_n$ on J . Hence by induction, we have

$$\alpha_0 \leq \alpha_1 \leq \cdots \leq \alpha_n \leq \beta_n \leq \cdots \leq \beta_1 \leq \beta_0. \quad (3.20)$$

8 GQ method for second-order BV problem

By the fact that α_n, β_n are lower and upper solutions of (3.1) with $\alpha_n \leq \beta_n$ and Corollary 2.3 we can conclude that there exists a solution $u(t)$ of (3.1) such that $\alpha_n \leq u \leq \beta_n$ on J . From this we can obtain that

$$\alpha_0 \leq \alpha_1 \leq \cdots \leq \alpha_n \leq u \leq \beta_n \leq \cdots \leq \beta_1 \leq \beta_0. \quad (3.21)$$

Using Green's function, we can write $\alpha_n(t)$ and $\beta_n(t)$ as follows:

$$\begin{aligned} \alpha_n(t) &= \int_0^1 K(t,s) \tilde{F}(s, \alpha_{n-1}(s), \beta_{n-1}(s); \alpha_n(s)) ds, \\ \beta_n(t) &= \int_0^1 K(t,s) \tilde{G}(s, \alpha_{n-1}(s), \beta_{n-1}(s); \beta_n(s)) ds. \end{aligned} \quad (3.22)$$

Here $K(t,s)$ is the Green's function given by

$$K(t,s) = \begin{cases} \frac{1}{c} x(s)y(t), & 0 \leq s \leq t \leq 1, \\ \frac{1}{c} x(t)y(s), & 0 \leq t \leq s \leq 1, \end{cases} \quad (3.23)$$

where $x(t) = (\tau_0/\nu_0)t + 1$, $y(t) = (\tau_1/\nu_1)(1-t) + 1$ are two linearly independent solutions of $-u'' = 0$ and $c = x(t)y'(t) - x'(t)y(t)$. We can prove that the sequences $\{\alpha_n(t)\}$ and $\{\beta_n(t)\}$ are equicontinuous and uniformly bounded. Now applying Ascoli-Arzelà's theorem, we can show that there exist subsequences $\{\alpha_{n,j}(t)\}$, $\{\beta_{n,j}(t)\}$ such that $\alpha_{n,j}(t) \rightarrow \rho(t)$ and $\beta_{n,j}(t) \rightarrow r(t)$ with $\rho(t) \leq u \leq r(t)$ on J . Since the sequences $\{\alpha_n(t)\}$, $\{\beta_n(t)\}$ are monotone, we have $\alpha_n(t) \rightarrow \rho(t)$ and $\beta_n(t) \rightarrow r(t)$. Taking the limit as $n \rightarrow \infty$, we get

$$\lim_{n \rightarrow \infty} \alpha_n(t) = \rho(t) \leq u \leq r(t) = \lim_{n \rightarrow \infty} \beta_n(t). \quad (3.24)$$

Next we show that $\rho(t) \geq r(t)$. From BVPs (3.6) and (3.7) we get

$$\begin{aligned} -\rho''(t) &= f(t, \rho) + g(t, \rho), & B\rho(\mu) &= b(\mu), \\ -r''(t) &= f(t, r) + g(t, r), & Br(\mu) &= b(\mu). \end{aligned} \quad (3.25)$$

Set $p(t) = r - \rho$ and note that $Bp(\mu) = 0$. We have

$$\begin{aligned} -p'' &= -r''(t) - (-\rho''(t)) = f(t, r) + g(t, r) - f(t, \rho) - g(t, \rho) \\ &= f_u(t, \xi)(r - \rho) + g_u(t, \eta)(r - \rho) = (f_u(t, \xi) + g_u(t, \eta))p, \end{aligned} \quad (3.26)$$

where ξ, η are between ρ and r . This implies that $-p'' \leq -kp$, where $f_u + g_u \leq -k < 0$. Now applying Corollary 2.2 we get $p \leq 0$ or $r(t) \leq \rho(t)$ on J . This proves $r(t) = \rho(t) = u(t)$. Hence $\{\alpha_n(t)\}$ and $\{\beta_n(t)\}$ converge uniformly and monotonically to the unique solution of (3.1).

Let us consider the order of convergence of $\{\alpha_n(t)\}$ and $\{\beta_n(t)\}$ to the unique solution $u(t)$ of (3.1). To do this, set

$$\begin{aligned} p_n(t) &= u(t) - \alpha_n(t) \geq 0, \\ q_n(t) &= \beta_n(t) - u(t) \geq 0, \end{aligned} \quad (3.27)$$

for $t \in J$ with $Bp_n(\mu) = Bq_n(\mu) = 0$.

Therefore we can write

$$p_{n+1} = \int_0^1 K(t, s) [f(s, u) + g(s, u) - \tilde{F}(s, \alpha_n, \beta_n; \alpha_{n+1})] ds, \quad (3.28)$$

where $K(t, s)$ is the Green's function given by (3.23).

Now using the Taylor series expansion with Lagrange remainder, and the mean value theorem together with (ii) of the hypothesis, we obtain

$$\begin{aligned} 0 &\leq p_{n+1} \\ &= \int_0^1 K(t, s) \left\{ f(s, u) + g(s, u) \right. \\ &\quad - \left[\sum_{i=0}^2 \frac{f^{(i)}(s, \alpha_n) (\alpha_{n+1} - \alpha_n)^i}{i!} \right. \\ &\quad \left. \left. + \sum_{i=0}^1 \frac{g^{(i)}(s, \alpha_n) (\alpha_{n+1} - \alpha_n)^i}{i!} + \frac{g^{(2)}(s, \beta_n) (\alpha_{n+1} - \alpha_n)^2}{2!} \right] \right\} ds \\ &= \int_0^1 K(t, s) \left\{ f(s, u) + g(s, u) \right. \\ &\quad - \left[f(s, \alpha_{n+1}) - \frac{f^{(3)}(s, \xi_1) (\alpha_{n+1} - \alpha_n)^3}{(3)!} + g(s, \alpha_{n+1}) \right. \\ &\quad \left. \left. - \frac{g^{(2)}(s, \xi_2) (\alpha_{n+1} - \alpha_n)^2}{2!} + \frac{g^{(2)}(s, \beta_n) (\alpha_{n+1} - \alpha_n)^2}{2!} \right] \right\} ds \\ &\leq \int_0^1 K(t, s) \left[f_u(s, \eta_1) (u - \alpha_{n+1}) + g_u(s, \eta_2) (u - \alpha_{n+1}) \right. \\ &\quad \left. + \frac{f^{(3)}(s, \xi_1) (u - \alpha_n)^3}{(3)!} - \frac{g^{(3)}(s, \eta_3) (\beta_n - \xi_2) (u - \alpha_n)^2}{2} \right] ds \\ &= \int_0^1 K(t, s) \left\{ [f_u(s, \eta_1) + g_u(s, \eta_2)] p_{n+1} \right. \\ &\quad \left. + \frac{f^{(3)}(s, \xi_1) p_n^3}{(3)!} - \frac{g^{(3)}(s, \eta_2) p_n^2 (q_n + p_n)}{2} \right\} ds, \end{aligned} \quad (3.29)$$

10 GQ method for second-order BV problem

where $\alpha_n \leq \xi_1$, $\xi_2 \leq \alpha_{n+1} \leq \eta_1$, $\eta_2 \leq u$, and $\xi_2 \leq \eta_3 \leq \beta_n$. Let $|K(t,s)| \leq A_1$, $|f_u(t,u) + g_u(t,v)| \leq A_2$, $|f^{(3)}(t,u)/3!| \leq A_3$, and $|g^{(3)}(t,u)/2| \leq A_4$. Then we have

$$\|p_{n+1}\| \leq k_1 \|p_n\|^3 + k_2 \|p_n\|^2 (\|q_n\| + \|p_n\|), \quad (3.30)$$

where $k_1 = A_1 A_3 / (1 - A_1 A_2)$ and $k_2 = A_1 A_4 / (1 - A_1 A_2)$.

Similarly, we can write

$$q_{n+1} = \int_0^1 K(t,s) [\tilde{G}(s, \alpha_n, \beta_n; \beta_{n+1}) - f(s, u) - g(s, u)] ds, \quad (3.31)$$

where $K(t,s)$ is the Green's function given by (3.23).

Using the Taylor series expansion with Lagrange remainder, and the mean value theorem together with (ii), we can show

$$\|q_{n+1}\| \leq k_1 \|q_n\|^3 + k_2 \|q_n\|^2 (\|q_n\| + \|p_n\|), \quad (3.32)$$

where $k_1 = A_1 A_3 / (1 - A_1 A_2)$ and $k_2 = A_1 A_4 / (1 - A_1 A_2)$.

Hence combining (3.30) and (3.32) we obtain

$$\begin{aligned} & \max_{t \in J} |u(t) - \alpha_{n+1}(t)| + \max_{t \in J} |\beta_{n+1}(t) - u(t)| \\ & \leq C \left[\max_{t \in J} |u(t) - \alpha_n(t)| + \max_{t \in J} |\beta_n(t) - u(t)| \right]^3, \end{aligned} \quad (3.33)$$

where C is an appropriate positive constant.

This completes the proof. $\square$

We note that the unique solution we have obtained is the unique solution of (3.1) in the sector determined by the lower and upper solutions.

Next we merely state a result without proof using coupled lower and upper solutions of (3.1). However, in order to show the existence of the unique solution of the iterates, we use the existence result [7, Theorem 2.4.1]. for systems and a special case of the comparison theorem of [7].

THEOREM 3.2. *Assume that*

- (i) $\alpha_0, \beta_0 \in C^2[J, R]$ are coupled lower and upper solutions of (3.1) with $\alpha_0(t) \leq \beta_0(t)$ on J such that

$$\begin{aligned} -\alpha_0'' &\leq f(t, \beta_0) + g(t, \alpha_0), & B\alpha_0(\mu) &\leq b_\mu & \text{on } J, \\ -\beta_0'' &\geq f(t, \alpha_0) + g(t, \beta_0), & B\beta_0(\mu) &\geq b_\mu & \text{on } J; \end{aligned} \quad (3.34)$$

- (ii) $f, g \in C^3[\Omega, \mathbb{R}]$ such that $f(t, u)$ is 2-hyperconvex in u on J [i.e., $f^{(3)}(t, u) \geq 0$ for $(t, u) \in \Omega$], $g(t, u)$ is 2-hyperconcave in u on J [i.e., $g^{(3)}(t, u) \leq 0$ for $(t, u) \in \Omega$], $f(t, u), g(t, u)$ are nonincreasing with $f_u - g_u > 0$ and

$$f_u(t, u) \leq -\max_{\Omega} [f^{(3)}(t, u)] (\beta_0 - \alpha_0)^2 \leq 0 \quad \text{on } \Omega. \quad (3.35)$$

Then there exist monotone sequences $\{\alpha_n(t)\}$ and $\{\beta_n(t)\}$, $n \geq 0$ such that

$$\begin{aligned} -\alpha_n'' &= \sum_{i=0}^1 \frac{f^{(i)}(t, \beta_{n-1}) (\beta_n - \beta_{n-1})^i}{i!} + \frac{f^{(2)}(t, \alpha_{n-1}) (\beta_n - \beta_{n-1})^2}{(2)!} \\ &\quad + \sum_{i=0}^1 \frac{g^{(i)}(t, \alpha_{n-1}) (\alpha_n - \alpha_{n-1})^i}{i!} + \frac{g^{(2)}(t, \beta_{n-1}) (\alpha_n - \alpha_{n-1})^2}{(2)!}, \\ B\alpha_n(\mu) &= b_\mu \quad \text{on } J; \\ -\beta_n'' &= \sum_{i=0}^1 \frac{f^{(i)}(t, \alpha_{n-1}) (\alpha_n - \alpha_{n-1})^i}{i!} + \frac{f^{(2)}(t, \beta_{n-1}) (\alpha_n - \alpha_{n-1})^2}{(2)!} \\ &\quad + \sum_{i=0}^1 \frac{g^{(i)}(t, \beta_{n-1}) (\beta_n - \beta_{n-1})^i}{i!} + \frac{g^{(2)}(t, \alpha_{n-1}) (\beta_n - \beta_{n-1})^2}{(2)!}, \\ B\beta_n(\mu) &= b_\mu \quad \text{on } J, \end{aligned} \quad (3.36)$$

which converge uniformly and monotonically to the unique solution of (3.1) and the convergence is of order 3.

Remark 3.3. Similar results can be obtained for the other two coupled upper and lower solutions of (3.1) and the numerical applications of these results can be demonstrated.

4. Numerical results

Next we will provide an example which satisfies all the hypotheses of Theorem 3.1 which demonstrates the application of Theorem 3.1.

Example 4.1. Let us consider the following BVP:

$$\begin{aligned} -u'' &= u^3 - 2u^4 - 0.1u + 0.4, \\ u(0) &= 0, \quad u(1) = 1. \end{aligned} \quad (4.1)$$

It is easy to check that $\alpha_0(t) \equiv 0$ and $\beta_0(t) \equiv 1$ are natural lower and upper solutions for (4.1), respectively. Let $H(t, u)$ denote the right-hand side of (4.1) and split it into nonincreasing and nondecreasing functions as $H(t, u) = f(t, u) + g(t, u)$ where

$$\begin{aligned} f(t, u) &= u^3, \\ g(t, u) &= -2u^4 - 0.1u + 0.4. \end{aligned} \quad (4.2)$$

Table 4.1. Table of three α, β -iterates of (4.1).

t	$\alpha_1(t)$	$\alpha_2(t)$	$\alpha_3(t)$	$\beta_3(t)$	$\beta_2(t)$	$\beta_1(t)$
0.1	0.071613	0.105795	0.114155	0.115155	0.121882	0.211998
0.2	0.139816	0.207722	0.224408	0.226396	0.239650	0.372556
0.3	0.206071	0.305997	0.330820	0.333780	0.352693	0.497196
0.4	0.272568	0.401158	0.433435	0.437334	0.460118	0.596776
0.5	0.342305	0.494293	0.532364	0.537104	0.561279	0.679076
0.6	0.419460	0.587139	0.627930	0.633249	0.656144	0.749881
0.7	0.510237	0.681965	0.720845	0.726212	0.745445	0.813727
0.8	0.624444	0.781200	0.812382	0.816952	0.830723	0.874419
0.9	0.778369	0.886852	0.904514	0.907236	0.914404	0.935401




Figure 4.1

It is easy to show that

$$\begin{aligned} f_{uuu} &= 6 > 0, \\ g_{uuu} &= -48u \leq 0 \end{aligned} \quad (4.3)$$

for $0 \leq u \leq 1$. Hence f is a 2-hyperconvex function and g is a 2-hyperconcave function. Now we need to check the following conditions in order to use Theorem 3.1:

$$\begin{aligned} f_u(t, u) &= 3u^2 \geq 0, \\ g_u(t, u) &= -8u^3 - 0.1 \leq 0, \\ f_u(t, u) + g_u(t, u) &= 3u^2 - 8u^3 - 0.1 < 0, \end{aligned} \quad (4.4)$$

whenever $0 \leq u \leq 1$. Hence we can apply the iterates of Theorem 3.1. Using the nonlinear finite-difference methods for BVPs and Mathematica we can find the α, β -iterates as given in Table 4.1.

The α -iterates (with broken line) and the β -iterates (with unbroken line) can be seen on Figure 4.1.

Given the specific finite difference scheme, we can apply it to obtain lower and upper solutions. Then, we can make the difference between upper and lower solutions arbitrarily small. The obtained numerical solution however, will be close to the actual solution of the nonlinear problem (4.1) only within the truncation error of the finite difference scheme chosen.

Now we will provide a numerical example to show the usefulness of Theorem 3.2.

Example 4.2. Let us discuss the following second-order BVP:

$$-u'' = 3 \cos u - 27e^{u/3} + 25.5, \quad u(0.1) = 0.1, \quad u(0.5) = 0.5. \quad (4.5)$$

Denote the right-hand side of (4.5) by $H(t, u)$. We can split the forcing function into two functions as $H(t, u) = f(t, u) + g(t, u)$ where

$$\begin{aligned} f(t, u) &= 3 \cos u, \\ g(t, u) &= -27e^{u/3} + 25.5. \end{aligned} \quad (4.6)$$

If we choose $\alpha_0(t) \equiv 0.1$, $\beta_0(t) \equiv 0.5$, and $0.1 \leq t \leq 0.5$ we get

$$\begin{aligned} 0 &\leq 3 \cos 0.5 - 27e^{0.1/3} + 25.5 = 0.21758, \\ 0 &\geq 3 \cos 0.1 - 27e^{0.5/3} + 25.5 = -3.41172, \\ 0.1 &\leq 0.1, \quad 0.5 \geq 0.5. \end{aligned} \quad (4.7)$$

Thus $\alpha_0(t) \equiv 0.1$ and $\beta_0(t) \equiv 0.5$ are coupled lower and upper solutions for (4.5) of the type defined in Theorem 3.2.

Next we can show that

$$\begin{aligned} f_{uuu} &= 3 \sin u > 0, \\ g_{uuu} &= -e^{u/3} < 0 \end{aligned} \quad (4.8)$$

for $0.1 \leq u \leq 0.5$. Hence f is 2-hyperconvex function and g is 2-hyperconcave function. Now we need to check the following conditions in order to apply Theorem 3.2:

$$\begin{aligned} f_u(t, u) &= -3 \sin u < 0, \\ g_u(t, u) &= -9e^{u/3} < 0, \\ f_u(t, u) - g_u(t, u) &= e^u + u^2 > 0, \\ -3 \sin 0.1 &\leq -3 \sin 0.5(0.5 - 0.1)^2 \leq 0, \end{aligned} \quad (4.9)$$

whenever $0.1 \leq u \leq 0.5$. Hence all the hypotheses of Theorem 3.2 are satisfied and we can apply the given iterates. Now using the nonlinear finite-difference methods for BVPs and Mathematica we can derive the α, β -iterates in Table 4.2.

The graph on Figure 4.2 shows α -iterates (with broken line) and the β -iterates (with unbroken line).

Table 4.2. Table of three α, β -iterates of (4.5).

t	$\alpha_1(t)$	$\alpha_2(t)$	$\alpha_3(t)$	$\beta_3(t)$	$\beta_2(t)$	$\beta_1(t)$
0.10	0.100000	0.100000	0.100000	0.100000	0.100000	0.100000
0.15	0.141017	0.141551	0.141573	0.141646	0.142080	0.145352
0.20	0.182354	0.182909	0.182942	0.183077	0.183878	0.189605
0.25	0.225015	0.225334	0.225365	0.225541	0.226590	0.233909
0.30	0.270074	0.270116	0.270138	0.270329	0.271473	0.279436
0.35	0.318505	0.318582	0.318595	0.318773	0.319842	0.327418
0.40	0.372012	0.372103	0.372110	0.372247	0.373077	0.379177
0.45	0.432077	0.432091	0.432095	0.432169	0.432624	0.436164
0.50	0.500000	0.500000	0.500000	0.500000	0.500000	0.500000




Figure 4.2

Remark 4.3. Note that the interval in the above example is due to the fact that f and g satisfies the hypothesis of Theorem 3.2 on the specific interval chosen.

5. Conclusion

We have used iterates of nonlinearity of order 2 when the forcing function is the sum of 2-hyperconvex and 2-hyperconcave. We develop two sequences depending on the type of the lower and upper solutions, which converge rapidly (order 3) to the unique solution of (3.1). We demonstrate the application of the results with numerical applications.

References

- [1] R. E. Bellman, *Methods of Nonlinear Analysis. Vol. 1*, Mathematics in Science and Engineering, vol. 61-I, Academic Press, New York, 1970.
- [2] R. E. Bellman and R. E. Kalaba, *Quasilinearization and Nonlinear Boundary-Value Problems*, Modern Analytic and Computational Methods in Science and Mathematics, vol. 3, American Elsevier, New York, 1965.
- [3] S. R. Bernfeld and V. Lakshmikantham, *An Introduction to Nonlinear Boundary Value Problems*, Mathematics in Science and Engineering, vol. 109, Academic Press, New York, 1974.

- [4] A. Cabada and J. J. Nieto, *Rapid convergence of the iterative technique for first order initial value problems*, Applied Mathematics and Computation **87** (1997), no. 2-3, 217–226.
- [5] ———, *Quasilinearization and rate of convergence for higher-order nonlinear periodic boundary-value problems*, Journal of Optimization Theory and Applications **108** (2001), no. 1, 97–107.
- [6] S. Heikkilä and V. Lakshmikantham, *Monotone Iterative Techniques for Discontinuous Nonlinear Differential Equations*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 181, Marcel Dekker, New York, 1994.
- [7] G. S. Ladde, V. Lakshmikantham, and A. S. Vatsala, *Monotone Iterative Techniques for Nonlinear Differential Equations*, Monographs, Advanced Texts and Surveys in Pure and Applied Mathematics, vol. 27, Pitman, Massachusetts, 1985.
- [8] V. Lakshmikantham and J. J. Nieto, *Generalized quasilinearization iterative method for initial value problems*, Nonlinear Studies **2** (1995), 1–9.
- [9] V. Lakshmikantham and A. S. Vatsala, *Generalized Quasilinearization for Nonlinear Problems*, Mathematics and Its Applications, vol. 440, Kluwer Academic, Dordrecht, 1998.
- [10] V. B. Mandelzweig, *Quasilinearization method and its verification on exactly solvable models in quantum mechanics*, Journal of Mathematical Physics **40** (1999), no. 12, 6266–6291.
- [11] V. B. Mandelzweig and F. Tabakin, *Quasilinearization approach to nonlinear problems in physics with application to nonlinear ODEs*, Computer Physics Communications **141** (2001), no. 2, 268–281.
- [12] T. Melton and A. S. Vatsala, *Generalized quasilinearization and higher order of convergence for first order initial value problems*, to appear in Dynamic Systems & Applications.
- [13] R. N. Mohapatra, K. Vajravelu, and Y. Yin, *Extension of the method of quasilinearization and rapid convergence*, Journal of Optimization Theory and Applications **96** (1998), no. 3, 667–682.
- [14] M. Sokol and A. S. Vatsala, *A unified exhaustive study of monotone iterative method for initial value problems*, Nonlinear Studies **8** (2001), no. 4, 429–438.
- [15] I. H. West and A. S. Vatsala, *Generalized monotone iterative method for initial value problems*, Applied Mathematics Letters **17** (2004), no. 11, 1231–1237.

Tanya G. Melton: Department of Mathematics, University of Louisiana at Lafayette, Lafayette,
LA 70504-1010, USA

E-mail address: tmelton@lsua.edu

A. S. Vatsala: Department of Mathematics, University of Louisiana at Lafayette, Lafayette,
LA 70504-1010, USA

E-mail address: vatsala@louisiana.edu

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br