

A TRANSMISSION PROBLEM FOR BEAMS ON NONLINEAR SUPPORTS

TO FU MA AND HIGIDIO PORTILLO OQUENDO

Received 20 October 2005; Revised 10 April 2006; Accepted 12 April 2006

A transmission problem involving two Euler-Bernoulli equations modeling the vibrations of a composite beam is studied. Assuming that the beam is clamped at one extremity, and resting on an elastic bearing at the other extremity, the existence of a unique global solution and decay rates of the energy are obtained by adding just one damping device at the end containing the bearing mechanism.

Copyright © 2006 T. F. Ma and H. Portillo Oquendo. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In this paper we consider the existence of a global solution and decay rates of the energy for a transmission problem involving two Euler-Bernoulli equations with nonlinear boundary conditions. More precisely, we are concerned with the system of equations

$$\rho_1 u_{tt} + \beta_1 u_{xxxx} = 0 \quad \text{in }]0, L_0[\times \mathbb{R}^+, \quad (1.1)$$

$$\rho_2 v_{tt} + \beta_2 v_{xxxx} = 0 \quad \text{in }]L_0, L[\times \mathbb{R}^+, \quad (1.2)$$

coupled by the “transmission” conditions

$$\begin{aligned} u(L_0, t) - v(L_0, t) &= 0, & u_x(L_0, t) - v_x(L_0, t) &= 0, \\ \beta_1 u_{xx}(L_0, t) - \beta_2 v_{xx}(L_0, t) &= 0, & \beta_1 u_{xxx}(L_0, t) - \beta_2 v_{xxx}(L_0, t) &= 0. \end{aligned} \quad (1.3)$$

To the system we add the nonlinear boundary conditions

$$u(0, t) = 0, \quad u_x(0, t) = 0, \quad (1.4)$$

$$v_{xx}(L, t) = 0, \quad \beta_2 v_{xxx}(L, t) = f(v(L, t)) + g(v_t(L, t)), \quad (1.5)$$

2 A transmission problem for beams on nonlinear supports




Figure 1.1. A composite beam on an elastic bearing.

and the initial data

$$\begin{aligned} u(x,0) &= u^0(x), & u_t(x,0) &= u^1(x) & \text{in } [0, L_0], \\ v(x,0) &= v^0(x), & v_t(x,0) &= v^1(x) & \text{in } [L_0, L]. \end{aligned} \quad (1.6)$$

The system (1.1)–(1.6) models the transverse vibrations of a composite beam of length L , constituted by two types of materials of different mass densities $\rho_1, \rho_2 > 0$ and flexural rigidities $\beta_1, \beta_2 > 0$. Because of the boundary condition (1.4), the beam is clamped at the left end $x = 0$. On the other extremity, the condition (1.5) implies that the bending moment is zero and that the shear force is equal to $f(v(L, t)) + g(v_t(L, t))$. This means that, at the end $x = L$, the beam is resting on a kind of bearing, described by the function f , and subjected to a frictional dissipation described by the function g (see Figure 1.1).

We notice that stabilization of transmission problems has been considered by some authors. In the beginning, Lions [8] studied the exact controllability of the transmission problem for the wave equation. Later, Liu and Williams [10] studied the boundary stabilization of transmission problems for linear systems of wave equations. In the case of beam equations, which involve fourth-order derivatives, there are more possibilities in the problem modeling and boundary conditions. For instance, Muñoz Rivera and Portillo Oquendo [14] studied a transmission problem for viscoelastic beams, by exploiting the dissipations due to the memory effects of the material. On the other hand, there are a few results on fourth-order equations with nonlinear boundary conditions involving third-order derivatives. That class of problems models elastic beams on elastic bearings, and one of the first results, with nonlinearities, was given by Feireisl [2], who studied the periodic solutions for a superlinear problem. Some related stationary problems were considered by Grossinho and Ma [3] and Ma [11]. We refer the reader to [1, 4–7, 12–16] for other interesting related works.

Our objective is to show that under suitable assumptions, the sole dissipation $g(v_t)$, acting on the boundary point $x = L$, will be sufficient to stabilize the whole system. The dissipation effect on the boundary $x = L$ will be transmitted to (1.1) through (1.2). The proof of the boundary stabilization is based on the arguments from Lagnese [6] and Lagnese and Leugering [7].

The paper is organized as follows. In Section 2 we define some notations and establish the global existence and uniqueness results (see Theorem 2.2). Weak solutions are also considered (see Theorem 2.6). In Section 3 we prove the decay of the energy of the system,

which is defined by

$$\begin{aligned} E(t) = E(t, u, v) &= \frac{1}{2} \int_0^{L_0} \{ \rho_1 |u_t| + \beta_1 |u_{xx}| \} dx \\ &+ \frac{1}{2} \int_{L_0}^L \{ \rho_2 |v_t| + \beta_2 |v_{xx}| \} dx + \hat{f}(v(L, t)), \end{aligned} \quad (1.7)$$

where $\hat{f}(w) = \int_0^w f(s) ds$ (see Theorem 3.1).

2. Global existence

In our study we assume that f is a C^1 function satisfying the sign condition

$$f(w)w \geq 0, \quad \forall w \in \mathbb{R}, \quad (2.1)$$

and that g is a C^1 for which there exists a constant $c_0 > 0$ such that

$$g(0) = 0, \quad (g(r) - g(s))(r - s) \geq c_0 |r - s|^2, \quad \forall r, s \in \mathbb{R}. \quad (2.2)$$

In particular it follows that $g(w)w \geq c_0 w^2$ for all $w \in \mathbb{R}$. In order to deal with the transmission conditions (1.3) and the boundary condition (1.4), we define the Sobolev space

$$X = \{(\varphi, \psi) \in \mathbb{H}^2 \mid (\varphi, \psi) \text{ satisfies (2.4)}\}, \quad (2.3)$$

where

$$\varphi(0) = \varphi_x(0) = \varphi(L_0) - \psi(L_0) = \varphi_x(L_0) - \psi_x(L_0) = 0, \quad (2.4)$$

$$\mathbb{H}^k = H^k(0, L_0) \times H^k(L_0, L). \quad (2.5)$$

We also write $\mathbb{L}^2 = L^2(0, L_0) \times L^2(L_0, L)$. Our study is based on the space

$$V = \{(\varphi, \psi) \in (H^2(0, L_0) \times H^3(L_0, L)) \cap X \mid \psi_{xx}(L) = 0\}, \quad (2.6)$$

so that the first part of condition (1.5) is also recovered. As a simple consequence of the trace theorem and (2.4) one has the following useful boundary estimate.

LEMMA 2.1. *Given $(u, v) \in C^1([0, T], X)$, there exists a constant $C > 0$ such that*

$$\begin{aligned} |v(L, t)| &\leq C \{ \|u_{xx}\|_2 + \|v_{xx}\|_2 \}, \quad \forall t \in [0, T], \\ |v_t(L, t)| &\leq C \{ \|u_{xxt}\|_2 + \|v_{xxt}\|_2 \}, \quad \forall t \in [0, T], \end{aligned} \quad (2.7)$$

where $\|\cdot\|_2$ denotes either $L^2(0, L_0)$ or $L^2(L_0, L)$ norms.

Now we prove the existence of global regular solutions.

4 A transmission problem for beams on nonlinear supports

THEOREM 2.2. *Assume that conditions (2.1)-(2.2) hold. Then for any initial data $(u^0, v^0) \in \mathbb{H}^4 \cap V$ and $(u^1, v^1) \in V$, satisfying the compatibility condition,*

$$\begin{aligned} v_{xxx}^0(L) - f(v^0(L)) - g(v^1(L)) &= 0, \\ \beta_1 u_{xx}^0(L_0, t) - \beta_2 v_{xx}^0(L_0, t) &= 0, \\ \beta_1 u_{xxx}^0(L_0, t) - \beta_2 v_{xxx}^0(L_0, t) &= 0, \end{aligned} \quad (2.8)$$

problem (1.1)–(1.6) has a unique strong solution (u, v) such that

$$(u, v) \in L^\infty(\mathbb{R}^+; \mathbb{H}^4), \quad (u_t, v_t) \in L^\infty(\mathbb{R}^+; X), \quad (u_{tt}, v_{tt}) \in L^\infty(\mathbb{R}^+; \mathbb{L}^2). \quad (2.9)$$

The proof of Theorem 2.2 is given in several steps, by using the Galerkin method.

Approximate problem. Let $\{(\varphi^n, \psi^n)\}_{n \in \mathbb{N}}$ be a Galerkin basis of V , which for convenience is chosen to satisfy

$$\{(u^0, v^0), (u^1, v^1)\} \subset V_2, \quad (2.10)$$

where

$$V_m = \text{span}\{(\varphi^1, \psi^1), \dots, (\varphi^m, \psi^m)\}. \quad (2.11)$$

Then the corresponding approximate variational problem to problem (1.1)–(1.6) reads as follows: find $(u^m(t), v^m(t)) \in V_m$ of the form

$$(u^m(t), v^m(t)) = \sum_{j=1}^m h_j^m(t) (\varphi^j, \psi^j) \quad (2.12)$$

such that

$$\begin{aligned} \int_0^{L_0} \{\rho_1 u_{tt}^m \varphi^j + \beta_1 u_{xxx}^m \varphi_{xx}^j\} dx + \int_{L_0}^L \{\rho_2 v_{tt}^m \psi^j + \beta_2 v_{xxx}^m \psi_{xx}^j\} dx \\ + \{\beta_2 f(v^m(L, t)) + g(v_t^m(L, t))\} \psi^j(L) = 0, \end{aligned} \quad (2.13)$$

$$(u^m(0), v^m(0)) = (u^0, v^0), \quad (u_t^m(0), v_t^m(0)) = (u^1, v^1). \quad (2.14)$$

As a matter of fact, (2.13) is an m -dimensional system of ODEs in $h_j^m(t)$ and has a local solution $(u^m(t), v^m(t))$ in an interval $[0, t_m]$. In the following, we derive uniform estimates, so that local solutions can be extended to the interval $[0, T]$ for any $T > 0$. Note that initial conditions in (2.14) are well defined because of (2.10).

Estimate 2.3. Replacing φ^i by u_t^m and ψ^i by v_t^m in (2.13), one concludes that

$$\frac{d}{dt} E(t, u^m, v^m) = -\beta_2 g(v_t^m(L, t)) v_t^m(L, t). \quad (2.15)$$

Then from condition (2.2) we see that $E(t, u^m, v^m)$ is decreasing and therefore there exists $M_1 > 0$ such that

$$\|u_t^m(t)\|_2^2 + \|v_t^m(t)\|_2^2 + \|u_{xx}^m(t)\|_2^2 + \|v_{xx}^m(t)\|_2^2 \leq M_1 \quad (2.16)$$

for all $m \in \mathbb{N}$, $t > 0$, where M_1 depends on $E(0, u^0, v^0)$.

Estimate 2.4. Let us obtain an estimate for $u_{tt}^m(0)$ and $v_{tt}^m(0)$ in L^2 norms. Replacing φ^i by $u_{tt}^m(0)$ and ψ^i by $v_{tt}^m(0)$ in (2.13), one concludes from the compatibility condition (2.8) that for some constant $C > 0$,

$$\|u_{tt}^m(0)\|_2^2 + \|v_{tt}^m(0)\|_2^2 \leq C \left(\|u_{xxx}^0\|_2^2 + \|v_{xxx}^0\|_2^2 \right). \quad (2.17)$$

Therefore, there exists $M = M(u^0, v^0) > 0$ such that

$$\|u_{tt}^m(0)\|_2^2 + \|v_{tt}^m(0)\|_2^2 \leq M \quad (2.18)$$

for all $m \in \mathbb{N}$.

Estimate 2.5. Here we use a finite-difference argument as in [12]. Let us fix $t, \xi > 0$ such that $\xi < T - t$, and take the difference of (2.13) with $t = t + \xi$ and $t = t$. Then replacing φ^j by $u_t^m(t + \xi) - u_t^m(t)$ and ψ^j by $v_t^m(t + \xi) - v_t^m(t)$, and putting

$$\begin{aligned} \hat{P}_m(t, \xi) = & \rho_1 \|u_t^m(t + \xi) - u_t^m(t)\|_2^2 + \rho_2 \|v_t^m(t + \xi) - v_t^m(t)\|_2^2 \\ & + \beta_1 \|u_{xx}^m(t + \xi) - u_{xx}^m(t)\|_2^2 + \beta_2 \|v_{xx}^m(t + \xi) - v_{xx}^m(t)\|_2^2, \end{aligned} \quad (2.19)$$

one infers that

$$\frac{1}{2} \frac{d}{dt} \hat{P}_m(t, \xi) \leq A + B, \quad (2.20)$$

where

$$\begin{aligned} A = & -(g(v^m(L, t + \xi)) - g(v^m(L, t))) (v_t^m(L, t + \xi) - v_t^m(L, t)), \\ B = & -\beta_2 (f(v^m(L, t + \xi)) - f(v^m(L, t))) (v_t^m(L, t + \xi) - v_t^m(L, t)). \end{aligned} \quad (2.21)$$

Taking $0 < \varepsilon < c_0$, and using the mean value theorem and Lemma 2.1, there exists $C_\varepsilon > 0$ such that

$$\begin{aligned} B \leq & C_\varepsilon \left(\|u_{xx}^m(t + \xi) - u_{xx}^m(t)\|_2^2 + \|v_{xx}^m(t + \xi) - v_{xx}^m(t)\|_2^2 \right) \\ & + \varepsilon \|v_t^m(L, t + \xi) - v_t^m(L, t)\|^2. \end{aligned} \quad (2.22)$$

Then from condition (2.2), we conclude that for a constant $C > 0$,

$$\frac{1}{2} \frac{d}{dt} \hat{P}_m(t, \xi) = C \hat{P}_m(t, \xi), \quad (2.23)$$

6 A transmission problem for beams on nonlinear supports

and therefore $\hat{P}_m(t, \xi) \leq \hat{P}_m(0, \xi)e^{2CT}$. So, dividing the inequality by ξ^2 and making $\xi \rightarrow 0$, we see that

$$\begin{aligned} & \rho_1 \|u_{tt}^m(t)\|_2^2 + \rho_2 \|v_{tt}^m(t)\|_2^2 + \beta_1 \|u_{xxt}^m(t)\|_2^2 + \beta_2 \|v_{xxt}^m(t)\|_2^2 \\ & \leq \left(\rho_1 \|u_{tt}^m(0)\|_2^2 + \rho_2 \|v_{tt}^m(0)\|_2^2 + \beta_1 \|u_{xx}^1\|_2^2 + \beta_2 \|v_{xx}^1\|_2^2 \right) e^{CT}. \end{aligned} \quad (2.24)$$

Hence there exists $M_2 > 0$ such that

$$\|u_{tt}^m(t)\|_2^2 + \|v_{tt}^m(t)\|_2^2 + \|u_{xxt}^m(t)\|_2^2 + \|v_{xxt}^m(t)\|_2^2 \leq M_2 \quad (2.25)$$

for all $m \in \mathbb{N}$ and $t \in [0, T]$.

Existence result. From Estimates 2.3 and 2.5, we can apply Aubin-Lions compactness theorem to pass to the limit the approximate problem. Then the proof of the existence result is complete.

Uniqueness. Let (u_1, v_1) and (u_2, v_2) be two solutions of problem (1.1)–(1.6). Writing $U = u_1 - u_2$ and $V = v_1 - v_2$, we see that (U, V) satisfies

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left\{ \rho_1 \|U_t(t)\|_2^2 + \rho_2 \|V_t(t)\|_2^2 + \beta_1 \|U_{xx}(t)\|_2^2 + \beta_2 \|V_{xx}(t)\|_2^2 \right\} \\ & \leq -\beta_2 [f(v_1(L, t)) - f(v_2(L, t))] V_t(L, t) \\ & \quad - \beta_2 [g(v_{1t}(L, t)) - g(v_{2t}(L, t))] V_t(L, t). \end{aligned} \quad (2.26)$$

Then using (2.2) and Lemma 2.1, as in Estimate 2.5, we deduce the existence of $C > 0$ such that

$$\frac{d}{dt} P(t) \leq C \left(\|U_{xx}(t)\|_2^2 + \|V_{xx}(t)\|_2^2 \right), \quad t \in [0, T], \quad (2.27)$$

where now $P(t) = P(U, V, t)$. Since we have $P(0) = 0$, from Gronwall lemma we get $U = V = 0$.

Weak solutions. We say that a pair (u, v) is a weak solution of problem (1.1)–(1.6) if

$$(u, v) \in L^\infty(\mathbb{R}^+, X), \quad (u_t, v_t) \in L^\infty(\mathbb{R}^+, \mathbb{L}^2), \quad (u_{tt}, v_{tt}) \in L^\infty(\mathbb{R}^+, \mathbb{H}^{-2}) \quad (2.28)$$

satisfy the initial conditions (1.6), the compatibility conditions (2.8), and the variational identity

$$\begin{aligned} & \frac{d}{dt} \left\{ \int_0^{L_0} \rho_1 u_t \varphi dx + \int_{L_0}^L \rho_2 v_t \psi dx \right\} \\ & + \int_0^{L_0} \beta_1 u_{xx} \varphi_{xx} dx + \int_{L_0}^L \beta_2 v_{xx} \psi_{xx} dx + \{ \beta_2 f(v(L, t)) + g(v_t(L, t)) \} \psi(L) = 0 \end{aligned} \quad (2.29)$$

for all $(\varphi, \psi) \in X$. In order to study the existence of weak solutions let us denote by $\mathcal{C}$ the set of all acceptable initial data for the existence of strong solutions, that is,

$$\mathcal{C} := \{((u^0, v^0), (u^1, v^1)) \in (\mathbb{H}^4 \cap V) \times V \mid (2.8) \text{ holds}\}. \quad (2.30)$$

Then we have the following existence result for weak solutions.

THEOREM 2.6. *Assume that conditions (2.1)–(2.2) hold. Then for any initial data satisfying*

$$((u^0, v^0), (u^1, v^1)) \in \overline{\mathcal{C}}^{\mathbb{H}^2 \times \mathbb{L}^2}, \quad (2.31)$$

problem (1.1)–(1.6) has a unique weak solution.

This theorem is proved using density arguments, similar to those used by Cavalcanti et al. [1]. In fact, from the assumption on the initial data, there exists a sequence $((u_\nu^0, v_\nu^0), (u_\nu^1, v_\nu^1)) \in \mathcal{C}$ such that

$$(u_\nu^0, v_\nu^0) \longrightarrow (u^0, v^0) \quad \text{in } \mathbb{H}^2, \quad (u_\nu^1, v_\nu^1) \longrightarrow (u^1, v^1) \quad \text{in } \mathbb{L}^2. \quad (2.32)$$

Now, for each $\nu \in \mathbb{N}$, the initial conditions (u_ν^0, v_ν^0) and (u_ν^1, v_ν^1) give a unique regular solution (u_ν, v_ν) of problem (1.1)–(1.6). From the estimates used in the proof of Theorem 2.2 it can be shown that (u_ν, v_ν) converges to a weak solution (u, v) of (1.1)–(1.6). The uniqueness is then proved by means of the regularization techniques as by Lions and Visik (see e.g. [9]).

3. Decay of the energy

In this section we study decay rates for the first-order energy (1.7) associated to system (1.1)–(1.6). Here we assume that the bearing device has a superlinear behavior, characterized by the condition

$$\exists \rho \geq 2 \quad \text{such that } \rho \hat{f}(w) - f(w)w \leq 0, \quad \forall w \in \mathbb{R}, \quad (3.1)$$

and that the material of the beam occupying $[L_0, L]$ is more dense and stiff than that in $[0, L_0]$, that is,

$$\rho_1 \leq \rho_2, \quad \beta_1 \geq \beta_2. \quad (3.2)$$

Then the rate of decay will depend on the behavior of the nonlinear dissipation g in a neighborhood of the origin, which is related to the following assumption: there exist $c_1, c_2 > 0$ and $q \geq 1$ such that

$$c_1 \min\{|w|, |w|^q\} \leq |g(w)| \leq c_2 \max\{|w|, |w|^{1/q}\}. \quad (3.3)$$

Our main result is given by the following theorem.

THEOREM 3.1. *Suppose that*

$$(u^0, v^0) \in \mathbb{H}^2 \cap X, \quad (u^1, v^1) \in \mathbb{L}^2. \quad (3.4)$$

8 A transmission problem for beams on nonlinear supports

Suppose in addition that conditions (3.1)–(3.3) also hold. Then if (u, v) is the solution of problem (1.1)–(1.6), one has the following decay rates:

(1) if $q > 1$, then there exists a positive constant $C = C(E(0))$ such that

$$E(t) \leq C(1+t)^{-2/(q-1)}; \quad (3.5)$$

(2) if $q = 1$, then there exist positive constants C and μ such that

$$E(t) \leq CE(0)e^{-\mu t}. \quad (3.6)$$

We will prove this theorem for strong solutions. Our conclusion follows by a standard density argument.

In order, we establish some auxiliary results related to the multipliers method. Let us introduce the functional

$$R_1(t) := \int_0^{L_0} \rho_1 u_t x u_x dx + \int_{L_0}^L \rho_2 v_t x v_x dx. \quad (3.7)$$

In the following lemma we retrieve a part of the energy.

LEMMA 3.2. *There exists a positive constant $C_1 = C_1(E(0))$ such that*

$$\begin{aligned} \frac{d}{dt} R_1(t) &\leq \frac{\rho_2 L}{2} |v_t(L, t)| + C_1 \{f(v(L, t))v(L, t) + |g(v_t(L, t))|\} \\ &\quad - \frac{1}{2} \int_0^{L_0} \rho_1 |u_t| + \beta_1 |u_{xx}| - \frac{1}{2} \int_{L_0}^L \rho_2 |v_t| + \beta_2 |v_{xx}| dx \end{aligned} \quad (3.8)$$

for any strong solution of (1.1)–(1.6).

Proof. Multiplying (1.1) by xu_x , (1.2) by xv_x , integrating by parts, and using the boundary conditions (1.4)–(1.5) and (1.3), we arrive at the following identity:

$$\begin{aligned} \frac{d}{dt} R_1(t) &= \frac{L_0}{2} (\rho_1 - \rho_2) |u_t(L_0, t)|^2 + \frac{L_0}{2} \frac{\beta_1}{\beta_2} (\beta_2 - \beta_1) |u_{xx}(L_0, t)|^2 \\ &\quad + \frac{\rho_2 L}{2} |v_t(L, t)|^2 - L(f(v(L, t)) + g(v_t(L, t))) v_x(L, t) \\ &\quad - \frac{1}{2} \int_0^{L_0} \rho_1 |u_t|^2 + 3\beta_1 |u_{xx}|^2 dx - \frac{1}{2} \int_{L_0}^L \rho_2 |v_t|^2 + 3\beta_2 |v_{xx}|^2 dx. \end{aligned} \quad (3.9)$$

In view of the inequalities (3.2), the above equation reduces to

$$\begin{aligned} \frac{d}{dt} R_1(t) &\leq \frac{\rho_2 L}{2} |v_t(L, t)|^2 - \underbrace{L(f(v(L, t)) + g(v_t(L, t))) v_x(L, t)}_{:=I_1} \\ &\quad - \frac{1}{2} \int_0^{L_0} \rho_1 |u_t|^2 + 3\beta_1 |u_{xx}|^2 dx - \frac{1}{2} \int_{L_0}^L \rho_2 |v_t|^2 + 3\beta_2 |v_{xx}|^2 dx. \end{aligned} \quad (3.10)$$

Now we will estimate I_1 . Lemma 2.1 implies that $|v(L, t)| \leq CE^{1/2}(0)$ for some $C > 0$, thus, as $f \in C^1(\mathbb{R})$ we have that $|f(v(L, t))| \leq C|v(L, t)|$ for some other positive constant $C = C(E^{1/2}(0))$. Applying Young's inequality and taking into account the preceding estimates, we get for $\eta > 0$,

$$\begin{aligned} I_1 &\leq \eta |v_x(L, t)|^2 + C_\eta \left\{ |f(v(L, t))|^2 + |g(v_t(L, t))|^2 \right\} \\ &\leq \eta |v_x(L, t)|^2 + C_\eta \left\{ f(v(L, t))v(L, t) + |g(v_t(L, t))|^2 \right\}, \end{aligned} \quad (3.11)$$

from where by Lemma 2.1 follows that

$$\begin{aligned} I_1 &\leq \eta C \left\{ \int_0^{L_0} \beta_1 |u_{xx}|^2 dx + \int_{L_0}^L \beta_2 |v_{xx}|^2 dx \right\} \\ &\quad + C_\eta \left\{ f(v(L, t))v(L, t) + |g(v_t(L, t))|^2 \right\}. \end{aligned} \quad (3.12)$$

Substitution of this inequality into (3.10) and fixing $\eta > 0$ small our conclusion follows. $\square$

Our next step is to retrieve the remainder part of the energy. Let (φ, ψ) be the solution of the stationary problem

$$\beta_1 \varphi_{xxxx} = 0 \quad \text{on }]0, L_0[\times \mathbb{R}^+, \quad (3.13)$$

$$\beta_2 \psi_{xxxx} = 0 \quad \text{on }]L_0, L[\times \mathbb{R}^+, \quad (3.14)$$

satisfying the boundary conditions

$$\begin{aligned} \varphi(0, t) &= \varphi_x(0, t) = 0, \\ \psi_{xx}(L, t) &= 0, \quad \psi(L, t) = v(L, t), \\ \varphi(L_0, t) - \psi(L_0, t) &= 0, \\ \varphi_x(L_0, t) - \psi_x(L_0, t) &= 0, \\ \beta_1 \varphi_{xx}(L_0, t) - \beta_2 \psi_{xx}(L_0, t) &= 0, \\ \beta_1 \varphi_{xxx}(L_0, t) - \beta_2 \psi_{xxx}(L_0, t) &= 0, \end{aligned} \quad (3.15)$$

which depend clearly on $v(L, t)$. We consider the following functional:

$$R_2(t) := \int_0^{L_0} \rho_1 u_t \varphi dx + \int_{L_0}^L \rho_2 v_t \psi dx. \quad (3.16)$$

LEMMA 3.3. *Given $\epsilon > 0$, there exists a positive constant C_ϵ such that*

$$\begin{aligned} \frac{d}{dt} R_2(t) &\leq \epsilon \left\{ \int_0^{L_0} \rho_1 |u_t|^2 + \beta_1 |u_{xx}|^2 dx + \int_{L_0}^L \rho_2 |v_t|^2 + \beta_2 |v_{xx}|^2 dx \right\} \\ &\quad + C_\epsilon \left\{ |v_t(L, t)|^2 + |g(v_t(L, t))|^2 \right\} - \frac{1}{2} f(v(L, t))v(L, t) \end{aligned} \quad (3.17)$$

for any strong solution of (1.1)–(1.6).

Proof. Multiplying (1.1) by φ , (1.2) by ψ , integrating by parts and using boundary conditions (1.3)–(1.5) and (3.15), we have the following identity:

$$\begin{aligned} \frac{d}{dt}R_2(t) &= \int_0^{L_0} \rho_1 u_t \varphi_t dx + \int_{L_0}^L \rho_2 v_t \psi_t dx - \int_0^{L_0} \beta_1 u_{xx} \varphi_{xx} dx \\ &\quad - \int_{L_0}^L \beta_2 v_{xx} \psi_{xx} dx - (f(v(L, t)) + g(v_t(L, t)))v(L, t). \end{aligned} \quad (3.18)$$

On the other hand, multiplying (3.13) by $u - \varphi$, (3.14) by $v - \psi$, integrating by parts and using boundary conditions (1.3)–(1.5) and (3.15), we obtain

$$\int_0^{L_0} \beta_1 u_{xx} \varphi_{xx} dx + \int_{L_0}^L \beta_2 v_{xx} \psi_{xx} dx = \int_0^{L_0} \beta_1 |\varphi_{xx}|^2 dx + \int_{L_0}^L \beta_2 |\psi_{xx}|^2 dx. \quad (3.19)$$

Since the right-hand side of this equality is positive, by substitution of this into (3.18) we arrive at

$$\frac{d}{dt}R_2(t) \leq \int_0^{L_0} \rho_1 u_t \varphi_t dx + \int_{L_0}^L \rho_2 v_t \psi_t dx - f(v(L, t))v(L, t) - g(v_t(L, t))v(L, t). \quad (3.20)$$

Now, we will estimate the last term of the above inequality. Using Young's inequality and Lemma 2.1, we have for $\eta > 0$,

$$\begin{aligned} |g(v_t(L, t))v(L, t)| &\leq \eta |v(L, t)|^2 + C_\eta |g(v_t(L, t))|^2 \\ &\leq \eta C \left\{ \int_0^{L_0} \beta_1 |u_{xx}|^2 dx + \int_{L_0}^L \beta_2 |v_{xx}|^2 dx \right\} + C_\eta |g(v_t(L, t))|^2. \end{aligned} \quad (3.21)$$

On the other hand, from the elliptic regularity of the system (3.13)–(3.15) there exists a constant $C > 0$ such that

$$\int_0^{L_0} |\varphi|^2 dx + \int_{L_0}^L |\psi|^2 dx \leq C |v(L, t)|^2, \quad (3.22)$$

and since the system (3.13)–(3.15) is linear we also have

$$\int_0^{L_0} |\varphi_t|^2 dx + \int_{L_0}^L |\psi_t|^2 dx \leq C |v_t(L, t)|^2. \quad (3.23)$$

Applying Young's inequality to the two first terms of the right-hand side of (3.20) and using the above estimate we have for $\eta > 0$,

$$\begin{aligned} \int_0^{L_0} \rho_1 u_t \varphi_t dx &\leq \eta \int_0^{L_0} \rho_1 |u_t|^2 dx + C_\eta |v_t(L, t)|^2, \\ \int_{L_0}^L \rho_2 v_t \psi_t dx &\leq \eta \int_{L_0}^L \rho_2 |v_t|^2 dx + C_\eta |v_t(L, t)|^2. \end{aligned} \quad (3.24)$$

By substitution of the estimates (3.21)–(3.24) into (3.20) and taking $\epsilon = \max\{\eta, \eta C\}$ we arrive at the desired result. This completes the proof of the lemma. $\square$

Now, we will summarize the results of the previous lemmas. Let us consider the following functional:

$$R(t) := R_1(t) + 2(C_1 + 1)R_2(t), \quad (3.25)$$

where C_1 is the constant considered in Lemma 3.2.

LEMMA 3.4. *There exists a positive constant C such that*

$$\frac{d}{dt}R(t) \leq -\frac{1}{2}E(t) + C\{g(v_t(L, t))v_t(L, t) + [g(v_t(L, t))v_t(L, t)]^{2/(q+1)}\} \quad (3.26)$$

for any strong solution of (1.1)–(1.6).

Proof. First, let ϵ_0 be the solution of

$$2(C_1 + 1)\epsilon_0 = \frac{1}{4}. \quad (3.27)$$

Combining Lemmas 3.2 and 3.3 with $\epsilon = \epsilon_0$ and using the superlinearity of the function f (see (3.1)) we arrive at

$$\frac{d}{dt}R(t) \leq -\frac{1}{2}E(t) + C\{|v_t(L, t)|^2 + |g(v_t(L, t))|^2\}. \quad (3.28)$$

Now, we will estimate the second term of the right-hand side of (3.28). From the hypothesis (3.3) we have the following estimates:

$$\begin{aligned} |v_t(L, t)| \geq 1 & \quad \text{then } |v_t(L, t)|^2 + |g(v_t(L, t))|^2 \leq Cg(v_t(L, t))v_t(L, t), \\ |v_t(L, t)| \leq 1 & \quad \text{then } |v_t(L, t)|^2 + |g(v_t(L, t))|^2 \leq C[g(v_t(L, t))v_t(L, t)]^{2/(q+1)}. \end{aligned} \quad (3.29)$$

Therefore, for any value of $v_t(L, t)$, we conclude that

$$|v_t(L, t)|^2 + |g(v_t(L, t))|^2 \leq C\{g(v_t(L, t))v_t(L, t) + [g(v_t(L, t))v_t(L, t)]^{2/(q+1)}\}. \quad (3.30)$$

In view of (3.28) the proof is complete. $\square$

Proof of Theorem 3.1. Now Lemma 3.4 plays an essential role. To prove the polynomial decay of the energy, we assume that $q > 1$. Using Young's inequality is not difficult to show that there exists a positive constant C such that

$$|R(t)| \leq CE(t). \quad (3.31)$$

Let us denote $\sigma := (q - 1)/2$. Since

$$\frac{d}{dt}E(t) = -g(v_t(L, t))v_t(L, t), \quad (3.32)$$

we get from estimate (3.31) that

$$\begin{aligned} \frac{d}{dt}[E^\sigma R](t) &\leq \sigma R(t)E^{\sigma-1}(t)\frac{d}{dt}E(t) + E^\sigma(t)\frac{d}{dt}R(t) \\ &\leq CE^\sigma(t)g(v_t(L, t))v_t(L, t) + E^\sigma(t)\frac{d}{dt}R(t). \end{aligned} \quad (3.33)$$

Using Lemma 3.4 and estimate $E(t) \leq E(0)$, the above inequality can be written as

$$\begin{aligned} \frac{d}{dt}[E^\sigma R](t) &\leq -\frac{1}{2}E^{\sigma+1}(t) + CE^\sigma(0)g(v_t(L, t))v_t(L, t) \\ &\quad + CE^{(q+1)/2}(0)E^{(q-1)/(q+1)}(t)[g(v_t(L, t))v_t(L, t)]^{2/(q+1)}. \end{aligned} \quad (3.34)$$

Using Young's inequality, the last term of the above inequality can be estimated by

$$\begin{aligned} CE^{(q+1)/2}(0)E^{(q-1)/(q+1)}(t)[g(v_t(L, t))v_t(L, t)]^{2/(q+1)} \\ \leq \eta E^{\sigma+1}(t) + C_\eta E^{(q+1)^{2/4}}(0)g(v_t(L, t))v_t(L, t). \end{aligned} \quad (3.35)$$

Taking $\eta = 1/4$, inequality (3.34) becomes

$$\frac{d}{dt}[E^\sigma R](t) \leq -\frac{1}{4}E^{\sigma+1}(t) + Cg(v_t(L, t))v_t(L, t), \quad (3.36)$$

where C is a constant which depends continuously on $E(0)$. Now, let us define the Lyapunov functional

$$F(t) := NE(t) + [E^\sigma R](t). \quad (3.37)$$

Combining identity (3.32) with inequality (3.36) and taking N large, we get

$$\frac{d}{dt}F(t) \leq -\frac{1}{4}E^{\sigma+1}(t). \quad (3.38)$$

On the other hand, in view of (3.31) we have that for N large,

$$\frac{N}{2}E(t) \leq F(t) \leq 2NE(t). \quad (3.39)$$

These two last inequalities imply that

$$\frac{d}{dt}F(t) \leq -\alpha F^{\sigma+1}(t), \quad \alpha = (2N)^{-(\sigma+1)}, \quad (3.40)$$

from where follows that

$$F(t) \leq \frac{1}{(F^{-\sigma}(0) + \alpha\sigma t)^{1/\sigma}}. \quad (3.41)$$

Finally, the equivalence relation (3.39) implies the polynomial decay of the energy E . This proves the first part of Theorem 3.1.

It remains to prove the exponential decay of the energy. To this end, we assume that $q = 1$. From identity (3.32) and inequality (3.36) we have that the Lyapunov functional

$$F(t) := NE(t) + R(t) \quad (3.42)$$

satisfies

$$\frac{d}{dt}F(t) \leq -\frac{1}{2}E(t), \quad (3.43)$$

from where in view of (3.39) follows that for N large,

$$\frac{d}{dt}F(t) \leq -\frac{1}{4N}F(t) \implies F(t) \leq F(0)e^{-t/4N}. \quad (3.44)$$

Finally, using equivalence relation (3.39) we have the exponential decay of the energy E . This completes the proof of Theorem 3.1. $\square$

Remarks 3.5. When considering 2-dimensional plates instead of 1-dimensional beams, there are mainly two kinds of difficulties. Firstly, the control of some unwanted tangential derivatives on the boundary where the support f is acting. However, it seems that a compactness argument similar to the one in [14] may be used to show exponential decay for $q = 1$. But polynomial decay for $q > 1$ seems to be a harder question. The second kind of difficulties lies in the lack of formal results on the existence and regularity for stationary plate equations with transmission conditions similar to (1.3), which is essential when using multipliers techniques.

Acknowledgment

This work was partially supported by CNPq/Brazil.

References

- [1] M. M. Cavalcanti, V. N. Domingos Cavalcanti, and T. F. Ma, *Exponential decay of the viscoelastic Euler-Bernoulli equation with a nonlocal dissipation in general domains*, Differential and Integral Equations **17** (2004), no. 5-6, 495–510.
- [2] E. Feireisl, *Nonzero time periodic solutions to an equation of Petrovsky type with nonlinear boundary conditions: slow oscillations of beams on elastic bearings*, Annali della Scuola Normale Superiore di Pisa. Classe di Scienze **20** (1993), no. 1, 133–146.
- [3] M. R. Grossinho and T. F. Ma, *Symmetric equilibria for a beam with a nonlinear elastic foundation*, Portugaliae Mathematica **51** (1994), no. 3, 375–393.
- [4] B. V. Kapitonov and G. Perla Menzala, *Energy decay and a transmission problem in electromagneto-elasticity*, Advances in Differential Equations **7** (2002), no. 7, 819–846.
- [5] ———, *Uniform stabilization and exact control of a multilayered piezoelectric body*, Portugaliae Mathematica **60** (2003), no. 4, 411–454.
- [6] J. E. Lagnese, *Boundary Stabilization of Thin Plates*, SIAM Studies in Applied Mathematics, vol. 10, SIAM, Pennsylvania, 1989.
- [7] J. E. Lagnese and G. Leugering, *Uniform stabilization of a nonlinear beam by nonlinear boundary feedback*, Journal of Differential Equations **91** (1991), no. 2, 355–388.
- [8] J.-L. Lions, *Contrôlabilité Exacte, Perturbations et Stabilisation de Systèmes Distribués. Tome 1*, Collection: Recherches en Mathématiques Appliquées, vol. 8, Masson, Paris, 1988.

- [9] J.-L. Lions and E. Magenes, *Problèmes aux Limites non Homogènes et Applications. Vol. 1*, Travaux et Recherches Mathématiques, no. 17, Dunod, Paris, 1968.
- [10] W. Liu and G. Williams, *The exponential stability of the problem of transmission of the wave equation*, Bulletin of the Australian Mathematical Society **57** (1998), no. 2, 305–327.
- [11] T. F. Ma, *Existence results for a model of nonlinear beam on elastic bearings*, Applied Mathematics Letters **13** (2000), no. 5, 11–15.
- [12] ———, *Boundary stabilization for a non-linear beam on elastic bearings*, Mathematical Methods in the Applied Sciences **24** (2001), no. 8, 583–594.
- [13] R. Ma and Y. An, *Uniqueness of positive solutions of a class of ODE with nonlinear boundary conditions*, Boundary Value Problems **2005** (2005), no. 3, 289–298.
- [14] J. E. Muñoz Rivera and H. Portillo Oquendo, *Transmission problem for viscoelastic beams*, Advances in Mathematical Sciences and Applications **12** (2002), no. 1, 1–20.
- [15] S. Nicaise, *Boundary exact controllability of interface problems with singularities. II. Addition of internal controls*, SIAM Journal on Control and Optimization **35** (1997), no. 2, 585–603.
- [16] A. F. Pazoto and G. Perla Menzala, *Uniform stabilization of a nonlinear beam model with thermal effects and nonlinear boundary dissipation*, Funkcialaj Ekvacioj **43** (2000), no. 2, 339–360.

To Fu Ma: Department of Mathematics, State University of Maringá, 87020-900 Maringá, PR, Brazil

E-mail address: matofu@icmc.usp.br

Higdio Portillo Oquendo: Department of Mathematics, Federal University of Paraná, 81531-990 Curitiba, PR, Brazil

E-mail address: higdio@mat.ufpr.br

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru