

*Research Article*

## A New Iterative Algorithm for Approximating Common Fixed Points for Asymptotically Nonexpansive Mappings

H. Y. Zhou, Y. J. Cho, and S. M. Kang

Received 28 February 2007; Accepted 13 April 2007

Recommended by Nan-Jing Huang

Suppose that  $K$  is a nonempty closed convex subset of a real uniformly convex and smooth Banach space  $E$  with  $P$  as a sunny nonexpansive retraction. Let  $T_1, T_2 : K \rightarrow E$  be two weakly inward and asymptotically nonexpansive mappings with respect to  $P$  with sequences  $\{K_n\}, \{l_n\} \subset [1, \infty)$ ,  $\lim_{n \rightarrow \infty} k_n = 1$ ,  $\lim_{n \rightarrow \infty} l_n = 1$ ,  $F(T_1) \cap F(T_2) = \{x \in K : T_1x = T_2x = x\} \neq \emptyset$ , respectively. Suppose that  $\{x_n\}$  is a sequence in  $K$  generated iteratively by  $x_1 \in K$ ,  $x_{n+1} = \alpha_n x_n + \beta_n (PT_1)^n x_n + \gamma_n (PT_2)^n x_n$ , for all  $n \geq 1$ , where  $\{\alpha_n\}$ ,  $\{\beta_n\}$ , and  $\{\gamma_n\}$  are three real sequences in  $[\epsilon, 1 - \epsilon]$  for some  $\epsilon > 0$  which satisfy condition  $\alpha_n + \beta_n + \gamma_n = 1$ . Then, we have the following. (1) If one of  $T_1$  and  $T_2$  is completely continuous or demicompact and  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ ,  $\sum_{n=1}^{\infty} (l_n - 1) < \infty$ , then the strong convergence of  $\{x_n\}$  to some  $q \in F(T_1) \cap F(T_2)$  is established. (2) If  $E$  is a real uniformly convex Banach space satisfying Opial's condition or whose norm is Fréchet differentiable, then the weak convergence of  $\{x_n\}$  to some  $q \in F(T_1) \cap F(T_2)$  is proved.

Copyright © 2007 H. Y. Zhou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### 1. Introduction

Let  $K$  be a nonempty closed convex subset of a real uniformly convex Banach space  $E$ . A self-mapping  $T : K \rightarrow K$  is said to be nonexpansive if  $\|T(x) - T(y)\| \leq \|x - y\|$  for all  $x, y \in K$ . A self-mapping  $T : K \rightarrow K$  is called asymptotically nonexpansive if there exist sequences  $\{k_n\} \subset [1, \infty)$ ,  $k_n \rightarrow 1$  as  $n \rightarrow \infty$  such that

$$\|T^n(x) - T^n(y)\| \leq k_n \|x - y\|, \quad \forall x, y \in K, n \geq 1. \quad (1.1)$$

## 2 Fixed Point Theory and Applications

A self-mapping  $T : K \rightarrow K$  is said to be uniformly  $L$ -Lipschitzian if there exists constant  $L > 0$  such that

$$\|T^n(x) - T^n(y)\| \leq L\|x - y\|, \quad \forall x, y \in K, n \geq 1. \quad (1.2)$$

A self-mapping  $T : K \rightarrow K$  is called asymptotically quasi-nonexpansive if  $F(T) \neq \emptyset$  and there exist sequences  $\{k_n\} \subset [1, \infty)$  with  $k_n \rightarrow 1$  as  $n \rightarrow \infty$  such that

$$\|T^n(x) - p\| \leq k_n\|x - p\|, \quad \forall x \in K, p \in F(T), n \geq 1. \quad (1.3)$$

It is clear that, if  $T$  is an asymptotically nonexpansive mapping from  $K$  into itself with a fixed point in  $K$ , then  $T$  is asymptotically quasi-nonexpansive, but the converse may be not true.

As a generalization of the class of nonexpansive maps, the class of asymptotically nonexpansive mappings was introduced by Goebel and Kirk [1] in 1972, who proved that if  $K$  is a nonempty bounded closed convex subset of a real uniformly convex Banach space and  $T$  is an asymptotically nonexpansive self-mapping of  $K$ , then  $T$  has a fixed point.

In 1978, Bose [2] first proved that if  $K$  is a nonempty bounded closed convex subset of a real uniformly convex Banach space  $E$  satisfying Opial's condition and  $T : K \rightarrow K$  is an asymptotically nonexpansive mapping, then the sequence  $\{T^n x\}$  converges weakly to a fixed point of  $T$ , provided that  $T$  is asymptotically regular at  $x \in K$ , that is,

$$\lim_{n \rightarrow \infty} \|T^n x - T^{n+1} x\| = 0. \quad (1.4)$$

In 1982, Passty [3] proved that Bose's weak convergence theorem still holds if Opial's condition is replaced by the condition that  $E$  has a Fréchet differentiable norm.

Furthermore, Tan and Xu [4, 5] later proved that the asymptotic regularity of  $T$  at  $x$  can be weakened to the weakly asymptotic regularity of  $T$  at  $x$ , that is,

$$\omega - \lim_{n \rightarrow \infty} (T^n x - T^{n+1} x) = 0. \quad (1.5)$$

In all the above results ( $x_n = T^n x$ ), the asymptotic regularity of  $T$  at  $x \in K$  is equivalent to  $x_n - T x_n \rightarrow 0$  as  $n \rightarrow \infty$ . We wish that the later is a conclusion rather than an assumption.

In 1991, Schu [6, 7] introduced a modified Mann iterative algorithm to approximate fixed points of asymptotically nonexpansive maps without assuming the asymptotic regularity of  $T$  at  $x \in K$ . Schu established the conclusion that  $x_n - T x_n \rightarrow 0$  as  $n \rightarrow \infty$  by choosing properly iterative parameters  $\{\alpha_n\}$ .

Schu's iterative algorithm was defined as follows:

$$\begin{aligned} x_1 &\in K, \\ x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T^n x_n, \quad \forall n \geq 1. \end{aligned} \quad (1.6)$$

Since then, many authors have developed Schu's algorithm and results. Rhoades [8] and Tan and Xu [4] generalized Schu's iterative algorithm to the modified Ishikawa iterative algorithm and extended the main results of Schu to uniformly convex Banach spaces.

Furthermore, Osilike and Aniagbosor [9] improved the main results of Schu [6]. Schu [7] and Rhoades [8], without assuming the boundedness condition, imposed on  $K$ . Recently, Chang et al. [10] established a more general demiclosed principle and improved the corresponding results of Bose [2], Górnicki [11], Passty [3], Reich [12], Schu [6, 7], and Tan and Xu [4, 5].

Some iterative algorithms for approximating fixed points of nonself nonexpansive mappings have been studied by various authors (see [13–18]). However, iterative algorithms for approximating fixed points of nonself asymptotically nonexpansive mappings have not been paid too much attention. The main reason is the fact that when  $T$  is not a self-mapping, the mapping  $T^n$  is nonsensical. Recently, in order to establish the convergence theorems for non-self-asymptotically nonexpansive mappings, Chidume et al. [19] introduced the following definition.

*Definition 1.1.* Let  $K$  be a nonempty subset of real-normed linear space  $E$ . Let  $P : E \rightarrow K$  be the nonexpansive retraction of  $E$  onto  $K$ .

(1) A non-self-mapping  $T : K \rightarrow E$  is called *asymptotically nonexpansive* if there exists a sequence  $\{k_n\} \subset [1, \infty)$  with  $k_n \rightarrow 1$  as  $n \rightarrow \infty$  such that

$$\|T(PT)^{n-1}(x) - T(PT)^{n-1}(y)\| \leq k_n \|x - y\|, \quad \forall x, y \in K, n \geq 1. \quad (1.7)$$

(2)  $T$  is said to be *uniformly  $L$ -Lipschitzian* if there exists a constant  $L > 0$  such that

$$\|T(PT)^{n-1}(x) - T(PT)^{n-1}(y)\| \leq L \|x - y\|, \quad \forall x, y \in K, n \geq 1. \quad (1.8)$$

By using the following iterative algorithm:

$$\begin{aligned} x_1 &\in K, \\ x_{n+1} &= P((1 - \alpha_n)x_n + \alpha_n T(PT)^{n-1}x_n), \quad \forall n \geq 1, \end{aligned} \quad (1.9)$$

Chidume et al. [19] established the following demiclosed principle, strong and weak convergence theorems for non-self-asymptotically nonexpansive mappings in uniformly convex Banach spaces.

**THEOREM 1.2** [19]. *Let  $E$  be a uniformly convex Banach space,  $K$  a nonempty closed convex subset of  $E$ . Let  $T : K \rightarrow E$  be an asymptotically nonexpansive mapping with a sequence  $\{k_n\} \subset [1, \infty)$  and  $k_n \rightarrow 1$  as  $n \rightarrow \infty$ . Then  $I - T$  is demiclosed at zero.*

**THEOREM 1.3** [19]. *Let  $E$  be a uniformly convex Banach space and let  $K$  be a nonempty closed convex subset of  $E$ . Let  $T : K \rightarrow E$  be completely continuous and asymptotically nonexpansive mapping with a sequence  $\{k_n\} \subset [1, \infty)$  such that  $\sum_{n=1}^{\infty} (k_n^2 - 1) < \infty$ , and  $F(T) \neq \emptyset$ . Let  $\{\alpha_n\} \subset (0, 1)$  be a sequence such that  $\epsilon \leq 1 - \alpha_n \leq 1 - \epsilon$  for all  $n \geq 1$  and some  $\epsilon > 0$ . For an arbitrary point  $x_1 \in K$ , define the sequence  $\{x_n\}$  by (1.9). Then,  $\{x_n\}$  converges strongly to some fixed point of  $T$ .*

**THEOREM 1.4** [19]. *Let  $E$  be a uniformly convex Banach space which has a Fréchet differentiable norm and let  $K$  be a nonempty closed convex subset of  $E$ . Let  $T : K \rightarrow E$  be an asymptotically nonexpansive mapping with a sequence  $\{k_n\} \subset [1, \infty)$  such that  $\sum_{n=1}^{\infty} (k_n^2 - 1) < \infty$  and  $F(T) \neq \emptyset$ . Let  $\{\alpha_n\} \subset (0, 1)$  be a sequence such that  $\epsilon \leq 1 - \alpha_n \leq 1 - \epsilon$  for all  $n \geq 1$*

## 4 Fixed Point Theory and Applications

and some  $\epsilon > 0$ . For an arbitrary point  $x_1 \in K$ , let  $\{x_n\}$  be the sequence defined by (1.9). Then  $\{x_n\}$  converges weakly to some fixed point of  $T$ .

We now introduce the following definition.

**Definition 1.5.** Let  $K$  be a nonempty subset of real normed linear space  $E$ . Let  $P : E \rightarrow K$  be a nonexpansive retraction of  $E$  onto  $K$ .

(1) A non-self-mapping  $T : K \rightarrow E$  is called *asymptotically nonexpansive* with respect to  $P$  if there exists a sequence  $\{k_n\} \subset [1, \infty)$  with  $k_n \rightarrow 1$  as  $n \rightarrow \infty$  such that

$$\|(PT)^n x - (PT)^n y\| \leq k_n \|x - y\|, \quad \forall x, y \in K, n \geq 1. \quad (1.10)$$

(2)  $T$  is said to be *uniformly  $L$ -Lipschitzian* with respect to  $P$  if there exists a constant  $L > 0$  such that

$$\|(PT)^n x - (PT)^n y\| \leq L \|x - y\|, \quad \forall x, y \in K, n \geq 1. \quad (1.11)$$

**Remark 1.6.** If  $T$  is self-mapping, then  $P$  becomes the identity mapping, so that (1.7), (1.8), and (1.9) reduce to (1.1), (1.2), and (1.6), respectively.

We remark in the passing that if  $T : K \rightarrow E$  is asymptotically nonexpansive in light of (1.7) and  $P : E \rightarrow K$  is a nonexpansive retraction, then  $PT : K \rightarrow K$  is asymptotically nonexpansive in light of (1.1). Indeed, by definition (1.7), we have

$$\begin{aligned} & \|(PT)^n x - (PT)^n y\| \\ &= \|PT(PT)^{n-1} x - PT(PT)^{n-1} y\| \\ &\leq \|T(PT)^{n-1} x - T(PT)^{n-1} y\| \\ &\leq k_n \|x - y\|, \quad \forall x, y \in K, n \geq 1. \end{aligned} \quad (1.12)$$

Conversely, it may not be true.

It is our purpose in this paper to introduce a new iterative algorithm (see (2.6)) for approximating common fixed points of two non-self-asymptotically nonexpansive mappings with respect to  $P$  and to prove some strong and weak convergence theorems for such mappings in uniformly convex Banach spaces. As a consequence, the main results of Chidume et al. [19] are deduced.

## 2. Preliminaries

In this section, we will introduce a new iterative algorithm and prove a new demiclosedness principle for a non-self-asymptotically nonexpansive mapping in the sense of (1.10).

Let  $E$  be a Banach space with dimension  $E \geq 2$ . The modulus of  $E$  is the function  $\delta_E : (0, 2] \rightarrow [0, 1]$  defined by

$$\delta_E(\epsilon) = \inf \left\{ 1 - \left\| \frac{1}{2}(x + y) \right\| : \|x\| = 1, \|y\| = 1, \epsilon = \|x - y\| \right\}. \quad (2.1)$$

A Banach space  $E$  is uniformly convex if and only if  $\delta_E(\epsilon) > 0$  for all  $\epsilon \in (0, 2]$ .

A subset  $K$  of  $E$  is said to be retract if there exists a continuous mapping  $P : E \rightarrow K$  such that  $Px = x$  for all  $x \in K$ . Every closed convex subset of a uniformly convex Banach space is a retraction. A mapping  $P : E \rightarrow E$  is said to be a retraction if  $P^2 = P$ . Note that if a mapping  $P$  is a retraction, then  $Pz = z$  for all  $z \in R(P)$ , the range of  $P$ .

Let  $E$  be a Banach space and let  $C, D$  be subsets of  $E$ . Then, a mapping  $P : C \rightarrow D$  is said to be sunny if

$$P(Px + t(x - Px)) = Px, \quad (2.2)$$

whenever  $Px + t(x - Px) \in C$  for all  $x \in C$  and  $t \geq 0$ .

Let  $K$  be a subset of a Banach space  $E$ . For all  $x \in K$ , define a set  $I_K(x)$  by

$$I_K(x) = \{x + \lambda(y - x) : \lambda > 0, y \in K\}. \quad (2.3)$$

A non-self-mapping  $T : K \rightarrow E$  is said to be inward if  $Tx \in I_k(x)$  for all  $x \in K$  and  $T$  is said to be weakly inward if  $Tx \in \overline{I_K(x)}$  for all  $x \in K$ .

The following facts are well known (see [20, 18]).

**LEMMA 2.1.** *Let  $C$  be a nonempty convex subset of a smooth Banach space  $E$ ,  $C_0 \subset C$ , let  $J : E \rightarrow E^*$  be the normalized duality mapping of  $E$ , and let  $P : C \rightarrow C_0$  be a retraction. Then, the following statements are equivalent:*

- (1)  $\langle x - Px, J(y - Px) \rangle \leq 0$  for all  $x \in C$  and  $y \in C_0$ ;
- (2)  $P$  is both sunny and nonexpansive.

**LEMMA 2.2.** *Let  $E$  be a real smooth Banach space, let  $K$  be a nonempty closed convex subset of  $E$  with  $P$  as a sunny nonexpansive retraction, and let  $T : K \rightarrow E$  be a mapping satisfying weakly inward condition. Then  $F(PT) = F(T)$ .*

A Banach space  $E$  is said to satisfy Opial's condition if for any sequence  $\{x_n\}$  in  $E$ ,  $x_n \rightharpoonup x$  implies that

$$\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\| \quad (2.4)$$

for all  $y \in E$  with  $y \neq x$ , where  $x_n \rightharpoonup x$  denotes that  $\{x_n\}$  converges weakly to  $x$ . It is well known that Hilbert space and  $l^p$  ( $1 < p < \infty$ ) admit Opial's property, while  $L^p$  does not unless  $p = 2$ .

Let  $E$  be a Banach space and  $S(E) = \{x \in E : \|x\| = 1\}$ . The space  $E$  is said to be smooth if

$$\lim_{t \rightarrow 0} \frac{\|x + ty\| - \|x\|}{t} \quad (2.5)$$

exists for all  $x, y \in S(E)$ . For any  $x, y \in E$  ( $x \neq 0$ ), we denote this limit by  $(x, y)$ . The norm  $\|\cdot\|$  of  $E$  is said to be Fréchet differentiable if for all  $x \in S(E)$ , the limit  $(x, y)$  exists uniformly for all  $y \in S(E)$ .

A mapping  $T$  with domain  $D(T)$  and range  $R(T)$  in  $E$  is said to be demiclosed at  $p$  if whenever  $\{x_n\}$  is a sequence in  $D(T)$  such that  $\{x_n\}$  converges to  $x^* \in D(T)$  and  $\{Tx_n\}$  converges strongly to  $p$ ,  $Tx^* = p$ .

## 6 Fixed Point Theory and Applications

Let  $E$  be a real normed linear space, let  $K$  be a nonempty closed convex subset of  $E$  which is also a nonexpansive retraction of  $E$  with a retraction  $P$ . Let  $T_1 : K \rightarrow E$  and  $T_2 : K \rightarrow E$  be two non-self-asymptotically nonexpansive mappings with respect to  $P$ . For approximating the common fixed points of two non-self-asymptotically nonexpansive mappings, we introduce the following iterative algorithm:

$$\begin{aligned} x_1 &\in K, \\ x_{n+1} &= \alpha_n x_n + \beta_n (PT_1)^n x_n + \gamma_n (PT_2)^n x_n, \quad \forall n \geq 1, \end{aligned} \tag{2.6}$$

where  $\{\alpha_n\}$ ,  $\{\beta_n\}$ , and  $\{\gamma_n\}$  are three real sequences in  $(0, 1)$  satisfying  $\alpha_n + \beta_n + \gamma_n = 1$ .

LEMMA 2.3 [21]. *Let  $\{\alpha_n\}$  and  $\{t_n\}$  be two nonnegative real sequences satisfying*

$$\alpha_{n+1} \leq \alpha_n + t_n, \quad \forall n \geq 1. \tag{2.7}$$

*If  $\sum_{n=1}^{\infty} t_n < \infty$ , then  $\lim_{n \rightarrow \infty} \alpha_n$  exists.*

The following lemma can be found in Zhou et al. [22].

LEMMA 2.4 [22]. *Let  $E$  be a real uniformly convex Banach space and let  $B_r(0)$  be the closed ball of  $E$  with centre at the origin and radius  $r > 0$ . Then, there exists a continuous strictly increasing convex function  $g : [0, \infty) \rightarrow [0, \infty)$  with  $g(0) = 0$  such that*

$$\|\lambda x + \mu y + \gamma z\|^2 \leq \lambda \|x\|^2 + \mu \|y\|^2 + \gamma \|z\|^2 - \lambda \mu g(\|x - y\|) \tag{2.8}$$

for all  $x, y, z \in B_r(0)$  and  $\lambda, \mu, \gamma \in [0, 1]$  with  $\lambda + \mu + \gamma = 1$ .

The following demiclosedness principle for non-self-mapping follows from [10, Theorem 1].

LEMMA 2.5. *Let  $E$  be a real smooth and uniformly convex Banach space and  $K$  a nonempty closed convex subset of  $E$  with  $P$  as a sunny nonexpansive retraction. Let  $T : K \rightarrow E$  be a weakly inward and asymptotically nonexpansive mapping with respect to  $P$  with a sequence  $\{k_n\} \subset [1, \infty)$  such that  $\{k_n\} \rightarrow 1$  as  $n \rightarrow \infty$ . Then  $I - T$  is demiclosed at zero, that is,  $x_n \rightarrow x$  and  $x_n - Tx_n \rightarrow 0$  imply that  $Tx = x$ .*

*Proof.* Suppose that  $\{x_n\} \subset K$  converges weakly to  $x^* \in K$  and  $x_n - Tx_n \rightarrow 0$  as  $n \rightarrow \infty$ . We will prove that  $Tx^* = x^*$ . Indeed, since  $\{x_n\} \subset K$ , by the property of  $P$ , we have  $Px_n = x_n$  for all  $n \geq 1$  and so  $x_n - PTx_n \rightarrow 0$  as  $n \rightarrow \infty$ . By Chang et al. [10, Theorem 1], we conclude that  $x^* = PTx^*$ . Since  $F(PT) = F(T)$  by Lemma 2.2, we have  $Tx^* = x^*$ . This completes the proof.  $\square$

Remark 2.6. Lemma 2.5 extends Chang et al. [10, Theorem 1] to non-self-mapping case.

Using the proof lines of Reich [12, Proposition], then we can prove the following lemma.

LEMMA 2.7. *Let  $K$  be a closed convex subset of a uniformly convex Banach space  $E$  with a Fréchet differentiable norm and let  $\{T_n : 1 \leq n \leq \infty\}$  be a family of Lipschitzian self-mappings of  $K$  with a nonempty common fixed point set  $F$  and a Lipschitzian constant*

sequence  $\{L_n\}$  such that  $\sum_{n=1}^{\infty} (L_n - 1) < \infty$ . If  $x_1 \in K$  and  $x_{n+1} = T_n x_n$  for  $n \geq 1$ , then  $\lim_{n \rightarrow \infty} (f_1 - f_2, x_n)$  exists for all  $f_1 \neq f_2 \in F$ .

*Remark 2.8.* Lemma 2.7 is an extension of a proposition due to Reich [12].

### 3. Main results

In this section, we present some several strong and weak convergence theorems for two non-self-asymptotically nonexpansive mappings with respect to  $P$ .

**LEMMA 3.1.** *Let  $K$  be a nonempty closed convex subset of a normed linear space  $E$ . Let  $T_1, T_2 : K \rightarrow E$  be two non-self-asymptotically nonexpansive mappings with respect to  $P$  with sequences  $\{k_n\}, \{l_n\} \subset [1, \infty)$ ,  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ ,  $\sum_{n=1}^{\infty} (l_n - 1) < \infty$ , respectively. Suppose that  $\{x_n\}$  is the sequence defined by (2.6). If  $F(T_1) \cap F(T_2) \neq \emptyset$ , then  $\lim_{n \rightarrow \infty} \|x_n - q\|$  and  $\lim_{n \rightarrow \infty} \|y_n - q\|$  exist for any  $q \in F(T_1) \cap F(T_2)$ .*

*Proof.* For any  $q \in F(T_1) \cap F(T_2)$ , using the fact that  $P$  is nonexpansive and (2.6), then we have

$$\begin{aligned} \|x_{n+1} - q\| &= \|(\alpha_n x_n + \beta_n (PT_1)^n x_n + \gamma_n (PT_2)^n x_n) - Pq\| \\ &\leq \alpha_n \|x_n - q\| + \beta_n k_n \|x_n - q\| + \gamma_n l_n \|x_n - q\| \\ &\leq m_n \|x_n - q\|, \end{aligned} \quad (3.1)$$

where  $m_n = \max\{k_n, l_n\}$  for all  $n \geq 1$ . It is clear that  $\sum_{n=1}^{\infty} (m_n - 1) < \infty$  by the assumptions on  $\{k_n\}$  and  $\{l_n\}$ . It follows from Lemma 2.3 that  $\lim_{n \rightarrow \infty} \|x_n - q\|$  exists. This completes the proof.  $\square$

**LEMMA 3.2.** *Let  $K$  be a nonempty closed convex subset of a real uniformly convex Banach space  $E$ . Let  $T_1, T_2 : K \rightarrow E$  be two non-self-asymptotically nonexpansive mappings with respect to  $P$  with sequences  $\{k_n\}, \{l_n\} \subset [1, \infty)$ ,  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ ,  $\sum_{n=1}^{\infty} (l_n - 1) < \infty$ , respectively. Suppose that  $\{x_n\}$  is the sequence defined by (2.6), where  $\{\alpha_n\}$ ,  $\{\beta_n\}$ , and  $\{\gamma_n\}$  are three sequences in  $[\epsilon, 1 - \epsilon]$  for some  $\epsilon > 0$ . If  $F(T_1) \cap F(T_2) \neq \emptyset$ , then*

$$\lim_{n \rightarrow \infty} \|x_n - (PT_1)x_n\| = \lim_{n \rightarrow \infty} \|x_n - (PT_2)x_n\| = 0. \quad (3.2)$$

*Proof.* From (2.6), by the property of  $P$ , and Lemma 2.4, we have

$$\begin{aligned} \|x_{n+1} - q\|^2 &\leq \|(\alpha_n x_n + \beta_n (PT_1)^n x_n + \gamma_n (PT_2)^n x_n) - q\|^2 \\ &= \|\alpha_n(x_n - q) + \beta_n((PT_1)^n x_n - q) + \gamma_n((PT_2)^n x_n - q)\|^2 \\ &\leq \alpha_n \|x_n - q\|^2 + \beta_n \|((PT_1)^n x_n - q)\|^2 + \gamma_n \|((PT_2)^n x_n - q)\|^2 \\ &\quad - \alpha_n \beta_n g(\|x_n - (PT_1)^n x_n\|) \\ &\leq m_n^2 \|x_n - q\|^2 - \epsilon^2 g(\|x_n - (PT_1)^n x_n\|), \end{aligned} \quad (3.3)$$

which implies that  $g(\|x_n - (PT_1)^n x_n\|) \rightarrow 0$  as  $n \rightarrow \infty$ . Since  $g : [0, \infty) \rightarrow [0, \infty)$  with  $g(0) = 0$  being a continuous strictly increasing convex function, we have  $x_n - (PT_1)^n x_n \rightarrow 0$  as

## 8 Fixed Point Theory and Applications

$n \rightarrow \infty$ . Consequently,  $x_n - (PT_1)x_n \rightarrow 0$  as  $n \rightarrow \infty$ . Similarly, we can prove that  $x_n - (PT_2)x_n \rightarrow 0$  as  $n \rightarrow \infty$ . This completes the proof.  $\square$

**THEOREM 3.3.** *Let  $K$  be a nonempty closed convex subset of a real smooth uniformly convex Banach space  $E$  with  $P$  as a sunny nonexpansive retraction. Let  $T_1, T_2 : K \rightarrow E$  be two weakly inward and asymptotically nonexpansive mappings with respect to  $P$  with sequences  $\{k_n\}, \{l_n\} \subset [1, \infty)$ ,  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ ,  $\sum_{n=1}^{\infty} (l_n - 1) < \infty$ , respectively. Let  $\{x_n\} \subset K$  be the sequence defined by (2.6), where  $\{\alpha_n\}$ ,  $\{\beta_n\}$ , and  $\{\gamma_n\}$  are three sequences in  $[\epsilon, 1 - \epsilon]$  for some  $\epsilon > 0$ . If one of  $T_1$  and  $T_2$  is completely continuous and  $F(T_1) \cap F(T_2) \neq \emptyset$ , then  $\{x_n\}$  converges strongly to a common fixed point of  $T_1$  and  $T_2$ .*

*Proof.* By Lemma 3.1,  $\lim_{n \rightarrow \infty} \|x_n - q\|$  exists for any  $q \in F$ . It is sufficient to show that  $\{x_n\}$  has a subsequence which converges strongly to a common fixed point of  $T_1$  and  $T_2$ . By Lemma 3.2,  $\lim_{n \rightarrow \infty} \|x_n - PT_1x_n\| = \lim_{n \rightarrow \infty} \|x_n - PT_2x_n\| = 0$ . Suppose that  $T_1$  is completely continuous. Noting that  $P$  is nonexpansive, we conclude that there exists subsequence  $\{PT_1x_{n_j}\}$  of  $\{PT_1x_n\}$  such that  $PT_1x_{n_j} \rightarrow q$ , and hence  $x_{n_j} \rightarrow q$  as  $j \rightarrow \infty$ . By the continuity of  $P$ ,  $T_1$ , and  $T_2$ , we have  $q = PT_1q = PT_2q$ , and so  $q \in F(T_1) \cap F(T_2)$  by Lemma 2.2. Thus,  $\{x_n\}$  converges strongly to a common fixed point  $q$  of  $T_1$  and  $T_2$ . This completes the proof.  $\square$

**THEOREM 3.4.** *Let  $K$  be a nonempty closed convex subset of a real smooth and uniformly convex Banach space  $E$  with  $P$  as a sunny nonexpansive retraction. Let  $T_1, T_2 : K \rightarrow E$  be two weakly inward asymptotically nonexpansive mappings with respect to  $P$  with sequences  $\{k_n\}, \{l_n\} \subset [1, \infty)$ ,  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ ,  $\sum_{n=1}^{\infty} (l_n - 1) < \infty$ , respectively. Let  $\{x_n\} \subset K$  be the sequence defined by (2.6), where  $\{\alpha_n\}$ ,  $\{\beta_n\}$ , and  $\{\gamma_n\}$  are three sequences in  $[\epsilon, 1 - \epsilon]$  for some  $\epsilon > 0$ . If one of  $T_1$  and  $T_2$  is demicompact and  $F(T_1) \cap F(T_2) \neq \emptyset$ , then  $\{x_n\}$  converges strongly to a common fixed point of  $T_1$  and  $T_2$ .*

*Proof.* Since one of  $T_1$  and  $T_2$  is demicompact, so is one of  $PT_1$  and  $PT_2$ . Suppose that  $PT_1$  is demicompact. Noting that  $\{x_n\}$  is bounded, we assert that there exists a subsequence  $\{PT_1x_{n_j}\}$  of  $\{PT_1x_n\}$  such that  $PT_1x_{n_j}$  converges strongly to  $q$ . By Lemma 3.2, we have  $x_{n_j} \rightarrow q$  as  $j \rightarrow \infty$ . Since  $P$ ,  $T_1$ , and  $T_2$  are all continuous, we have  $q = PT_1q = PT_2q$  and  $q \in F(T_1) \cap F(T_2)$  by Lemma 2.2. By Lemma 3.1, we know that  $\lim_{n \rightarrow \infty} \|x_n - q\|$  exists. Therefore,  $\{x_n\}$  converges strongly to  $q$  as  $n \rightarrow \infty$ . This completes the proof.  $\square$

**THEOREM 3.5.** *Let  $K$  be a nonempty closed convex subset of a real smooth and uniformly convex Banach space  $E$  satisfying Opial's condition or whose norm is Fréchet differentiable. Let  $T_1, T_2 : K \rightarrow E$  be two weakly inward and asymptotically nonexpansive mappings with respect to  $P$  with sequences  $\{k_n\}, \{l_n\} \subset [1, \infty)$ ,  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ ,  $\sum_{n=1}^{\infty} (l_n - 1) < \infty$ , respectively. Let  $\{x_n\} \subset K$  be the sequence defined by (2.6), where  $\{\alpha_n\}$ ,  $\{\beta_n\}$ , and  $\{\gamma_n\}$  are three sequences in  $[\epsilon, 1 - \epsilon]$  for some  $\epsilon > 0$ . If  $F(T_1) \cap F(T_2) \neq \emptyset$ , then  $\{x_n\}$  converges weakly to a common fixed point of  $T_1$  and  $T_2$ .*

*Proof.* For any  $q \in F(T_1) \cap F(T_2)$ , by Lemma 3.1, we know that  $\lim_{n \rightarrow \infty} \|x_n - q\|$  exists. We now prove that  $\{x_n\}$  has a unique weakly subsequential limit in  $F(T_1) \cap F(T_2)$ . First of all, Lemmas 2.2, 2.5, and 3.2 guarantee that each weakly subsequential limit of  $\{x_n\}$  is

a common fixed point of  $T_1$  and  $T_2$ . Secondly, Opial's condition and Lemma 2.7 guarantee that the weakly subsequential limit of  $\{x_n\}$  is unique. Consequently,  $\{x_n\}$  converges weakly to a common fixed point of  $T_1$  and  $T_2$ . This completes the proof.  $\square$

*Remark 3.6.* The main results of this paper can be extended to a finite family of non-self-asymptotically nonexpansive mappings  $\{T_i : 1 \leq i \leq m\}$ , where  $m$  is a fixed positive integer, by introducing the following iterative algorithm:

$$x_1 \in K, \quad (3.4)$$

$$x_{n+1} = \alpha_{n1}x_n + \alpha_{n2}(PT_1)^n x_n + \alpha_{n3}(PT_2)^n x_n + \cdots + \alpha_{n(m+1)}(PT_m)^n x_n,$$

where  $\{\alpha_{n1}\}, \{\alpha_{n2}\}, \dots$ , and  $\{\alpha_{n(m+1)}\}$  are  $m+1$  real sequences in  $(0, 1)$  satisfying  $\alpha_{n1} + \alpha_{n2} + \cdots + \alpha_{n(m+1)} = 1$ .

We close this section with the following open question.

How to devise an iterative algorithm for approximating common fixed points of an infinite family of non-self-asymptotically nonexpansive mappings?

### Acknowledgment

The first author was supported by National Natural Science Foundation of China Grant no. 10471033.

### References

- [1] K. Goebel and W. A. Kirk, "A fixed point theorem for asymptotically nonexpansive mappings," *Proceedings of the American Mathematical Society*, vol. 35, no. 1, pp. 171–174, 1972.
- [2] S. C. Bose, "Weak convergence to the fixed point of an asymptotically nonexpansive map," *Proceedings of the American Mathematical Society*, vol. 68, no. 3, pp. 305–308, 1978.
- [3] G. B. Passty, "Construction of fixed points for asymptotically nonexpansive mappings," *Proceedings of the American Mathematical Society*, vol. 84, no. 2, pp. 212–216, 1982.
- [4] K.-K. Tan and H. K. Xu, "The nonlinear ergodic theorem for asymptotically nonexpansive mappings in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 114, no. 2, pp. 399–404, 1992.
- [5] K.-K. Tan and H. K. Xu, "Fixed point iteration processes for asymptotically nonexpansive mappings," *Proceedings of the American Mathematical Society*, vol. 122, no. 3, pp. 733–739, 1994.
- [6] J. Schu, "Iterative construction of fixed points of asymptotically nonexpansive mappings," *Journal of Mathematical Analysis and Applications*, vol. 158, no. 2, pp. 407–413, 1991.
- [7] J. Schu, "Weak and strong convergence to fixed points of asymptotically nonexpansive mappings," *Bulletin of the Australian Mathematical Society*, vol. 43, no. 1, pp. 153–159, 1991.
- [8] B. E. Rhoades, "Fixed point iterations for certain nonlinear mappings," *Journal of Mathematical Analysis and Applications*, vol. 183, no. 1, pp. 118–120, 1994.
- [9] M. O. Osilike and S. C. Aniagbosor, "Weak and strong convergence theorems for fixed points of asymptotically nonexpansive mappings," *Mathematical and Computer Modelling*, vol. 32, no. 10, pp. 1181–1191, 2000.
- [10] S.-S. Chang, Y. J. Cho, and H. Y. Zhou, "Demi-closed principle and weak convergence problems for asymptotically nonexpansive mappings," *Journal of the Korean Mathematical Society*, vol. 38, no. 6, pp. 1245–1260, 2001.

## 10 Fixed Point Theory and Applications

- [11] J. Górnicki, "Weak convergence theorems for asymptotically nonexpansive mappings in uniformly convex Banach spaces," *Commentationes Mathematicae Universitatis Carolinae*, vol. 30, no. 2, pp. 249–252, 1989.
- [12] S. Reich, "Weak convergence theorems for nonexpansive mappings in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 67, no. 2, pp. 274–276, 1979.
- [13] S. Ishikawa, "Fixed points and iteration of a nonexpansive mapping in a Banach space," *Proceedings of the American Mathematical Society*, vol. 59, no. 1, pp. 65–71, 1976.
- [14] J. S. Jung and S. S. Kim, "Strong convergence theorems for nonexpansive nonself-mappings in Banach spaces," *Nonlinear Analysis*, vol. 33, no. 3, pp. 321–329, 1998.
- [15] S. H. Khan and H. Fukhar-ud-din, "Weak and strong convergence of a scheme with errors for two nonexpansive mappings," *Nonlinear Analysis*, vol. 61, no. 8, pp. 1295–1301, 2005.
- [16] S.-Y. Matsushita and D. Kuroiwa, "Strong convergence of averaging iterations of nonexpansive nonself-mappings," *Journal of Mathematical Analysis and Applications*, vol. 294, no. 1, pp. 206–214, 2004.
- [17] N. Shahzad, "Approximating fixed points of non-self nonexpansive mappings in Banach spaces," *Nonlinear Analysis*, vol. 61, no. 6, pp. 1031–1039, 2005.
- [18] Y. Song and R. Chen, "Viscosity approximation methods for nonexpansive nonself-mappings," *Journal of Mathematical Analysis and Applications*, vol. 321, no. 1, pp. 316–326, 2006.
- [19] C. E. Chidume, E. U. Ofoedu, and H. Zegeye, "Strong and weak convergence theorems for asymptotically nonexpansive mappings," *Journal of Mathematical Analysis and Applications*, vol. 280, no. 2, pp. 364–374, 2003.
- [20] W. Takahashi, *Nonlinear Functional Analysis. Fixed Point Theory and Its Applications*, Yokohama Publishers, Yokohama, Japan, 2000.
- [21] K.-K. Tan and H. K. Xu, "Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process," *Journal of Mathematical Analysis and Applications*, vol. 178, no. 2, pp. 301–308, 1993.
- [22] H. Y. Zhou, G. T. Guo, H. J. Hwang, and Y. J. Cho, "On the iterative methods for nonlinear operator equations in Banach spaces," *PanAmerican Mathematical Journal*, vol. 14, no. 4, pp. 61–68, 2004.

H. Y. Zhou: Department of Applied Mathematics, North China Electric Power University, Baoding 071003, China

*Email address:* witman66@yahoo.com.cn

Y. J. Cho: Department of Mathematics Education and RINS, College of Natural Sciences, Gyeongsang National University, Chinju 660-701, South Korea

*Email address:* yjcho@gsnu.ac.kr

S. M. Kang: Department of Mathematics Education and RINS, College of Natural Sciences, Gyeongsang National University, Chinju 660-701, South Korea

*Email address:* smkang@gsnu.ac.kr

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)