

Research Article

A Note on Asymptotic Contractions

Marina Arav, Francisco Eduardo Castillo Santos,
Simeon Reich, and Alexander J. Zaslavski

Received 27 August 2006; Revised 16 October 2006; Accepted 16 October 2006

Recommended by Brailey Sims

We provide sufficient conditions for the iterates of an asymptotic contraction on a complete metric space X to converge to its unique fixed point, uniformly on each bounded subset of X .

Copyright © 2007 Marina Arav et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let (X, d) be a complete metric space. The following theorem is the main result of Chen [1]. It improves upon Kirk's original theorem [2]. In this connection, see also [3, 4].

THEOREM 1.1. *Let $T : X \rightarrow X$ be such that*

$$d(T^n x, T^n y) \leq \phi_n(d(x, y)) \quad (1.1)$$

for all $x, y \in X$ and all natural numbers n , where $\phi_n : [0, \infty) \rightarrow [0, \infty)$ and $\lim_{n \rightarrow \infty} \phi_n = \phi$, uniformly on any bounded interval $[0, b]$. Suppose that ϕ is upper semicontinuous and that $\phi(t) < t$ for all $t > 0$. Furthermore, suppose that there exists a positive integer n_ such that ϕ_{n_*} is upper semicontinuous and $\phi_{n_*}(0) = 0$. If there exists $x_0 \in X$ which has a bounded orbit $O(x_0) = \{x_0, Tx_0, T^2x_0, \dots\}$, then T has a unique fixed point $x_* \in X$ and $\lim_{n \rightarrow \infty} T^n x = x_*$ for all $x \in X$.*

Note that Theorem 1.1 does not provide us with uniform convergence of the iterates of T on bounded subsets of X , although this does hold for many classes of mappings of contractive type (e.g., [5, 6]). This property is important because it yields stability of the

2 Fixed Point Theory and Applications

convergence of iterates even in the presence of computational errors [7]. In the present paper we show that this conclusion can be derived in the setting of Theorem 1.1. To this end, we first prove a somewhat more general result (Theorem 1.2) which, when combined with Theorem 1.1, yields our strengthening of Chen's result (Theorem 1.3).

THEOREM 1.2. *Let $x_* \in X$ be a fixed point of $T : X \rightarrow X$. Assume that*

$$d(T^n x, x_*) \leq \phi_n(d(x, x_*)) \quad \forall x \in X \text{ and all natural numbers } n, \quad (1.2)$$

where $\phi_n : [0, \infty) \rightarrow [0, \infty)$ and $\lim_{n \rightarrow \infty} \phi_n = \phi$, uniformly on any bounded interval $[0, b]$. Suppose that ϕ is upper semicontinuous and that $\phi(t) < t$ for all $t > 0$. Then $T^n x \rightarrow x_*$ as $n \rightarrow \infty$, uniformly on each bounded subset of X .

THEOREM 1.3. *Let $T : X \rightarrow X$ be such that*

$$d(T^n x, T^n y) \leq \phi_n(d(x, y)) \quad (1.3)$$

for all $x, y \in X$ and all natural numbers n , where $\phi_n : [0, \infty) \rightarrow [0, \infty)$ and $\lim_{n \rightarrow \infty} \phi_n = \phi$, uniformly on any bounded interval $[0, b]$. Suppose that ϕ is upper semicontinuous and that $\phi(t) < t$ for all $t > 0$. Furthermore, suppose that there exists a positive integer n_* such that ϕ_{n_*} is upper semicontinuous and $\phi_{n_*}(0) = 0$. If there exists $x_0 \in X$ which has a bounded orbit $O(x_0) = \{x_0, Tx_0, T^2 x_0, \dots\}$, then T has a unique fixed point $x_* \in X$ and $\lim_{n \rightarrow \infty} T^n x = x_*$, uniformly on each bounded subset of X .

2. Proof of Theorem 1.2

We may assume without loss of generality that $\phi(0) = 0$ and $\phi_n(0) = 0$ for all integers $n \geq 1$.

For each $x \in X$ and each $r > 0$, set

$$B(x, r) = \{y \in X : d(x, y) \leq r\}. \quad (2.1)$$

We first prove three lemmas.

LEMMA 2.1. *Let $K > 0$. Then there exists a natural number \bar{q} such that for all integers $s \geq \bar{q}$,*

$$T^s(B(x_*, K)) \subset B(x_*, K + 1). \quad (2.2)$$

Proof. There exists a natural number \bar{q} such that for all integers $s \geq \bar{q}$,

$$|\phi_s(t) - \phi(t)| < 1 \quad \forall t \in [0, K]. \quad (2.3)$$

Let $s \geq \bar{q}$ be an integer. Then for all $x \in B(x_*, K)$,

$$d(T^s x, x_*) \leq \phi_s(d(x, x_*)) < \phi(d(x, x_*)) + 1 < d(x, x_*) + 1 < K + 1. \quad (2.4)$$

Lemma 2.1 is proved. \square

LEMMA 2.2. *Let $0 < \epsilon_1 < \epsilon_0$. Then there exists a natural number q such that for each integer $j \geq q$,*

$$T^j(B(x_*, \epsilon_1)) \subset B(x_*, \epsilon_0). \quad (2.5)$$

Proof. There exists an integer $q \geq 1$ such that for each integer $j \geq q$,

$$|\phi_j(t) - \phi(t)| < (\epsilon_0 - \epsilon_1)/2 \quad \forall t \in [0, \epsilon_0]. \quad (2.6)$$

Assume that

$$j \in \{q, q+1, \dots\}, \quad x \in B(x_*, \epsilon_1). \quad (2.7)$$

By (1.2) and (2.6),

$$\begin{aligned} d(T^j x, x_*) &\leq \phi_j(d(x, x_*)) < \phi(d(x, x_*)) + \frac{(\epsilon_0 - \epsilon_1)}{2} \\ &\leq \epsilon_1 + \frac{(\epsilon_0 - \epsilon_1)}{2} = \frac{(\epsilon_0 + \epsilon_1)}{2}. \end{aligned} \quad (2.8)$$

Lemma 2.2 is proved. \square

LEMMA 2.3. *Let $K, \epsilon > 0$. Then there exists a natural number q such that for each $x \in B(x_*, K)$,*

$$\min \{d(T^j x, x_*): j = 1, \dots, q\} \leq \epsilon. \quad (2.9)$$

Proof. By Lemma 2.1, there is a natural number \bar{q} such that

$$T^n(B(x_*, K)) \subset B(x_*, K+1) \text{ for all natural numbers } n \geq \bar{q}. \quad (2.10)$$

We may assume without loss of generality that $\epsilon < K/8$. Since the function $t - \phi(t)$, $t \in (0, \infty)$, is lower semicontinuous and positive, there is

$$\delta \in \left(0, \frac{\epsilon}{8}\right) \quad (2.11)$$

such that

$$t - \phi(t) \geq 2\delta \quad \forall t \in \left[\frac{\epsilon}{2}, K+1\right]. \quad (2.12)$$

4 Fixed Point Theory and Applications

There is a natural number $s \geq \bar{q}$ such that

$$|\phi(t) - \phi_s(t)| \leq \delta \quad \forall t \in [0, K+1]. \quad (2.13)$$

By (2.12) and (2.13), we have, for all $t \in [\epsilon/2, K+1]$,

$$\phi_s(t) \leq \phi(t) + \delta \leq t - 2\delta + \delta = t - \delta. \quad (2.14)$$

In view of (2.13) and (2.11), we have, for all $t \in [0, \epsilon/2]$,

$$\phi_s(t) \leq \phi(t) + \delta \leq t + \delta \leq \frac{\epsilon}{2} + \delta < \frac{3}{4}\epsilon. \quad (2.15)$$

Choose a natural number p such that

$$p > 4 + \delta^{-1}(K+1). \quad (2.16)$$

Let

$$x \in B(x_*, K). \quad (2.17)$$

We will show that

$$\min \{d(T^j x, x_*): j = 1, 2, \dots, ps\} \leq \epsilon. \quad (2.18)$$

Let us assume the contrary. Then

$$d(T^j x, x_*) > \epsilon \quad \forall j = s, \dots, ps. \quad (2.19)$$

By (2.17) and (2.10),

$$T^j x \in B(x_*, K+1), \quad j = s, \dots, ps. \quad (2.20)$$

Let a natural number i satisfy $i \leq p-1$. By (2.19) and (2.20),

$$d(T^{is} x, x_*) > \epsilon, \quad d(T^{is} x, x_*) \leq K+1. \quad (2.21)$$

It follows from (1.2), (2.21), and (2.14) that

$$d(T^s(T^{is} x), x_*) \leq \phi_s(d(T^{is} x, x_*)) \leq d(T^{is} x, x_*) - \delta. \quad (2.22)$$

Thus for each natural number $i \leq p - 1$,

$$d(T^{(i+1)s}x, x_*) \leq d(T^{is}x, x_*) - \delta. \quad (2.23)$$

This inequality implies that

$$d(T^{ps}x, x_*) \leq d(T^{(p-1)s}x, x_*) - \delta \leq \dots \leq d(T^s x, x_*) - (p-1)\delta. \quad (2.24)$$

When combined with (2.20) and (2.16), this implies, in turn, that

$$d(T^{ps}x, x_*) \leq K + 1 - (p-1)\delta < 0. \quad (2.25)$$

The contradiction we have reached proves (2.18) and completes the proof of Lemma 2.3. \square

Completion of the proof of Theorem 1.2. Let $K, \epsilon > 0$. Choose $\epsilon_1 \in (0, \epsilon)$. By Lemma 2.2, there exists a natural number q_1 such that

$$T^j(B(x_*, \epsilon_1)) \subset B(x_*, \epsilon) \text{ for all integers } j \geq q_1. \quad (2.26)$$

By Lemma 2.3, there exists a natural number q_2 such that

$$\min \{d(T^j x, x_*): j = 1, \dots, q_2\} \leq \epsilon_1 \quad \forall x \in B(x_*, K). \quad (2.27)$$

Assume that

$$x \in B(x_*, K). \quad (2.28)$$

By (2.27), there is a natural number $j_1 \leq q_2$ such that

$$d(T^{j_1}x, x_*) \leq \epsilon_1. \quad (2.29)$$

In view of (2.29) and (2.26),

$$T^j(T^{j_1}x) \in B(x_*, \epsilon) \text{ for all integers } j \geq q_1. \quad (2.30)$$

Inclusion (2.30) and the inequality $j_1 \leq q_2$ now imply that

$$T^i x \in B(x_*, \epsilon) \text{ for all integers } i \geq q_1 + q_2. \quad (2.31)$$

Theorem 1.2 is proved.

Acknowledgments

Part of the first author's research was carried out when she was visiting the Technion—Israel Institute of Technology. The third author was partially supported by the Fund for the Promotion of Research at the Technion and by the Technion VPR Fund—B. and G. Greenberg Research Fund (Ottawa).

6 Fixed Point Theory and Applications

References

- [1] Y.-Z. Chen, “Asymptotic fixed points for nonlinear contractions,” *Fixed Point Theory and Applications*, vol. 2005, no. 2, pp. 213–217, 2005.
- [2] W. A. Kirk, “Fixed points of asymptotic contractions,” *Journal of Mathematical Analysis and Applications*, vol. 277, no. 2, pp. 645–650, 2003.
- [3] I. D. Arandelić, “On a fixed point theorem of Kirk,” *Journal of Mathematical Analysis and Applications*, vol. 301, no. 2, pp. 384–385, 2005.
- [4] J. Jachymski and I. Józwik, “On Kirk’s asymptotic contractions,” *Journal of Mathematical Analysis and Applications*, vol. 300, no. 1, pp. 147–159, 2004.
- [5] F. E. Browder, “On the convergence of successive approximations for nonlinear functional equations,” *Indagationes Mathematicae*, vol. 30, pp. 27–35, 1968.
- [6] E. Rakotch, “A note on contractive mappings,” *Proceedings of the American Mathematical Society*, vol. 13, no. 3, pp. 459–465, 1962.
- [7] D. Butnariu, S. Reich, and A. J. Zaslavski, “Asymptotic behavior of inexact orbits for a class of operators in complete metric spaces,” to appear in *Journal of Applied Analysis*.

Marina Arav: Department of Mathematics and Statistics, Georgia State University, Atlanta, GA 30303, USA
Email address: matmxa@langate.gsu.edu

Francisco Eduardo Castillo Santos: School of Mathematical and Physical Sciences, The University of Newcastle, Newcastle, NSW 2308, Australia
Email address: francisco.castillosantos@studentmail.newcastle.edu.au

Simeon Reich: Department of Mathematics, The Technion–Israel Institute of Technology, Haifa 32000, Israel
Email address: sreich@techunix.technion.ac.il

Alexander J. Zaslavski: Department of Mathematics, The Technion–Israel Institute of Technology, Haifa 32000, Israel
Email address: ajzasl@tx.technion.ac.il

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2009
First Round of Reviews	May 1, 2009
Publication Date	August 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Department of Physics, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk