

Research Article

Hybrid Iteration Method for Fixed Points of Nonexpansive Mappings in Arbitrary Banach Spaces

M. O. Osilike, F. O. Isiogugu, and P. U. Nwokoro

Received 20 June 2007; Accepted 23 November 2007

Recommended by Nanjing Huang

We prove that recent results of Wang (2007) concerning the iterative approximation of fixed points of nonexpansive mappings using a hybrid iteration method in Hilbert spaces can be extended to arbitrary Banach spaces without the strong monotonicity assumption imposed on the hybrid operator.

Copyright © 2007 M. O. Osilike et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let E be a real Banach space. A mapping $T : E \rightarrow E$ is said to be *L-Lipschitzian* if there exists $L > 0$ such that

$$\|Tx - Ty\| \leq L\|x - y\|, \quad \forall x, y \in E. \quad (1.1)$$

T is said to be *nonexpansive* if $L = 1$ in (1.1).

Several authors have studied various methods for the iterative approximation of fixed points of nonexpansive mappings. Recently, Wang [1] studied the following iteration method in Hilbert spaces.

The hybrid iteration method. Let H be a Hilbert space, $T : H \rightarrow H$ a nonexpansive mapping with $F(T) = \{x \in H : Tx = x\} \neq \emptyset$, and $F : H \rightarrow H$ an L -Lipschitzian mapping which is also η -strongly monotone, where T is η -strongly monotone if there exists $\eta > 0$ such that

$$\langle Tx - Ty, x - y \rangle \geq \eta\|x - y\|^2, \quad \forall x, y \in H. \quad (1.2)$$

2 Fixed Point Theory and Applications

Let $\{\alpha_n\}_{n=1}^{\infty}$ and $\{\lambda_n\}_{n=1}^{\infty}$ be real sequences in $[0, 1)$, and $\mu > 0$, then the sequence $\{x_n\}_{n=1}^{\infty}$ is generated from an arbitrary $x_1 \in H$ by

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T^{\lambda_{n+1}} x_n, \quad n \geq 1, \quad (1.3)$$

where $T^{\lambda_{n+1}} x_n := T x_n - \lambda_{n+1} \mu F(T x_n)$, $\mu > 0$. Wang's work was motivated by earlier results of Xu and Kim [2] and Yamada [3], in addition to several other related results. Using this iteration method, Wang proved the following main results.

LEMMA 1.1 (see [1, page 3]). *Let H be a Hilbert space, $T : H \rightarrow H$ a nonexpansive mapping with $F(T) = \{x \in H : Tx = x\} \neq \emptyset$, and $F : H \rightarrow H$ an η -strongly monotone and L -Lipschitzian mapping. Let $\{x_n\}_{n=1}^{\infty}$ be the sequence generated from an arbitrary $x_1 \in H$ by*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T^{\lambda_{n+1}} x_n, \quad n \geq 1, \quad (1.4)$$

where $T^{\lambda_{n+1}} x_n := T x_n - \lambda_{n+1} \mu F(T x_n)$, $\mu > 0$, and let $\{\alpha_n\}_{n=1}^{\infty}$ and $\{\lambda_n\}_{n=1}^{\infty}$ be real sequences in $[0, 1)$ satisfying the following conditions:

- (i) $0 < \alpha \leq \alpha_n \leq \beta < 1$, for some $\alpha, \beta \in (0, 1)$,
- (ii) $\sum_{n=1}^{\infty} \lambda_n < \infty$,
- (iii) $0 < \mu < 2\eta/L^2$.

Then,

- (a) $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists for each $x^* \in F(T)$,
- (b) $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$.

THEOREM 1.2 (see [1, page 5]). *Let H , T , $F(T)$, F , $\{T^{\lambda_{n+1}}\}_{n=1}^{\infty}$, $\{x_n\}_{n=1}^{\infty}$, $\{\alpha_n\}_{n=1}^{\infty}$, $\{\lambda_n\}_{n=1}^{\infty}$, μ, α , and β be as in Lemma 1.1. Let $\{x_n\}_{n=1}^{\infty}$ be the sequence generated from an arbitrary $x_1 \in H$ by*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T^{\lambda_{n+1}} x_n, \quad n \geq 1. \quad (1.5)$$

Then,

- (a) $\{x_n\}_{n=1}^{\infty}$ converges weakly to a fixed point of T ,
- (b) $\{x_n\}_{n=1}^{\infty}$ converges strongly to a fixed point of T if and only if $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$, where $d(x, F(T)) := \inf \{\|x - p\| : p \in F(T)\}$.

It is our purpose in this paper to extend Lemma 1.1 and Theorem 1.2 from Hilbert spaces to arbitrary Banach spaces. Our results are much more general and applicable than the results of Wang [1] because the strong monotonicity condition imposed on F by Wang is not required in our results.

2. Preliminaries

In the sequel, we will need what follows.

A Banach space E is said to satisfy *Opial's condition* (see, e.g., [4]) if for each sequence $\{x_n\}_{n=1}^{\infty}$ in E which converges weakly to a point $x \in E$, we have

$$\liminf_{n \rightarrow \infty} \|x_n - x\| < \liminf_{n \rightarrow \infty} \|x_n - y\|, \quad \forall y \in E. \quad (2.1)$$

Let E be a Banach space. A mapping T with domain $D(T)$ and range $R(T)$ in E is said to be *demiclosed at a point* $p \in D(T)$ if, whenever, $\{x_n\}_{n=1}^\infty$ is a sequence in E which converges weakly to a point $x \in E$ and $\{Tx_n\}_{n=1}^\infty$ converges strongly to p , then $Tx = p$. Furthermore, T is said to be *demicompact* if, whenever, $\{x_n\}_{n=1}^\infty$ is a bounded sequence in $D(T)$ such that $\{x_n - Tx_n\}_{n=1}^\infty$ converges strongly, then $\{x_n\}_{n=1}^\infty$ has a subsequence which converges strongly. T is said to satisfy condition (A) if $F(T) \neq \emptyset$ and there exists a nondecreasing function $f : [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$ and $f(t) > 0$ for all $t \in (0, \infty)$ such that $\|x - Tx\| \geq f(d(x, F(T)))$ for all $x \in D(T)$, where $d(x, F(T)) := \inf \{\|x - p\| : p \in F(T)\}$.

LEMMA 2.1 (see [5]). *Let E be a reflexive Banach space satisfying Opial's condition and let K be a nonempty closed convex subset of E . Let $T : K \rightarrow E$ be a nonexpansive mapping. Then, $(I - T)$ is demiclosed on K , where I is the identity mapping.*

LEMMA 2.2 (see [6, page 1184], [7]). *Let $\{a_n\}_{n=1}^\infty$, $\{b_n\}_{n=1}^\infty$, and $\{\delta_n\}_{n=1}^\infty$ be sequences of nonnegative real numbers satisfying the inequality*

$$a_{n+1} \leq (1 + \delta_n) a_n + b_n, \quad n \geq 1. \quad (2.2)$$

If $\sum_{n=1}^\infty \delta_n < \infty$ and $\sum_{n=1}^\infty b_n < \infty$, then $\lim_{n \rightarrow \infty} a_n$ exists. In particular, if $\{a_n\}_{n=1}^\infty$ has a subsequence which converges strongly to zero, then $\lim_{n \rightarrow \infty} a_n = 0$.

LEMMA 2.3 (see [8, page 770]). *Let E be an arbitrary normed space and let $\{t_n\}_{n=1}^\infty$ be a real sequence satisfying the following conditions:*

- (i) $0 \leq t_n \leq t < 1$ for all $n \geq 1$ and for some $t \in (0, 1)$,
- (ii) $\sum_{n=1}^\infty t_n = \infty$.

Let $\{u_n\}_{n=1}^\infty$ and $\{v_n\}_{n=1}^\infty$ be two sequences in E such that

- (iii) $u_{n+1} = (1 - t_n)u_n + t_n v_n$, $n \geq 1$,
- (iv) $\lim_{n \rightarrow \infty} \|u_n\| = d$ for some $d \in [0, \infty)$,
- (v) $\limsup_{n \rightarrow \infty} \|v_n\| \leq d$,
- (vi) $\{\sum_{j=1}^n t_j v_j\}_{n=1}^\infty$ is bounded.

Then, $d = 0$.

3. Main results

THEOREM 3.1. *Let E be an arbitrary real Banach space, $T : E \rightarrow E$ a nonexpansive mapping with $F(T) \neq \emptyset$, and $F : E \rightarrow E$ an L -Lipschitzian mapping. Let $\{x_n\}_{n=1}^\infty$ be the sequence generated from an arbitrary $x_1 \in E$ by*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T^{\lambda_{n+1}} x_n, \quad n \geq 1, \quad (3.1)$$

where $T^{\lambda_{n+1}} x_n := Tx_n - \lambda_{n+1} \mu F(Tx_n)$, $\mu > 0$, and $\{\alpha_n\}_{n=1}^\infty$ and $\{\lambda_n\}_{n=2}^\infty$ are real sequences in $[0, 1)$ satisfying the following conditions:

- (i) $0 < \alpha \leq \alpha_n < 1$ for all $n \geq 1$ and for some $\alpha \in (0, 1)$,
- (ii) $\sum_{n=1}^\infty (1 - \alpha_n) = \infty$,
- (iii) $\sum_{n=2}^\infty \lambda_n < \infty$.

4 Fixed Point Theory and Applications

Then,

- (a) $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists for each $x^* \in F(T)$,
- (b) $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$,
- (c) $\{x_n\}_{n=1}^{\infty}$ converges strongly to a fixed point of T if and only if $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$.

Proof. Let $x^* \in F(T)$ be arbitrary, then,

$$\begin{aligned}
\|x_{n+1} - x^*\| &= \|\alpha_n(x_n - x^*) + (1 - \alpha_n)(Tx_n - x^*) - (1 - \alpha_n)\lambda_{n+1}\mu F(Tx_n)\| \\
&\leq \|\alpha_n(x_n - x^*) + (1 - \alpha_n)(Tx_n - x^*)\| + (1 - \alpha_n)\lambda_{n+1}\mu \|F(Tx_n)\| \\
&\leq \alpha_n \|x_n - x^*\| + (1 - \alpha_n) \|Tx_n - x^*\| + (1 - \alpha_n)\lambda_{n+1}\mu \|F(Tx_n) - F(x^*)\| \\
&\quad + (1 - \alpha_n)\lambda_{n+1}\mu \|F(x^*)\| \\
&\leq \|x_n - x^*\| + (1 - \alpha_n)\lambda_{n+1}\mu L \|x_n - x^*\| + (1 - \alpha_n)\lambda_{n+1}\mu \|F(x^*)\| \\
&= [1 + \delta_n] \|x_n - x^*\| + \sigma_n,
\end{aligned} \tag{3.2}$$

where $\delta_n = (1 - \alpha_n)\lambda_{n+1}\mu L$ and $\sigma_n = (1 - \alpha_n)\lambda_{n+1}\mu \|F(x^*)\|$.

Since $\sum_{n=1}^{\infty} \delta_n < \infty$ and $\sum_{n=1}^{\infty} \sigma_n < \infty$, it follows from Lemma 2.2 that $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists. This completes the proof of (a).

Since $\{\|x_n - x^*\|\}_{n=1}^{\infty}$ is bounded, there exists $M > 0$ such that

$$\|x_n - x^*\| \leq M, \quad \forall n \geq 1. \tag{3.3}$$

Observe that

$$\begin{aligned}
\|x_{n+1} - Tx_{n+1}\| &= \|\alpha_n x_n + (1 - \alpha_n) T^{\lambda_{n+1}} x_n - Tx_{n+1}\| \\
&\leq \|\alpha_n(x_n - Tx_{n+1}) + (1 - \alpha_n)(Tx_n - Tx_{n+1})\| \\
&\quad + (1 - \alpha_n)\lambda_{n+1}\mu \|F(Tx_n)\| \\
&\leq \alpha_n \|x_n - Tx_{n+1}\| + (1 - \alpha_n) \|Tx_n - Tx_{n+1}\| \\
&\quad + (1 - \alpha_n)\lambda_{n+1}\mu \|F(Tx_n)\| \\
&\leq \alpha_n \|x_n - x_{n+1}\| + \alpha_n \|x_{n+1} - Tx_{n+1}\| + (1 - \alpha_n) \|x_n - x_{n+1}\| \\
&\quad + (1 - \alpha_n)\lambda_{n+1}\mu \|F(Tx_n)\|.
\end{aligned} \tag{3.4}$$

Thus,

$$\begin{aligned}
\|x_{n+1} - Tx_{n+1}\| &\leq \frac{1}{1 - \alpha_n} \|x_n - x_{n+1}\| + \lambda_{n+1}\mu \|F(Tx_n)\| \\
&\leq \|x_n - Tx_n\| + 2\lambda_{n+1}\mu \|F(Tx_n)\|.
\end{aligned} \tag{3.5}$$

It follows from (3.3) that

$$\|F(Tx_n)\| \leq L \|x_n - x^*\| + \|F(x^*)\| \leq LM + \|F(x^*)\| \leq D \tag{3.6}$$

for all $n \geq 1$ and for some $D > 0$. Using (3.6) in (3.5), we obtain

$$\|x_{n+1} - Tx_{n+1}\| \leq \|x_n - Tx_n\| + 2\lambda_{n+1}\mu D = \|x_n - Tx_n\| + \gamma_n, \quad (3.7)$$

where $\gamma_n = 2\lambda_{n+1}\mu D$. Since $\sum_{n=1}^{\infty} \gamma_n < \infty$, it follows from Lemma 2.2 that $\lim_{n \rightarrow \infty} \|x_n - Tx_n\|$ exists. Let $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = d$, and set $u_n = x_n - Tx_n$ so that

$$u_{n+1} = (1 - t_n)u_n + t_n v_n, \quad (3.8)$$

where $t_n = 1 - \alpha_n$ and $v_n = (1/(1 - \alpha_n))(Tx_n - Tx_{n+1}) - \lambda_{n+1}\mu F(Tx_n)$. Observe that $0 < t_n \leq 1 - \alpha = t \in (0, 1)$ and $\sum_{n=1}^{\infty} t_n = \infty$. Furthermore,

$$\begin{aligned} \|v_n\| &\leq \frac{1}{1 - \alpha_n} \|Tx_n - Tx_{n+1}\| + \lambda_{n+1}\mu \|F(Tx_n)\| \\ &\leq \frac{1}{1 - \alpha_n} \|x_n - x_{n+1}\| + \lambda_{n+1}\mu \|F(Tx_n)\| \\ &\leq \|x_n - Tx_n\| + 2\lambda_{n+1}\mu \|F(Tx_n)\| \\ &\leq \|x_n - Tx_n\| + 2\lambda_{n+1}\mu D. \end{aligned} \quad (3.9)$$

Thus, $\limsup_{n \rightarrow \infty} \|v_n\| \leq d$. Also,

$$\begin{aligned} \left\| \sum_{j=1}^n t_j v_j \right\| &= \left\| \sum_{j=1}^n (1 - \alpha_j) \left[\frac{1}{(1 - \alpha_j)} (Tx_j - Tx_{j+1}) - \lambda_{j+1}\mu F(Tx_j) \right] \right\| \\ &\leq \left\| \sum_{j=1}^n (Tx_j - Tx_{j+1}) \right\| + \sum_{j=1}^n (1 - \alpha_j) \lambda_{j+1}\mu \|F(Tx_j)\| \\ &< \|Tx_1 - Tx_{n+1}\| + \mu D \sum_{j=1}^n \lambda_{j+1} \\ &\leq \|x_1 - x_{n+1}\| + \mu D \sum_{j=1}^n \lambda_{j+1} \\ &\leq \|x_1 - x^*\| + \|x_{n+1} - x^*\| + \mu D \sum_{j=1}^{\infty} \lambda_{j+1} \leq K \end{aligned} \quad (3.10)$$

for all $n \geq 1$ and for some $K > 0$. Hence, $\{\sum_{j=1}^n t_j v_j\}_{n=1}^{\infty}$ is bounded. It now follows from Lemma 2.3 that $\lim_{n \rightarrow \infty} \|v_n\| = \lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$. This completes the proof of (b).

From (3.2), we obtain

$$\|x_{n+1} - x^*\| \leq [1 + \delta_n] \|x_n - x^*\| + \sigma_n \leq \|x_n - x^*\| + M\delta_n + \sigma_n = \|x_n - x^*\| + \beta_n, \quad (3.11)$$

where $\beta_n = M\delta_n + \sigma_n$. Hence, $d(x_{n+1}, F(T)) \leq d(x_n, F(T)) + \beta_n$. Since $\sum_{n=1}^{\infty} \beta_n < \infty$, it follows from Lemma 2.2 that $\lim_{n \rightarrow \infty} d(x_n, F(T))$ exists.

6 Fixed Point Theory and Applications

If $\{x_n\}_{n=1}^\infty$ converges strongly to a fixed point p of T , then $\lim_{n \rightarrow \infty} \|x_n - p\| = 0$. Since

$$0 \leq d(x_n, F(T)) \leq \|x_n - p\|, \quad (3.12)$$

we have $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$.

Conversely, suppose $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$, then we have $\lim_{n \rightarrow \infty} d(x_n, F(T)) = 0$. Thus for arbitrary $\epsilon > 0$, there exists a positive integer N_1 such that $d(x_n, F(T)) < \epsilon/4$ for all $n \geq N_1$. Furthermore, $\sum_{n=1}^\infty \beta_n < \infty$ implies that there exists a positive integer N_2 such that $\sum_{j=N}^\infty \beta_j < \epsilon/4$ for all $n \geq N_2$. Choose $N = \max\{N_1, N_2\}$, then $d(x_N, F(T)) < \epsilon/4$ and $\sum_{j=N}^\infty \beta_j < \epsilon/4$. It follows from (3.11) that for all $n, m \geq N$ and for all $p \in F(T)$, we have

$$\begin{aligned} \|x_n - x_m\| &\leq \|x_n - p\| + \|x_m - p\| \\ &\leq \|x_N - p\| + \sum_{j=N+1}^n \beta_j + \|x_N - p\| + \sum_{j=N+1}^m \beta_j \\ &\leq 2\|x_N - p\| + 2 \sum_{j=N}^\infty \beta_j. \end{aligned} \quad (3.13)$$

Taking infimum over all $p \in F(T)$, we obtain

$$\|x_n - x_m\| \leq 2d(x_N, F(T)) + 2 \sum_{j=N}^\infty \beta_j < \epsilon, \quad \forall n, m \geq N. \quad (3.14)$$

Thus, $\{x_n\}_{n=1}^\infty$ is Cauchy. Suppose $\lim_{n \rightarrow \infty} x_n = u$, then since $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$, we have $u \in F(T)$. This completes the proof of (c). \square

THEOREM 3.2. *Let E be a real reflexive Banach space satisfying Opial's condition, $T : E \rightarrow E$ a nonexpansive mapping with $F(T) \neq \emptyset$, and $F : E \rightarrow E$ an L -Lipschitzian mapping. Let $\{x_n\}_{n=1}^\infty$ be the sequence generated from an arbitrary $x_1 \in H$ by*

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T^{\lambda_{n+1}} x_n, \quad n \geq 1, \quad (3.15)$$

where $T^{\lambda_{n+1}} x_n := Tx_n - \lambda_{n+1} \mu F(Tx_n)$, $\mu > 0$, and $\{\alpha_n\}_{n=1}^\infty$ and $\{\lambda_n\}_{n=2}^\infty$ are real sequences in $[0, 1)$ satisfying the following conditions:

- (i) $0 < \alpha \leq \alpha_n < 1$ for all $n \geq 1$ and for some $\alpha \in (0, 1)$,
- (ii) $\sum_{n=1}^\infty (1 - \alpha_n) = \infty$,
- (iii) $\sum_{n=2}^\infty \lambda_n < \infty$.

Then, $\{x_n\}_{n=1}^\infty$ converges weakly to a fixed point of T .

Proof. From Lemma 2.1, $(I - T)$ is demiclosed at zero, and since $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ and E satisfies Opial's condition, it follows from standard argument that $\{x_n\}_{n=1}^\infty$ converges weakly to a fixed point of T . \square

Remark 3.3. It follows from Lemma 2.2 and Theorem 3.1 that under the hypothesis of Theorem 3.1, $\{x_n\}_{n=1}^\infty$ converges strongly to a fixed point p of T if and only if $\{x_n\}_{n=1}^\infty$ has a subsequence $\{x_{n_j}\}_{j=1}^\infty$ which converges strongly to p . Thus, under the hypothesis

of Theorem 3.1, if T is in addition completely continuous or demicompact, then $\{x_n\}_{n=1}^{\infty}$ converges strongly to a fixed point of T .

Furthermore, if T satisfies condition (A), then $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$; so under the conditions of Theorem 3.1, if T satisfies condition (A), then $\{x_n\}_{n=1}^{\infty}$ converges strongly to a fixed point of T .

Remark 3.4. Theorems 3.1 and 3.2 and Remark 3.3 extend the results of [1] from Hilbert spaces to much more general Banach spaces as considered here. Furthermore, the strong monotonicity condition imposed on F in [1] is not required in our results.

Prototypes of our real sequences $\{\alpha_n\}_{n=1}^{\infty}$ and $\{\lambda_n\}_{n=1}^{\infty}$ are $\alpha_n = n/(n+1), n \geq 1$ and $\lambda_n = 1/(n+1)^2, n \geq 1$.

References

- [1] L. Wang, “An iteration method for nonexpansive mappings in Hilbert spaces,” *Fixed Point Theory and Applications*, vol. 2007, Article ID 28619, 8 pages, 2007.
- [2] H. K. Xu and T. H. Kim, “Convergence of hybrid steepest-descent methods for variational inequalities,” *Journal of Optimization Theory and Applications*, vol. 119, no. 1, pp. 185–201, 2003.
- [3] I. Yamada, “The hybrid steepest descent method for the variational inequality problem over the intersection of fixed point sets of nonexpansive mappings,” in *Inherently Parallel Algorithms in Feasibility and Optimization and Their Applications (Haifa, 2000)*, D. Butnariu, Y. Censor, and S. Reich, Eds., vol. 8 of *Studies in Computational Mathematics*, pp. 473–504, North-Holland, Amsterdam, The Netherlands, 2001.
- [4] Z. Opial, “Weak convergence of the sequence of successive approximations for nonexpansive mappings,” *Bulletin of the American Mathematical Society*, vol. 73, pp. 591–597, 1967.
- [5] J. S. Jung, “Iterative approaches to common fixed points of nonexpansive mappings in Banach spaces,” *Journal of Mathematical Analysis and Applications*, vol. 302, no. 2, pp. 509–520, 2005.
- [6] M. O. Osilike and S. C. Aniagbosor, “Weak and strong convergence theorems for fixed points of asymptotically nonexpansive mappings,” *Mathematical and Computer Modelling*, vol. 32, no. 10, pp. 1181–1191, 2000.
- [7] M. O. Osilike, S. C. Aniagbosor, and B. G. Akuchu, “Fixed points of asymptotically demicontractive mappings in arbitrary Banach spaces,” *Panamerican Mathematical Journal*, vol. 12, no. 2, pp. 77–88, 2002.
- [8] L. Deng, “Convergence of the Ishikawa iteration process for nonexpansive mappings,” *Journal of Mathematical Analysis and Applications*, vol. 199, no. 3, pp. 769–775, 1996.

M. O. Osilike: Department of Mathematics, University of Nigeria, Nsukka, Nigeria
Email address: osilike@yahoo.com

F. O. Isiogugu: Department of Mathematics, University of Nigeria, Nsukka, Nigeria
Email address: obifeli2001@yahoo.com

P. U. Nwokoro: Department of Mathematics, University of Nigeria, Nsukka, Nigeria
Email address: petaun28@yahoo.com

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2009
First Round of Reviews	May 1, 2009
Publication Date	August 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Department of Physics, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk