

ITERATIVE METHODS FOR NONLINEAR QUASI COMPLEMENTARITY PROBLEMS

MUHAMMAD ASLAM NOOR

Mathematics Department, College of Science
King Saud University, Riyadh 11451, Saudi Arabia.

(Received April 13, 1985 and in revised form July 1, 1986)

ABSTRACT. In this paper, we consider and study an iterative algorithm for finding the approximate solution of the nonlinear quasi complementarity problem of finding $u \in K(u)$, such that

$$Tu \in K^*(u) \quad \text{and} \quad (u - m(u), Tu) = 0$$

where m is a point-to-point mapping, T is a (nonlinear) continuous mapping from a real Hilbert space H into itself and $K^*(u)$ is the polar cone of the convex cone $K(u)$ in H . We also discuss the convergence criteria and several special cases, which can be obtained from our main results.

KEY WORDS AND PHRASES. *Complementarity problems, Variational inequalities, Iterative methods, Convergence criteria.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES. 90C33, 49D20, 65K10.

1. INTRODUCTION.

The relationship between variational inequalities and complementarity is well known. Variational inequalities have been extended and generalized in various directions to study a wide class of difficult problems arising in different branches of mathematical and engineering sciences. Bensoussan and Lions [1] introduced a general class of variational inequalities known as quasi variational inequality, which has many applications in fluid dynamics, impulse control and general theory of equilibrium. In a quasi variational inequality formulation, the convex set also depends on the solution. Related to the variational inequality problem, there is also a complementarity problem, which has many applications in operations research, economics equilibrium and management sciences. In fact, it is known that if the underlying set in both these problems is a convex cone, then complementarity problem and variational inequality problem are equivalent. This equivalent has been used quite effectively in suggesting unified and general algorithms for solving complementarity problems, see Ahn [2] and Noor [3,4,5].

Motivated and inspired by the research activities in these fields, we introduce a new class of complementarity problems, which is the natural

generalization of linear and nonlinear complementarity problems studied extensively in recent years. After introducing the problem, we show that its solution can be obtained from an iterative scheme. We also discuss the convergence criteria for the approximate solution. Several special cases, which can be obtained from our main results, are discussed. Our results are an extension and improvements of the previously known results.

2. PRELIMINARIES AND BASIC RESULTS.

Let H be a real Hilbert space with its dual H' , whose inner product and norm are denoted by (\cdot, \cdot) and $\|\cdot\|$ respectively. We identify the Hilbert space H with its dual space H' . We define by \leq partial ordering on H , which is consistent with the topological vector structure of H that is the cone of all non-negative elements $\{v \in H : v \geq 0\}$ is a closed convex cone. Let K be a closed convex set in H . If $T:K \rightarrow H$ is a continuous mapping, then we consider the problem of finding $u \in K$ such that

$$(Tu, v-u) \geq 0, \quad \text{for all } v \in K. \quad (1.1)$$

Inequality of type (1.1) is known as variational inequality. If the convex set K also depends upon the solution, then variational inequality (1.1) is called quasi variational inequality. To be more specific, the quasi variational inequality problem is to find $u \in K(u)$ such that

$$(Tu, v-u) \geq 0, \quad \text{for all } v \in K(u). \quad (1.2)$$

In many important applications, $K(u)$ has the following form

$$K(u) = m(u) + K, \quad (1.3)$$

where m is a point-to-point mapping. In this case $K^*(u) = (m(u) + K)^* = m(u) \cap K^*$. For applications of quasi variational inequalities, see [6, 7, 8, 9].

Related to quasi variational inequalities, we now introduce a new class of complementarity problems, which will be called quasi complementarity problem of finding u such that

$$u - m(u) \geq 0, \quad Tu \geq 0, \quad (u - m(u), Tu) = 0 \quad (1.4)$$

If T is a nonlinear transformation, then problem (1.4) is called the nonlinear quasi complementarity problem (NQCP). If T is an affine transformation of the form $u \rightarrow Mu + q$, where M is a $n \times n$ matrix, q is a n -vector, then problem (1.4) is equivalent to finding u such that

$$u - m(u) \geq 0, \quad Mu + q \geq 0, \quad (u - m(u), Mu + q) = 0, \quad (1.5)$$

which is called linear quasi complementarity problem. Problem of type (1.5) was introduced and studied by Dolcetta [10], Mosco [7] and Pang [11, 12].

In particular, if the mapping m is zero, then problems (1.4) and (1.5) are equivalent to the following complementarity problems;

$$u \geq 0, \quad Tu \geq 0, \quad (u, Tu) = 0 \quad (1.6)$$

and

$$u \geq 0, \quad Mu + q \geq 0, \quad (u, Mu + q) = 0, \quad (1.7)$$

which have been extensively studied by Cottle [13,14], Karamardian [15], Mangasarian [16] and Ahn [17] in recent years.

Let $K^* = \{u \in H, (u, v) \geq 0 \text{ for all } v \in K\}$, be the polar cone of the convex cone K in H . If the convex cone also depends upon the solution itself, then we introduce the generalized quasi complementarity problem of finding $u \in K(u)$ such that

$$Tu \in K^*(u) \text{ and } (u - m(u), Tu) = 0, \quad (1.8)$$

It is clear that the nonlinear quasi complementarity problem (1.4) is a special case of the generalized nonlinear quasi complementarity problem (1.8). Furthermore, if the point-to-point mapping m is zero, then problem (1.8) is equivalent to finding $u \in K$ such that

$$Tu \in K^* \text{ and } (u, Tu) = 0,$$

which is known as the generalized nonlinear complementarity problem considered and studied by Karamardian [15], Cottle [14], Noor [3] and many others.

3. MAIN RESULTS.

We need the following results, the first is a generalization of a result of Karamardian [15] and Pang [11].

LEMMA 3.1. If K is the positive cone in H , then $u \in K(u)$, defined by (1.3), is a solution of (1.8) if and only if u satisfies the quasi variational inequality (1.2).

PROOF. The proof is similar to that of lemma 3.1 in [18,11].

THEOREM 3.1 [19,20]. For $K(u)$ given by (1.3), $u \in K(u)$ is a solution of quasi variational inequality (1.2) if and only if u satisfies the following relation

$$u = m(u) + P_K[u - \rho(Tu) - m(u)] \quad (3.1)$$

for some $\rho > 0$, where m is an arbitrary point-to-point mapping and P_K is the projection of H into K , the positive cone in H . For the description of P_K , see [8].

From lemma 3.1 and theorem 3.1, we conclude that the solution to problem (1.8) may be obtained by computing the fixed point of the function defined by (3.1). In fact, theorem 3.1 enables us to find the approximate solution u_n by the following iterative scheme:

$$u_{n+1} = m(u_n) + P_K[u_n - \rho T(u_n) - m(u_n)], \quad n=0,1,2,\dots \quad (3.2)$$

We also need the following concepts.

DEFINITION 3.1. A mapping $T: H \rightarrow H$ is said to be

i. *Strongly Monotone*, if there exists a constant $\alpha > 0$ such that

$$\langle Tu - Tv, u - v \rangle \geq \alpha \|u - v\|^2, \quad \text{for all } u, v \in H.$$

ii. *Lipschitz Continuous*, if there exists a constant $\beta > 0$ such that

$$\|Tu - Tv\| < \beta \|u - v\|, \quad \text{for all } u, v \in H.$$

In particular, it follows that $\alpha \leq \beta$.

In the next theorem, we study the conditions under which the approximate solution obtained from (3.2) converges to the exact solution u of quasi complementarity problem (QCP, 1.8).

THEOREM 3.2. Let the mapping T be strongly monotone and Lipschitz continuous. If the arbitrary mapping m is Lipschitz continuous, u_n and u are solutions satisfying (3.1) and (1.8), then u_n converges to u strongly in H , for

$$\left| \rho - \frac{\alpha}{\beta^2} \right| < \frac{\sqrt{\alpha^2 - 4\beta^2(\gamma - \gamma^2)}}{\beta^2}, \quad \alpha \geq 2\beta\sqrt{\gamma - \gamma^2} \quad \text{and } \gamma < 1/2,$$

where γ is the Lipschitz constant of m .

PROOF. The method of proof is similar to the one given in [5]. From lemma 3.1 and theorem 3.1, we observe that the solution u of (1.8) can be characterized by the relation (3.1). Hence from (3.1) and (3.2), we have

$$\begin{aligned} \|u_{n+1} - u\| &\leq \|m(u_n) - m(u) + P_K[u_n - \rho(Tu_n) - m(u_n)] - P_K[u - \rho(Tu) - m(u)]\| \\ &\leq \|m(u_n) - m(u)\| + \|u_n - u - \rho(Tu_n - Tu) - m(u_n) + m(u)\|, \\ &\quad \text{since } P_K \text{ is non-expansive, see [21,6].} \\ &\leq 2\gamma \|u_n - u\| + \|u_n - u - \rho(Tu_n - Tu)\| \end{aligned}$$

Now by the strongly monotonicity and Lipschitz continuity of T , we obtain

$$\|u_n - u - \rho(Tu_n - Tu)\|^2 \leq (1 - 2\rho\alpha + \beta^2\rho^2) \|u_n - u\|^2.$$

Thus using the above inequality, we have

$$\begin{aligned} \|u_{n+1} - u\| &\leq [\sqrt{(1 - 2\rho\alpha + \beta^2\rho^2) + 2\gamma}] \|u_n - u\| \\ &\leq \theta \|u_n - u\|, \end{aligned}$$

where

$$\theta = [\sqrt{(1 - 2\rho\alpha + \beta^2\rho^2)} + 2\gamma] < 1 \quad \text{for } \left| \rho - \frac{\alpha}{\beta^2} \right| < \frac{\sqrt{\alpha^2 - 4\beta^2(\gamma - \gamma^2)}}{\beta^2},$$

$$\gamma < \frac{1}{2} \text{ and } \alpha \geq 2\beta\sqrt{\gamma - \gamma^2}$$

Since $\theta < 1$, so the fixed point problem (3.1) has a unique solution u and consequently the Picard iterate u_{n+1} converges to u ; see [22], which is the required result.

It is clear that the use of algorithms as constructive methods for proving the existence of solution brings theory and computation closer together. Theorem 3.2 enables us to find the approximate solution of (1.8) by the following iterative scheme:

i. Given $u_0 \in H$.

ii. $u_{n+1} = m(u_n) + P_K[u_n - \rho(Tu_n) - m(u_n)], \quad n = 0, 1, 2, 3, \dots$

where P_K is the projection of H into K , m is a point-to-point mapping and $\rho > 0$ is required to satisfy the following conditions:

$$|\rho - \frac{\alpha}{\beta^2}| < \frac{\sqrt{\alpha^2 - 4\beta^2(\gamma - \gamma^2)}}{\beta^2}, \gamma < \frac{1}{2} \text{ and } \alpha > 2\beta\sqrt{\gamma - \gamma^2}.$$

REMARK 3.1. If the point-to-point mapping is zero, then it implies that the Lipschitz constant γ is zero. Thus our algorithm is exactly the same as considered by Fang [23] and Gana [24]. Furthermore, theorem 3.1 reduces to a result of Noor and Noor [21, 25] for the corresponding variational inequalities. When T is of the form $Tu = Mu + q$, one can easily prove that T is strongly monotone if and only if M is positive definite, see [15] for proof. Obviously T is also Lipschitz continuous. For the linear quasi complementarity problem, see [18] where one can find a general algorithm for finding the approximate solution along with the necessary convergence criteria.

ACKNOWLEDGEMENT. I wish to express my gratitude to Prof. Dr. W. Oettli for his valuable suggestions and comments.

REFERENCES

1. BENOUSSAN, A. and LIONS, J. *Applications des inéquations variationnelles en contrôle stochastique*, Dunod, Paris, 1978.
2. AHN, B.H. Iterative methods for linear complementarity problems with upper bounds on primary variables, *Math. Programming*, 26(1983), 295-315.
3. NOOR, M. ASLAM. Generalized nonlinear complementarity problem, *J. Nat. Sci. & Math.*, 26(1986), 9-20.
4. NOOR M. ASLAM. On the nonlinear complementarity problem, *J. Math. Anal. Appl.*, to appear.
5. NOOR, M. ASLAM. Generalized quasi complementarity problems, *J. Math. Anal. Appl.* (1986), to appear.
6. BAIOCCHI, C. and CAPELO, A. *Variational and quasi variational inequalities*, John Wiley and Sons, New York, 1984.
7. MOSCO, U. Implicit variational problems and quasi variational inequalities, *Lect. Notes Math.*, Springer-Verlag, Berlin, 1976.
8. ODEN, J. and KIKUCHI, N. Theory of variational inequalities with applications to problems of flow through porous media, *Int. J. Engng. Science*, 18(1986), 1173-1284.
9. CRANK, J. *Free and moving boundary problems*, Calendron Press, London, 1984.
10. DOLCETTA, I. *Systemi di complementarità a deseguaglianze variazionali*, Ph.D. Thesis, University of Rome, Italy, 1972.
11. PANG, J.S. On the convergence of a basic iterative method for the implicit complementarity problem, *J. Opt. Theor. Appl.* 37(1982), 149-162.
12. PANG J.S. *The implicit complementarity problem in Nonlinear Programming 4*, (ed.) by Mangasarian, Meyer and Robinson, Academic Press, London, (1981), 487-518.
13. COTTLE, R. Complementarity and variational problems, *Symposia Math.* 19(1976), 177-208.

14. COTTLE, R. Note on a fundamental theorem in quadratic programming, SIAM J. Appl. Math., 12(1964), 663-665.
15. KARAMARDIAN, S. Generalized complementarity problem, J. Opt. Theor. Appl. 8(1971), 161-168.
16. MANGASARIAN, O.L. Solution of symmetric linear complementarity problem by iterative methods, J. Opt. Theor. Appl. 22(1976), 465-485.
17. AHN B.H. Solution of non symmetric linear complementarity problems by iterative methods, J. Opt. Theor. Appl. 33(1981), 175-185.
18. NOOR, M. ASLAM. Convergence analysis of the iterative methods for quasi complementarity problems, to appear.
19. NOOR, M. ASLAM. An iterative scheme for a class of quasi variational inequalities, J. Math. Anal. Appl. 110(1985), 463-468.
20. CHAN, P. and PANG, J.S. The generalized quasi variational inequality problem, Math. Op. Res. 7(1982), 221-222.
21. NOOR, M. ASLAM and NOOR, K. INAYAT. Iterative methods for variational inequalities and nonlinear programming, Methods Op. Res. 31(1979), 455-463.
22. ORTEGA, J.M. and RHEINBOLD, W. Iterative solutions of nonlinear equations in several variables, Academic Press, London, 1970.
23. FANG, S.C. An iterative method for generalized complementarity problems, IEEE Trans. Aut. Control, 25(1980), 1225-1227.
24. GANA, A. Studies in complementarity problems, Ph.D. thesis, University of Michigan, U.S.A., 1982.
25. NOOR, K. INAYAT and NOOR, M. ASLAM. Iterative methods for a class of variational inequalities in Numerical Analysis of Singular Perturbation Problems, (ed.) by Hemker and Miller, Academic Press, London, (1979), 441-448.

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk