

## HOMOMORPHISMS AND RELATED CONTRACTIONS OF GRAPHS

ROBERT D. GIRSE and RICHARD A. GILLMAN

Department of Mathematics  
Idaho State University  
Pocatello, ID 83209

(Received August 6, 1985 and in revised form July 16, 1986)

**ABSTRACT.** For every homomorphism  $\phi$  of a graph  $G$  there exists a contraction  $\theta_\phi$  on  $\bar{G}$ , the complement of  $G$ . Here we study the graph equation  $\phi(G) = \theta_\phi(\bar{G})$ . In the course of our work we show that Hadwiger's Conjecture is true for every self-complementary graph.

**KEYS WORDS AND PHRASES.** Homomorphisms of graphs, contractions of graphs, self-complementary graph.

1980 AMS SUBJECT CLASSIFICATION CODE. 05C15.

### 1. INTRODUCTION.

By a graph of order  $n$  we mean a set  $V(G)$  of  $n$  vertices together with a set  $E(G)$  of unordered pairs of distinct vertices in  $V(G)$  called edges. A graph  $G$  is isomorphic to a graph  $H$  if there is a bijection from  $V(G)$  onto  $V(H)$  which preserves both adjacency and non-adjacency, in which case we will write  $G = H$ . A graph with the property that  $G = \bar{G}$ , where  $\bar{G}$  denotes the complement of  $G$ , is called self-complementary.

An elementary homomorphism of a graph  $G$  is the identification of two non-adjacent vertices of  $G$ , and a homomorphism is a sequence of elementary homomorphisms. Thus a homomorphism of a graph  $G$  onto a graph  $H$  is a function  $\phi$  from  $V(G)$  onto  $V(H)$  such that whenever  $u$  and  $v$  are adjacent in  $G$ ,  $\phi(u)$  and  $\phi(v)$  are adjacent in  $H$ . Likewise, an elementary contraction of a graph  $G$  is the identification of two adjacent vertices of  $G$ , and a contraction is a sequence of elementary contractions. Thus for every homomorphism  $\phi$  of  $G$  there is a contraction  $\theta_\phi$  of  $\bar{G}$  that we may construct as follows:  $\phi$  is a sequence of elementary homomorphisms  $\epsilon_1, \epsilon_2, \dots, \epsilon_n$  each of which identifies two non-adjacent vertices in  $G$ , so we let  $\theta_\phi$  be the sequence of elementary contractions  $\theta_1, \theta_2, \dots, \theta_n$  such that  $\theta_i$  identifies the same vertices in  $\bar{G}$  that  $\epsilon_i$  identifies in  $G$ .

Recently [1] the graph equation  $\overline{\phi(G)} = \theta_\phi(\bar{G})$  was studied. Here we consider a similar equation, namely,

$$\phi(G) = \theta_\phi(\bar{G}) . \quad (1.1)$$

We will employ the following notation as the need arises:  $\rho_G(u)$  will denote the valency of the vertex  $u$  in  $G$  and  $A_G(u)$  will be the set of all vertices in  $G$  that are

adjacent to  $u$ . Thus  $\rho_G(u) = |A_G(u)|$ , where  $|A_G(u)|$  is the cardinality of the set  $A_G(u)$ . As usual  $\chi(G)$  will denote the chromatic number of  $G$ .

## 2. SOME GENERAL RESULTS.

THEOREM 1. There is no graph  $G$  of order  $n > 1$  such that (1.1) is satisfied for every homomorphism  $\phi$  of  $G$ .

If  $G$  is not self-complementary then the identity homomorphism and its related null contraction suffice to satisfy the theorem. We will postpone the remainder of the proof until section 3, where we restrict our attention to results on self-complementary graphs.

THEOREM 2. If there exists a homomorphism  $\phi$  of  $G$  that satisfies (1.1) then:

- (a)  $\bar{G}$  must be connected.
- (b)  $\bar{G}$  cannot be a tree of order  $n \geq 5$ .

PROOF. (a) If  $\bar{G}$  is not connected, then no contraction of  $\bar{G}$  is connected.

However,  $\bar{G}$  not connected implies  $G$  is connected, thus every homomorphic image of  $G$  will be connected.

(b) If  $\bar{G}$  is a tree of order  $n \geq 5$  then  $G$  and every homomorphic image of  $G$  contains  $K_3$ , the complete graph on three vertices, as a subgraph. But every contraction of  $\bar{G}$  will be a tree and so cannot contain  $K_3$  as a subgraph.

THEOREM 3. If  $\chi(G) = 2$  and  $\bar{G}$  is connected then there exists a homomorphism  $\phi$  such that  $\phi(G) = K_2 = \theta_\phi(\bar{G})$ .

PROOF. Since  $\bar{G}$  is connected, the image of  $\bar{G}$  under any contraction will be connected. Thus every contraction of  $\bar{G}$  onto two points must have  $K_2$  as its image. From [3] we know that there exists a homomorphism  $\phi$  of  $G$  such that  $\phi(G) = K_{\chi(G)}$ . Hence, using this homomorphism and its related contraction, we have  $\phi(G) = K_2 = \theta_\phi(\bar{G})$ .

Since every homomorphism is a sequence of elementary homomorphisms, we now turn our attention to the equation

$$\varepsilon(G) = \theta_\varepsilon(\bar{G}), \quad (2.1)$$

where  $\varepsilon$  is an elementary homomorphism.

THEOREM 4. If  $G$  satisfies (2.1) then

$$\chi(\bar{G}) - 2 \leq \chi(G) \leq \chi(\bar{G}) + 1.$$

PROOF. In [3] Harary et. al. proved the following inequalities:

$$\chi(G) \leq \chi(\varepsilon(G)) \leq \chi(G) + 1$$

$$\chi(\bar{G}) - 1 \leq \chi(\theta_\varepsilon(\bar{G})) \leq \chi(\bar{G}) + 1.$$

Since (2.1) holds we have  $\chi(\varepsilon(G)) = \chi(\theta_\varepsilon(\bar{G}))$ , and so from the first inequality  $\chi(\theta_\varepsilon(\bar{G}))$  equals either  $\chi(G)$  or  $\chi(G) + 1$ . Putting these values into the second inequality above yields:

$$\chi(\bar{G}) - 1 \leq \chi(G) \leq \chi(\bar{G}) + 1$$

$$\chi(\bar{G}) - 2 \leq \chi(G) \leq \chi(\bar{G}),$$

which completes the proof.

The following result will be needed in the next section.

LEMMA 1. Let  $G$  be of order  $n$  and  $\epsilon$  an elementary homomorphism that identifies  $u_1$  and  $u_2$  in  $G$ . If (2.1) holds then

$$|A_G(u_1) \cap A_G(u_2)| = |A_{\bar{G}}(u_1) \cap A_{\bar{G}}(u_2)| + 1 + 2|E(G)| - \frac{n(n-1)}{2} .$$

PROOF. If  $\epsilon$  identifies  $u_1$  and  $u_2$  in  $G$  we have  $|E(\epsilon(G))| = |E(G)| - |A_{\bar{G}}(u_1) \cap A_{\bar{G}}(u_2)|$ , since when an elementary homomorphism  $\epsilon$  is applied to a graph  $G$ ,  $\epsilon(G)$  will lose one edge for each vertex which is adjacent to both vertices that were identified, and all other edges will remain in  $\epsilon(G)$ . Likewise, by the same reasoning as above,  $|E(\theta_{\epsilon}(\bar{G}))| = |E(\bar{G})| - |A_{\bar{G}}(u_1) \cap A_{\bar{G}}(u_2)| - 1$ , where the extra edge removed is the one which was contracted. Now if (2.1) holds both  $\epsilon(G)$  and  $\theta_{\epsilon}(\bar{G})$  must contain the same number of edges. Thus

$$|E(G)| - |A_G(u_1) \cap A_G(u_2)| = |E(\bar{G})| - |A_{\bar{G}}(u_1) \cap A_{\bar{G}}(u_2)| - 1,$$

where  $|E(\bar{G})| = \frac{n(n-1)}{2} - |E(G)|$ , which proves the lemma.

### 3. SELF-COMPLEMENTARY GRAPHS.

We first summarize some results from [4], [5] and [6] that will be needed. Let  $S$  be a self-complementary graph. If  $S$  is of order  $n$  then  $n \equiv 0, 1 \pmod{4}$ . Suppose  $f$  is an isomorphism such that  $f(S) = \bar{S}$ . If  $n \equiv 0 \pmod{4}$  then  $f$  has no fixed vertex, that is  $f(u) \neq u$  for all  $u \in V(S)$ . However if  $n \equiv 1 \pmod{4}$  there exists precisely one fixed vertex for  $f$ , and moreover any such  $S$  can be constructed by appropriately adding this fixed vertex, of valency  $\frac{n-1}{2}$ , to a self-complementary graph of order  $n-1$ . Lastly there are regular self-complementary graphs of degree  $\frac{n-1}{2}$  if and only if  $n \equiv 1 \pmod{4}$ .

We will assume throughout this section that  $S$  is nontrivial, that is  $n > 1$ . The following sequence of four lemmas will complete the proof of Theorem 1.

LEMMA 2. If  $S$  is of order  $n \equiv 0 \pmod{4}$  then there exist vertices  $u_1, u_2 \in V(S)$  such that  $u_1 u_2 \notin E(S)$  and  $\rho_S(u_1) = \rho_S(u_2)$ .

PROOF. Since  $n \equiv 0 \pmod{4}$ , any given valency that occurs in  $S$  will occur an even number of times, [4]. Thus there are vertices  $u_1, u_2 \in V(S)$  such that  $\rho_S(u_1) = \rho_S(u_2)$ . Now let  $f(S) = \bar{S}$ , then  $\rho_S(u_1) = \rho_S(u_2)$  implies  $\rho_{\bar{S}}(f(u_1)) = \rho_{\bar{S}}(f(u_2))$  and also  $\rho_{\bar{S}}(f(u_1)) = \rho_{\bar{S}}(f(u_2))$ . But  $u_1 u_2 \in E(S)$  if and only if  $f(u_1) f(u_2) \notin E(\bar{S})$  and so  $u_1 u_2$  or  $f(u_1) f(u_2)$  satisfy the lemma.

REMARK. For any self-complementary graph  $S$  of order  $n$ ,  $|E(S)| = \frac{n(n-1)}{4}$ . Thus from Lemma 1, if  $\epsilon$  is an elementary homomorphism of  $S$  which identifies the vertices  $u_1$  and  $u_2$  and  $\epsilon(S) = \theta_{\epsilon}(\bar{S})$ , then  $|A_S(u_1) \cap A_S(u_2)| = |A_{\bar{S}}(u_1) \cap A_{\bar{S}}(u_2)| + 1$ .

LEMMA 3. If  $S$  is of order  $n \equiv 0 \pmod{4}$  then there is an elementary homomorphism  $\epsilon$  of  $S$  such that  $\epsilon(S) \neq \theta_{\epsilon}(\bar{S})$ .

PROOF. Using Lemma 2 choose  $u_1, u_2 \in V(S)$  such that  $u_1 u_2 \notin E(S)$  but  $\rho_S(u_1) = \rho_S(u_2)$ . Now for any  $u \in V(S)$  that is distinct from  $u_1$  and  $u_2$ ,  $u$  can be adjacent to both  $u_1$  and  $u_2$ , neither  $u_1$  nor  $u_2$ , or adjacent to one and not the other.

Thus we have

$$A_s(u_1) = \{u_2, \dots, a_{k_1}, b_1, \dots, b_\ell\} \quad A_s(u_2) = \{a_1, \dots, a_{k_1}, c_1, \dots, c_\ell\}$$

$$A_{\bar{s}}(u_1) = \{u_2, d_1, \dots, d_{k_2}, c_1, \dots, c_\ell\} \quad A_{\bar{s}}(u_2) = \{u_1, d_1, \dots, d_{k_2}, b_1, \dots, b_\ell\}$$

where  $k_1 = |A_s(u_1) \cap A_s(u_2)|$  and  $k_2 = |A_{\bar{s}}(u_1) \cap A_{\bar{s}}(u_2)|$ . Let  $\varepsilon$  identify  $u_1$  and  $u_2$  and suppose  $\varepsilon(S) = \theta_\varepsilon(\bar{S})$ , so that from the preceding Remark  $k_1 = k_2 + 1$ . Then

$$n-1 = \rho_s(u_1) + \rho_{\bar{s}}(u_1) = (k_1 + \ell) + (k_2 + 1 + \ell) = 2k_1 + 2\ell.$$

Hence  $n = 2k_1 + 2\ell + 1 \not\equiv 0 \pmod{4}$ , a contradiction.

We now consider those self-complementary graphs of order  $n \equiv 1 \pmod{4}$  and let  $v$  be the fixed vertex, of valency  $\frac{n-1}{2}$ , under the isomorphism  $f(S) = \bar{S}$ .

LEMMA 4. Let  $S$  be of order  $n \equiv 1 \pmod{4}$  and  $\varepsilon$  be an elementary homomorphism that identifies any  $u \in V(S)$  with  $v$ . If  $\varepsilon(S) = \theta_\varepsilon(\bar{S})$  then  $S$  is regular.

PROOF. Let  $f(S) = \bar{S}$ , then for any  $u' \in V(S)$  distinct from  $v$ ,  $v u' \in E(S)$  if and only if  $v f(u') \notin E(S)$  while  $\rho_s(u') = \rho_{\bar{s}}(f(u'))$ .

Suppose  $S$  is not regular, then there is a vertex  $u \in V(S)$  such that  $\rho_s(u) \neq \frac{n-1}{2}$  and, from the observation above,  $v u \notin E(S)$ . Now let  $\varepsilon$  identify this vertex  $u$  and  $v$  and suppose  $\varepsilon(S) = \theta_\varepsilon(\bar{S})$ . We have, as in the proof of Lemma 3,

$$A_s(v) = \{a_1, \dots, a_{k_1}, b_1, \dots, b_{\frac{n-1}{2} - k_1}\} \quad A_s(u) = \{a_1, \dots, a_{k_1}, c_1, \dots, c_\ell\}$$

$$A_{\bar{s}}(v) = \{u, d_1, \dots, d_{k_2}, c_1, \dots, c_\ell\} \quad A_{\bar{s}}(u) = \{v, d_1, \dots, d_{k_2}, b_1, \dots, b_{\frac{n-1}{2} - k_1}\}$$

where  $k_1 = |A_s(u) \cap A_s(v)|$  and  $k_2 = |A_{\bar{s}}(u) \cap A_{\bar{s}}(v)|$ . But  $\rho_s(v) = \frac{n-1}{2}$  and so

$\ell = \frac{n-1}{2} - (k_2 + 1)$ , and  $k_1 = k_2 + 1$  since  $\varepsilon(S) = \theta_\varepsilon(\bar{S})$ . Hence

$$\rho_s(u) = k_1 + \ell = k_1 + \frac{n-1}{2} - (k_2 + 1) = \frac{n-1}{2}, \text{ a contradiction.}$$

LEMMA 5. If  $S$  is a regular self-complementary graph then there is an elementary homomorphism  $\varepsilon$  such that  $\varepsilon(S) \neq \theta_\varepsilon(\bar{S})$

PROOF. First note that  $S v$  is self-complementary, with  $\frac{n-1}{2}$  vertices of valency  $\frac{n-1}{2}$  and  $\frac{n-1}{2}$  vertices of valency  $\frac{n-3}{2}$ . Thus there are  $\frac{n-1}{2}$  edges joining every vertex of valency  $\frac{n-1}{2}$  to the vertices of valency  $\frac{n-3}{2}$ , and these must appear in either  $S v$  or  $\bar{S} v$ . Since  $S v = \bar{S} v$  these edges must be equally divided between  $S v$  and its complement. When  $v$  is added to  $S v$  to form  $S$  it is adjacent to every vertex of valency  $\frac{n-3}{2}$ . Thus if  $u \in V(S v)$  such that  $\rho_{S v}(u) = \frac{n-1}{2}$ , we have

$$|A_s(u) \cap A_s(v)| = \frac{n-1}{4}.$$

Now  $S$  is regular of degree  $\frac{n-1}{2}$  and we let  $\varepsilon$  identify  $u$ , as described above, and  $v$ . Assume  $\varepsilon(S) = \theta_\varepsilon(\bar{S})$  and  $u' \in V(\varepsilon(S))$  is the image of  $u$  and  $v$  under  $\varepsilon$ . Then

$$\begin{aligned} \rho_{\varepsilon(S)}(u') &= \rho_s(u) + \rho_{\bar{s}}(v) - |A_s(u_1) \cap A_s(u_2)| \\ &= \frac{n-1}{2} + \frac{n-1}{2} - \frac{n-1}{4} = 3(\frac{n-1}{4}), \end{aligned}$$

since every vertex adjacent to both  $u$  and  $v$  in  $S$  will only account for one edge incident to  $u'$  in  $\epsilon(S)$ . Similarly

$$\rho_{\theta_{\epsilon}(\bar{S})}(u') = \rho_{\bar{S}}(u) + \rho_{\bar{S}}(v) - |A_{\bar{S}}(u) \cap A_{\bar{S}}(v)| - 2,$$

where the 2 is subtracted since the edge contracted under  $\theta_{\epsilon}$  is incident to both  $u$  and  $v$ . However since  $\epsilon(S) = \theta_{\epsilon}(\bar{S})$  we must have  $|A_{\bar{S}}(u) \cap A_{\bar{S}}(v)| = |A_{\bar{S}}(u) \cap A_{\bar{S}}(v)| + 1$ , and so

$$\rho_{\theta_{\epsilon}(\bar{S})}(u') = \frac{n-1}{2} + \frac{n-1}{2} - \frac{n-1}{4} - 1 = 3(\frac{n-1}{4}) - 1.$$

Every other  $\bar{u} \in V(\theta_{\epsilon}(\bar{S}))$  has valency  $\rho_{\bar{S}}(\bar{u}) - 1 = \frac{n-3}{2}$  or  $\rho_{\bar{S}}(\bar{u}) = \frac{n-1}{2}$  depending on whether or not  $\bar{u} \in A_{\bar{S}}(u) \cap A_{\bar{S}}(v)$ . Therefore  $\rho_{\epsilon(S)}(u') > \rho_{\theta_{\epsilon}(\bar{S})}(\bar{u})$  for every vertex  $\bar{u}$  in  $\theta_{\epsilon}(\bar{S})$ , and hence  $\epsilon(S) \neq \theta_{\epsilon}(\bar{S})$  since isomorphisms must preserve valencies.

Now that the proof of Theorem 1 is complete we will show that for every self-complementary graph there is a homomorphism  $\phi$  which satisfies (1.1).

LEMMA 6. Let  $S$  be of order  $n$  and  $f(S) = \bar{S}$ . Then in the set

$V = \{(u_i, f(u_i)) \mid u_i \in V(S), i=1, \dots, n\}$  precisely  $m = \lceil \frac{n}{2} \rceil$  are non-adjacent pairs of distinct vertices in  $S$ .

PROOF. Let  $u \in V(S)$  such that  $f(u) \neq u$ . Then  $u$  and  $f(u)$  are adjacent in  $S$  if and only if  $f(u)$  and  $f(f(u))$  are non-adjacent in  $S$ . Now  $f$  is an isomorphism, and so  $f(u)$  and  $f(f(u))$  must also be distinct vertices of  $S$ . Hence for every pair of distinct non-adjacent vertices in  $V$ , there is a pair of adjacent vertices in  $V$  and conversely. Thus there must be  $m = \lceil \frac{n}{2} \rceil$  pairs of distinct non-adjacent vertices in  $V$ .

THEOREM 5. Let  $S$  be of order  $n$  and  $m = \lceil \frac{n}{2} \rceil$ . There exists a homomorphism  $\phi$  of  $S$  such that  $\phi(S) = K_{n-m} = \theta_{\phi}(\bar{S})$ .

PROOF. For each of the  $m$  pairs of distinct non-adjacent vertices provided by Lemma 6, there exists an elementary homomorphism  $\epsilon_i$  of  $S$  that identifies  $u_i$  and  $f(u_i)$ . Let  $\phi = \epsilon_1, \epsilon_2, \dots, \epsilon_m$  so that both  $\phi(S)$  and  $\theta_{\phi}(\bar{S})$  will have  $n-m$  vertices.

Suppose  $\phi(S) \neq K_{n-m}$ , then there are vertices  $u_1, u_2 \in V(\phi(S))$  such that  $u_1 u_2 \notin E(\phi(S))$ . Let  $\phi^{-1}(u_1) = \{a_1, a_2\}$  and  $\phi^{-1}(u_2) = \{b_1, b_2\}$ . Thus  $a_1 \neq b_1$ ,  $f(a_1) = a_2$  and  $f(b_1) = b_2$  where  $f(S) = \bar{S}$ . Now  $u_1 u_2 \notin E(\phi(S))$  implies  $a_1 b_1 \notin E(S)$  and so  $f(a_1) f(b_1) \notin E(\bar{S})$ . But then  $f(a_1) f(b_1) = a_1 b_2 \in E(S)$  and so  $u_1 u_2 \in E(\phi(S))$ , a contradiction. Thus  $\phi(S) = K_{n-m}$ . By an argument virtually identical to the one just given it follows that  $\theta_{\phi}(\bar{S}) = K_{n-m}$ , which proves the theorem.

From [3] we know that the smallest homomorphic image of any graph  $G$  is a complete graph of order  $\chi(G)$ . Thus from Theorem 5 we have  $\chi(S) \leq n - \lceil \frac{n}{2} \rceil$  for every self-complementary graph  $S$  of order  $n$ . However if a graph is contractable to a complete graph of order  $t$ , then it has a complete contraction of order  $k$  for  $1 \leq k \leq t$ , also from [3]. Hence Theorem 5 shows that every self-complementary graph satisfies Hadwiger's Conjecture [2], that is, every self-complementary graph  $S$  has a complete contraction of order  $\chi(S)$ .

#### 4. CONCLUDING REMARKS.

It would certainly be desirable to find some simple necessary and sufficient

conditions on the graph  $G$ , and perhaps on  $\bar{G}$ , to ensure the existence of a homomorphism  $\phi$  such that the graph equation  $\phi(G) = \theta_{\phi}(\bar{G})$  holds. Given that such conditions can be found, we would like to be able to construct the appropriate homomorphisms for which the equation holds or perhaps enumerate those homomorphisms of  $G$  that satisfy the equation. All of these problems remain open.

## REFERENCES

1. GIRSE, R. Homomorphisms of Complete  $n$ -Partite Graphs, Internat. J. Math. Math. Sci., to appear.
2. HADWIGER, H. Über eine Klassifikation der Streckenkomplexe, Vierteljschr. Naturforsch. Ges. Zurich 88 (1943), 133-142.
3. HARARY, F. HEDETNIEMI, S. and PRINS, G. An Interpolation Theorem for Graphical Homomorphisms, Portugal Math. 26 (1969), 453-462.
4. MORRIS, P. Self-Complementary Graphs and Digraphs, Math. Comp. 27 (1973), 216-217.
5. RINGEL, G. Selbstkomplementare Graphen, Archiv d. Math. 14 (1963), 354-358.
6. SACHS, H. Über Selbstkomplementare Graphen, Publ. Math. Debrecen 9 (1962), 270-288.

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)