

A MAXIMAL CHAIN APPROACH TO TOPOLOGY AND ORDER

R. VAINIO

Department of mathematics
Åbo Akademi
SF-20500 Åbo, Finland

(Received May 18, 1987 and in revised form August 26, 1987)

Abstract. On ordered sets (posets, lattices) we regard topologies (or, more general convergence structures) which on any maximal chain of the ordered set induce its own interval topology. This construction generalizes several well-known intrinsic structures, and still contains enough to produce interesting results on for instance compactness and connectedness. The "maximal chain compatibility" between topology (convergence structure) and order is preserved by formation of arbitrary products, at least in case the involved order structures are conditionally complete lattices.

AMS Subject Classification Codes: Primary 54A20, Secondary 54F05.

Key words and phrases: Intrinsic topologies (convergence structures) on ordered sets, order convergence, maximal chains.

INTRODUCTION

Parts of §§2 and 3 of the present note can be regarded as an extension of Vainio [9]. Terminology is as in [9]; crucial definitions and notations are recapitulated in next section. Let P be a partially ordered set (short: poset), endow P with a topology (or, more generally: a convergence structure) q such that *all maximal chains of P inherit their own interval topologies*. The resulting space (P, q) is called an *i-space*, and the structure q an *i-structure* (*i-topology*, or *i-convergence*) on P .

Specially on lattices, *i*-space compatibility provides an excellent realm for study of i.a. connectedness and compactness (§3). Partly, this is because completeness (conditional completeness) can be equivalently described as completeness (conditional completeness) of all maximal chains.

Examples of *i*-spaces abound. Whenever a poset is endowed with a topology, which is finer than interval topology and for which all maximal chains are compact sets, *i*-space compatibility follows. Every poset admits a *finest i-convergence*, as well as a *finest i-topology*. Several well-known intrinsic convergences are *i*-structures, as demonstrated in §2.

For conditionally complete lattices, an arbitrary product of *i*-spaces is again an *i*-space (§4). This is of interest, because intrinsic topologies (convergences) of ordered structures do not behave well visavi formation of products (cf. Erné [1]). In §4 a few category theoretical remarks on *i*-convergences are included.

The use of convergence structures instead of (the more special) topologies is motivated by works of i.a. M. Erné and D. C. Kent, which show a filter theoretical approach to intrinsic topologies on ordered sets provides an elegant and powerful method. Several important and natural structures, such as *order convergence* (cf. §1) are not topological. Moreover, R. N. Ball has created completion theory for lattices using *Cauchy structures* (i.e. Cauchy filters of uniform convergence structures) — still another example where classical topology does not suffice.

1. PRELIMINARIES

A *convergence (structure)* on a set S is a map $q : S \rightarrow 2^{F(S)}$ ($F(S) \equiv$ the set of all proper filters on S), which for all $x \in S$ satisfies

- (1) $[x] \in q(x)$. ($[x]$ is the trivial ultrafilter derived from x .)
- (2) $\mathcal{F} \in q(x)$ and $\mathcal{G} \supseteq \mathcal{F} \Rightarrow \mathcal{G} \in q(x)$.
- (3) $\mathcal{F} \in q(x) \Rightarrow \mathcal{F} \cap [x] \in q(x)$.

Since we wish *order convergence* (see below) to constitute a special example of our theory, we can not assume $\mathcal{F}, \mathcal{G} \in q(x) \Rightarrow \mathcal{F} \cap \mathcal{G} \in q(x)$. The definition of convergence as stated in axioms (1), (2), (3) above dates back to Kent [5]. The couple (S, q) is called a *convergence space*.

For $A \subseteq S$, let q_A denote the convergence q inherited to A . Of course, all topologies are convergences. The *topological modification* $\text{top}(q)$ is the finest topology on S coarser than q . The category of convergence spaces (morphisms: continuous maps) is *cartesian closed*, a fact we will apply in §4. In Gähler [3] convergence spaces are treated in considerable depth. Following [3] and Vainio [9] a convergence space (S, q) is called *connected*, if the topological modification $(S, \text{top}(q))$ is, i.e. if all continuous maps from (S, q) to the two-point discrete space are constant maps. A set in a given convergence space is a *connected set*, if the corresponding subspace is a connected convergence space.

Partially ordered sets (posets) will be denoted by P , lattices by L . Quasi-ordered sets (no anti-symmetry assumed!) will be used in §4, primarily as tools. Totally ordered subsets of P are called *chains*; a chain which is not a proper subset of any other chain of P is called a *maximal chain* in P . For $A \subseteq P$, $A^*(A^+)$ is the set of all upper bounds (lower bounds) of A . If $A = \{a\}$, we write $a^*(a^+)$. A poset is *order dense*, if $[x, y] \neq \emptyset$ for all $x < y$. Definitions of *complete* (*conditionally complete*) lattices and *subcomplete* (*conditionally subcomplete*) sublattices are the standard ones. The *lattice translations* $L \rightarrow L$ are the maps $x \rightarrow a \vee x, x \rightarrow a \wedge x$, for any $a \in L$. Any chain J in L is, of course, a lattice in its own right, and for $S \subseteq J$ the indices in the expressions $\vee_J S, \vee_L S$ tell in which lattice the l.u.b. is formed, provided it exists (dual notation for g.l.b.).

A poset endowed with a convergence is called *T_1 -ordered*, if all sets a^*, a^+ are closed. We will use several intrinsic convergence structures on a given poset P , i.a. *interval topology* $t(P)$ (O. Frink) and *order convergence* $o(P)$ (D. C. Kent). The former is defined as the coarsest T_1 -ordered topology on P , and the latter is given by

$$\mathcal{F} \in o(P)(x) \Leftrightarrow \wedge \mathcal{F}^* \text{ and } \vee \mathcal{F}^+ \text{ both exist, and both equal } x.$$

Hereby,

$$\mathcal{F}^* = \cup\{F^* : F \in \mathcal{F}\} \text{ and } \mathcal{F}^+ = \cup\{F^+ : F \in \mathcal{F}\}.$$

In any lattice L , the classical *order topology* (G. Birkhoff) equals $\text{top}(o(L))$.

We are now ready to proceed to ordered topological spaces and their maximal chains.

2. DEFINITION AND EXAMPLES OF i -SPACES

Let q be a convergence on an arbitrary poset P . The pair (P, q) is an *i -space* (q is an *i -convergence*, or an *i -topology* on P) , if for any maximal chain J of P the equality $t(J) = q_J$ holds. Since interval topologies on chains are Hausdorff (even T_5), any *i -space* (P, q) satisfies

$$x \leq y \text{ and } \mathcal{F} \in q(x) \cap q(y) \Rightarrow \mathcal{F} \text{ has no trace to any chain containing } x \text{ and } y.$$

We also note any T_1 -ordered space (P, q) is an *i -space*, if and only if $t(J) \geq q_J$ for all maximal chains J of P . However, there are *i -spaces* which are not T_1 -ordered. Clearly, on any poset all T_1 -ordered topologies yielding compact maximal chains are automatically *i -topologies*.

EXAMPLE 1. Every poset P admits a *finest i -convergence* $s(P)$, which determines *Rennie's chain topology* $r(P)$. (A set S in P is $r(P)$ -open, if and only if $S \cap J$ is a $t(J)$ -open set for all maximal chains J . This structure was referred to by Rennie [7].) Of course, $r(P)$ is the *finest i -topology on P* . In general, there is no coarsest *i -convergence* or coarsest *i -topology* on a given poset. Indeed, let \mathcal{S}_1 (\mathcal{S}_2) be the set of all open angular regions in the plane \mathbf{R}^2 with vertex at $(1,0)$ (at $(0,1)$). The restriction of \mathcal{S}_1 (\mathcal{S}_2) to $]0, 1[\times]0, 1[$ determines the topology τ_1 (τ_2) on $]0, 1[\times]0, 1[$. Now, both τ_1 and τ_2 are *i -topologies* on the poset $]0, 1[\times]0, 1[$ (endowed with natural order), but neither their g.l.b. topology nor their g.l.b. convergence are *i -structures*.

EXAMPLE 2. Among all *i -topologies* on a given poset P admitting closed maximal chains, there is a coarsest one denoted by $w(P)$. Indeed, given the maximal chains J of P , let $w(P)$ arise from the sub-base of closed sets $\{a^* \cap J, a^+ \cap J : a \in J\}$. It is easily seen $w(P)$ has the desired properties; also note that for $P \equiv]0, 1[\times]0, 1[$, no set of the form a^* or a^+ is $w(P)$ -closed. There are even *i -spaces* in which no maximal chain is closed, cf. the lattice $\{0, a, b, 1\}$, a and b non-related, endowed with the topology whose closed sets are generated by $\{a, b\}$, $\{0\}$, $\{1\}$. We note that if the maximal chains of an *i -space* (P, q) are closed sets, then $\text{top}(q)$ is an *i -topology* on P .

The following example shows that on posets, which are not lattices, order convergence misbehaves.

EXAMPLE 3. Let P be the subposet of \mathbf{R}^2 consisting of the strictly negative part of the x -axis, all y -axis for $y \geq 1$, and the point $(1, 0) = \alpha$. Clearly, $o(P)(\alpha) = \{[\alpha]\}$, which means the restriction of $o(P)$ to the maximal chain containing α is strictly finer than the interval topology of that chain.

From now on, in this section we will restrict our attention to lattices only. Our first aim is to generalize two lemmata of Vainio [9], originally proved assuming conditional

completeness. First we mention the obvious

LEMMA 4. *Let L be any lattice and \mathcal{F} a filter on L arising from some filter base \mathcal{B} of sets B for which $\vee_L B$ and $\wedge_L B$ exist. Then, for all $x \in X$*

$\mathcal{F} \in o(L)(x) \Leftrightarrow \vee_L \{\wedge_L B : B \in \mathcal{B}\}, \wedge_L \{\vee_L B : B \in \mathcal{B}\}$ both exist and both equal x .

LEMMA 5 (cf. Vainio [9, Lemma 3]). *Let L be any lattice, J a maximal chain of L , and S a subchain of J . Then,*

$\vee_J S$ exists $\Rightarrow \vee_L S$ exists and equals $\vee_J S$, and dually.

PROOF. Assume $\vee_J S = \alpha, \alpha \notin S$, and let $\beta \leq \alpha$ be an L upper bound of S . Then, $s \leq \beta \leq \alpha$ for all $s \in S$. Of course, $\beta \notin S$. Thus, $\beta < \alpha$ would mean J is not a maximal chain. Hence $\beta = \alpha$, and we have revealed α is the smallest L upper bound of S . \diamond

Example 3 shows Lemma 5 cannot be extended to posets.

LEMMA 6. (cf. Vainio [9, Lemma 7]). *For an arbitrary lattice L , any convergence structure between $t(L)$ and $o(L)$ is an i -structure.*

PROOF. Let J be any maximal chain of L , take $x \in J$, and denote the $t(J)$ neighborhood filter of x by \mathcal{F} . Since $t(J)$ equals $o(J)$, Lemma 4 gives \mathcal{F} has a base \mathcal{B} consisting of bounded J -intervals B such that $\vee_J \{\wedge_J B : B \in \mathcal{B}\}$ and $\wedge_J \{\vee_J B : B \in \mathcal{B}\}$ both exist and both equal x . (Note that any bounded J -interval possesses l.u.b. and g.l.b. in J .) According to Lemma 5, the index L can replace the index J in the above expressions, and thus, \mathcal{F} is a base of a filter on L order-converging towards x . Hence, $o(L)$ is an i -structure. Since $t(L)_J \geq t(J)$ and $t(L) \leq o(L)$, we get $t(L)$ is an i -structure, too. \diamond

COROLLARY 7. *On every lattice L , $o(L) \leq s(L)$. This result is not true for arbitrary posets (cf. Ex. 3).*

EXAMPLE 8. In an arbitrary lattice, there is a host of well-known convergence structures in between interval topology and order convergence. Erné [2] mentions among the topologies, e.g. interval topology, new interval topology, Lawson topology, bi-Scott topology, Rennie's L -topology, lim-inf topology, Birkhoff's order topology, and among the convergences (through which some of the previous topologies can be described), e.g. interval convergence, lim-inf convergence, and order convergence. In view of Lemma 6, we know all of them are i -structures.

The remainder of the paper will give further motivation for investigating i -spaces.

3. COMPACTNESS AND CONNECTIVITY

Let L be an arbitrary lattice. We say a property p of L is *described by maximal chains*, if

L satisfies $p \Leftrightarrow$ All maximal chains of L satisfy p .

Kogalovskii [6] proves completeness is described by maximal chains. We will note Lemma 5 and Rennie [7, Theorem 1] together imply that conditional completeness is described by maximal chains (also, cf. Vainio [9]), and from this Kogalovskii's result follows.

LEMMA 9. *In any lattice, the following properties are described by maximal chains: (1) convexity, (2) order density, (3) conditional completeness, (4) completeness.*

PROOF. (3) Assume all maximal chains of L are conditionally complete, let S be a chain of L with upper bound m , and let J be a maximal chain containing S and m . Hence $\vee_L S$ exists (Lemma 5), and L is conditionally complete according to Rennie [7, Theorem 1]. The converse statement is trivial (cf. for instance Vainio [9, Lemma 3]). (4) Assume all maximal chains of L are complete, and let J_1 and J_2 be maximal chains with greatest elements a_1, a_2 . Since $a_1 \vee a_2 \in J_1 \cap J_2$, then all maximal chains have the same greatest element, which proves L has a greatest element. Applying (3) above the proof is completed.◊

COROLLARY 10. *A lattice is (conditionally) complete, if and only if all maximal chains of the lattice are (conditionally) subcomplete.*

Using i.a. Lemma 9 and the well-known fact that any chain J is complete (conditionally complete), if and only if $t(J)$ is compact (all bounded ultrafilters on J are $t(J)$ -convergent), we obtain the following two theorems.

THEOREM 11. *For q an i -convergence on any lattice L , the following are equivalent:*

- (1) L is complete.
- (2) (L, q) has compact maximal chains.

If $q \leq t(L)$, but all maximal chains of (L, q) still are closed sets, then expressions

(1) and (2) are equivalent to

- (3) (L, q) is compact.

PROOF. For part (3), use the well-known fact that a lattice is complete, if and only if it is compact in its interval topology.◊

For L an arbitrary lattice, denote by $w(L)$ the coarsest i -convergence on L admitting closed maximal chains ($w(L)$ is always a topology, cf. Ex. 2), and take $w(L) \leq q \leq s(L)$.

THEOREM 12. *A lattice L is (conditionally) complete, if and only if all (bounded) ultrafilters on L containing some maximal chain of L are q -convergent.*

We next improve Theorem 4 of Vainio [9]. Proofs are omitted; the reader is referred to the well-known fact that any chain J is conditionally complete and order dense, if and only if $t(J)$ is connected, to Lemma 9, and to the proof of [9, Theorem 4]. Note that any convex subspace of an i -space is again an i -space. Definitions regarding connectivity of convergence spaces are as in Gähler [3] or Vainio [9]. Below, L is always a lattice.

THEOREM 13. *Every conditionally complete, order dense, convex sublattice of an i -space (L, q) is connected.*

THEOREM 14. *Let L be conditionally complete, and let (L, q) be a T_1 i -space in which all translations are continuous maps. Then, all connected components of (L, q) are order dense, convex (and hence, conditionally complete) sublattices of L .*

COROLLARY 15. *In conditionally complete T_1 i -spaces (L, q) with continuous translations, the connected components can be described as the maximal order dense, convex*

sublattices of L .

Let $X \equiv (X, q)$ be a convergence space and recall that (X, q) and $(X, \text{top}(q))$ yield the same real-valued continuous maps, denoted by $C(X)$. It is known (cf. Stone [8]) that the lattice $C(X)$ is conditionally complete, if and only if $\text{top}(q)$ is extremely disconnected. We next characterize conditional completeness of $C(X)$ in terms of any i -structure q on $C(X)$.

THEOREM 16. *The lattice $C(X)$ is conditionally complete, if and only if all maximal chains in $C(X)$ are q -connected sets.*

The proof is an easy consequence of Lemma 9 and of the fact that $C(X)$ is order dense.

4. PRODUCTS OF i -SPACES

Category theoretical definitions are as in Herrlich [4].

At first, we will regard a somewhat more general (but from a category theoretical point of view more natural) situation than i -space compatibility. The results will then be applied to i.a. formation of products of i -spaces.

Let S be a *quasi-ordered* set endowed with a convergence q such that for any maximal chain J of S the inequality $q_J \geq t(J)$ holds true. The corresponding category (morphisms: increasing continuous maps) is denoted by $QCON$ and the full subcategory determined by the partially ordered objects is called $PCON$. The first category is mainly a tool, used to obtain Remark 18 below.

LEMMA 17. *The concrete category $QCON$ is topological.*

PROOF. Let $(S_i, q_i)_{i \in I}$ be an arbitrary family of $QCON$ -objects, consider a set S and set theoretical maps $f_i : S \rightarrow S_i, i \in I$. Then, for any $x, y \in S$, define

$$x \leq y \text{ in } S \Leftrightarrow f_i(x) \leq f_i(y) \text{ in } S_i, \text{ all } i \in I.$$

$$\mathcal{F} \in q(x) \Leftrightarrow f_i(\mathcal{F}) \in q_i(f_i(x)), \text{ all } i \in I.$$

Obviously, S is thus endowed with a quasi-order \leq and a convergence q . It remains to prove the resulting space is a $QCON$ -object. Therefore, let J be a maximal chain in S , and let J_i be *one* maximal chain in S_i containing the chain $f_i(J)$. For any $i \in I$, denote the restriction of f_i to J by g_i , and note that for $\alpha \in J$

$$\alpha^+ = \bigcap_{i \in I} g_i^{-1}(g_i(\alpha)^+)$$

where α^+ and $g_i(\alpha)^+$ are formed in the chains J and J_i , respectively. Since all maps g_i are continuous, the sets α^+ (and dually, α^*) are closed sets in q_J , and hence $q_J \geq t(J)$. \diamond

There are interesting consequences of Lemma 17. Since $QCON$ is a topological category, Herrlich [4; Section 2.1, T(5)] quite easily gives $QCON$ is a cartesian closed category. Power objects in $QCON$ are the spaces $CI_c(S, T)$, i.e. the set of $QCON$ -morphisms (between given $QCON$ -objects S and T) endowed with product order and continuous convergence structure. Indeed, we obtain (all proofs omitted)

REMARK 18. The topological category $QCON$ is cartesian closed, and contains $PCON$ as a quotient-reflective subcategory; hence $PCON$ is cartesian closed.

Needless to say, Lemma 17 and Remark 18 do not apply to i -space compatibility.

It is an immediate consequence of the proof of Lemma 17 that $QCON$ (and thus, $PCON$) is closed under formation of arbitrary products and subspaces. This implies a few remarks on orderability. We call a convergence space p -orderable, if there is a partial order on X which makes (X, q) a $PCON$ -object. (If the po-structure in question is a total order and q a topology, we get the well-known orderable spaces.) Then let $PORD$ denote the full subcategory of the convergence space category CON consisting of all p -orderable spaces. Since CON is a topological category, any subcategory of this category which is closed under formation of products and subspaces is reflective in it. Thus, $PORD$ is a reflective subcategory of CON .

The remainder of the paper deals with a special construction; *products of i -spaces*. Consider a family $(P_i, q_i)_{i \in I}$ of i -spaces, each P_i being a poset and each q_i a convergence. Denote the product space by (P, q) , i.e. P is the product poset and q the product convergence. Since $PCON$ is closed under formation of arbitrary products, (P, q) is an i -space, if and only if $\prod q_i \leq s(P)$. This condition is equivalent to continuity of all projections $pr_i : (J, t(J)) \rightarrow (P_i, q_i)$, where J is an arbitrary maximal chain of P and $i \in I$. This is satisfied at least, if for all i, J as above, each set $pr_i(J)$ is a *convex chain in P_i* (i. e. $pr_i(J)$ is a convex subset of any maximal chain of P_i containing it). The following lemma gives a natural sufficient condition.

Let $(L_i)_{i \in I}$ be *conditionally complete lattices*, denote $L \equiv \prod L_i$, and let J be a maximal chain of L .

LEMMA 19. *For each $i \in I$, $pr_i(J)$ is a convex chain in L_i .*

PROOF. Assume $pr_\alpha(J)$ not convex, and let h_α be a "hole" in it. Define the subset A of J by

$$x \in A \Leftrightarrow x \in J \text{ and } pr_\alpha(x) > h_\alpha.$$

Let $z \in L$ be the element defined by $pr_i(z) = \wedge pr_i(A)$ for $i \in I, i \neq \alpha$, and $pr_\alpha(z) = h_\alpha$. (The conditional completeness assumption implies $\wedge pr_i(A)$ exists.) We now prove $z \in J$ by showing z is order related to an arbitrary $s \in J$.

If $pr_\alpha(s) > h_\alpha$, then $s \in A$, and for $i \neq \alpha$ $pr_i(s) \geq \wedge pr_i(A) = pr_i(z)$, which gives $s > z$.

If $pr_\alpha(s) < h_\alpha$, then $s < x$ for all $x \in A$, and hence for $i \neq \alpha$ $pr_i(s) \leq \wedge pr_i(A) = pr_i(z)$, and $s < z$ follows.

Now, $h_\alpha = pr_\alpha(z) \in pr_\alpha(J)$, a contradiction. \diamond

THEOREM 20. *The product of given i -spaces $(L_i, q_i)_{i \in I}$ is an i -space, if every L_i is a conditionally complete lattice.*

REFERENCES

1. M. ERNÉ, *Topologies on products of partially ordered sets I-III*, Algebra Univ. **11** (1980) no 3, pp. 295-311; ibid. **11** (1980) no 3, pp. 312-319; ibid. **13** (1981) no 1, pp. 1-23.
2. —and H. GATZKE, *Convergence and continuity in partially ordered sets and semilattices*, in Continuous Lattices and their Appl., Proc. Bremen 1982, M.Dekker.
3. W. GÄHLER, *Grundstrukturen der Analysis I-II*, Akademie-Verlag und Birkhäuser Verlag, 1977-78.
4. H. HERRLICH, *Categorical topology 1971-1981*, Proc. Symp. Gen. Top. Appl. Prague 1981, pp. 279-383.
5. D. C. KENT, *On convergence groups and convergence uniformities*, Fund. Math. **60** (1967), pp. 213-222.
6. S. R. KOGALOVSKIĬ, *On linearly complete ordered sets (Russian)*, Uspehi Mat. Nauk. **19** (1964) no 2, pp. 147-150.
7. B. C. RENNIE, *Lattices*, Proc. London Math. Soc. **52** (1951), pp. 386-400.
8. M. H. STONE, *Boundedness properties in function-lattices*, Canad. J. Math. **1** (1949), pp. 176-186.
9. R. VAINIO, *Connectedness properties of lattices*, Canad. Math. Bull. **29** (1986) no 3, pp. 314-320.

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk