

## ON A GLOBAL CLASSICAL SOLUTION OF A QUASILINEAR HYPERBOLIC EQUATION

Y. EBIHARA

Department of Applied Mathematics  
Faculty of Science, Fukuoka University  
Fukuoka 814-01 Japan

D.C. PEREIRA

Instituto de Mathematica-UFRJ  
CP 68530, CEP 21944  
Rio de Janeiro-Brasil

(Received June 16, 1987)

**ABSTRACT.** In this paper we establish the existence and uniqueness of global classical solutions for the equation which arises in the study of the extensional vibrations of thin rod, or torsional vibrations of thin rod.

### 1. INTRODUCTION.

In this paper we study the existence and uniqueness of global classical solutions of the first initial-boundary value problem for the equation:

$$u_{tt} - \Delta u - M \left( \int_{\Omega} |\nabla u|^2 dx \right) \Delta u_{tt} = f \quad (1.1)$$

in  $Q = \Omega \times ]0, T[$ , where  $\Omega (\partial \Omega)$ : the boundary is a smooth bounded domain in  $\mathbb{R}^n$ ,  $T$  is a positive number,  $\nabla u$  is the gradient of  $u$ ,  $\Delta$  is the Laplace operator and  $M(\lambda)$ ,  $\lambda > 0$ , is a real valued function with  $M(\lambda) > \rho > 0$ ,  $\lambda > 0$ , for some  $\rho > 0$ . We have mathematical interest in solving the equation (1.1) by the following reasons.

First, the equation (1.1) with  $M(\lambda) \equiv 1$  arises in the study of the extensional vibrations of thin rods, see Love [1], and it was studied by one of the authors in [2] and [3]. Second, the equation (1) with  $M(\lambda) \equiv \lambda_0$ ,  $\lambda_0 = \int_{\Omega} \phi^2 dx$ , where  $\phi$  is the

torsion-function arises in the study of the torsional vibrations of thin rods, see Love [1]. Third, the function  $M(\lambda)$  in (1.1) has its motivation in the mathematical description of the vibrations of an elastic stretched string, that is, the equation:

$$u_{tt} - M \left( \int_{\Omega} |\nabla u|^2 dx \right) \Delta u = 0 \quad (1.2)$$

what for  $M(\lambda) > \rho > 0$  was studied by Pohozaev [4], Nishihara [5] and Lions [6]. When  $M(\lambda) > 0$  was treated by Arosio-Spagnolo [7], Ebihara-Medeiros-Milla Miranda [8] and Yamada [9].

In this paper, we establish the existence and uniqueness of global classical solutions for the equation (1.1). For that we use the Faedo-Galerkin method and compactness argument with some technical idea.

## 2. NOTATIONS, ASSUMPTIONS AND MAIN RESULT.

Let  $(w_j)_{j \in \mathbb{N}}$  be a system of eigen functions of  $-\Delta$  which is defined on  $H^2(\Omega) \cap H_0^1(\Omega)$ . We denote by  $V \equiv V(\Omega)$  the set of all finite linear combinations of  $(w_j)_{j \in \mathbb{N}}$ . Putting  $(f, g) = \int_{\Omega} f(x)g(x)dx$ , we set  $(\cdot, \cdot)_m = ((-\Delta)^m \cdot, \cdot)$ ,  $m=1, 2, \dots$ , then  $(\cdot, \cdot)_m$  define an inner product on  $V$ . We put  $V_m \equiv V_m(\Omega)$  as the closure of  $V$  by the topology of norm  $|\cdot|_m^2 = (\cdot, \cdot)_m$ . Then we see that

$$H_0^1(\Omega) \equiv V_1 \hookrightarrow V_2 \hookrightarrow \dots \hookrightarrow V_m \hookrightarrow \dots,$$

$V_m \subset H^m(\Omega)$ ,  $m = 1, 2, \dots$ , and the norm  $|\cdot|_m$  is equivalent in  $V_m$  to the standard norm of  $H^m(\Omega)$ . We see that all the above injections " $\hookrightarrow$ " are compact.

Let  $T$  be a positive number and  $B$  a Banach space with a norm  $\|\cdot\|_B$ . We shall represent by  $L^p(0, T; B)$ ,  $1 < p < \infty$ , the Banach space of vector-valued functions  $u: [0, T] \rightarrow B$  which are measurable in  $B$  and  $\|u(t)\|_B \in L^p(0, T)$  with the norm

$$\|u\|_{L^p(0, T; B)}^p \equiv \int_0^T \|u(t)\|_B^p dt$$

and by  $L^\infty(0, T; B)$  the Banach space of vector-valued functions  $u: [0, T] \rightarrow B$  which are measurable in  $B$  and  $\|u(t)\|_B \in L^\infty(0, T)$  with the norm

$$\|u\|_{L^\infty(0, T; B)} \equiv \sup_{t \in [0, T]} \|u(t)\|_B.$$

We denote by  $C^j(0, T; B)$  the space of all vector-valued functions  $u: [0, T] \rightarrow B$ , which are  $j$ -times differentiable in the sense of  $B$ .

Let  $M(\lambda)$ ,  $\lambda > 0$  be a real valued function such that:

(A.1)  $M(\lambda) \in C^1[0, \infty)$  and there exist constants  $\alpha > 0$  and  $\rho > 0$  that

verify  $M(\lambda) > \alpha \lambda^{1/2} + \rho$ ,  $\forall \lambda \in [0, \infty)$ ,

(A.2)  $|M'(\lambda)|\lambda^{1/2} < \beta(\lambda)M(\lambda)$  where  $\beta(\lambda) \in C^0[0, \infty)$ ,  $\beta(\lambda) > 0$ ,  $\lambda > 0$ , then we have the following result:

THEOREM 2.1. Suppose that

$$u_0, u_1 \in V_m, (m \geq 2), \quad (2.1)$$

$$f, f' \in C(0, T; V_{m-1}). \quad (2.2)$$

Then there exists a unique function  $u: [0, T] \rightarrow L^2(\Omega)$  in the class:

$$u \in C^2(0, T; V_m) \quad (2.3)$$

that verifies

$$u'' - \Delta u - M(|u|_1^2) \Delta u'' = f \text{ in } Q \quad (2.4)$$

$$u(0) = u_0 \quad (2.5)$$

$$u'(0) = u_1 \quad (2.6)$$

$$u|_{\partial\Omega} = 0 \quad (2.7)$$

### 3. PROOF OF THEOREM 1.

We divide the proof in four parts:

- a) Approximated solutions
- b) A priori estimates
- c) Passage to the limit
- d) Uniqueness

#### a) APPROXIMATED SOLUTIONS.

Let  $[w_1, \dots, w_k]$  be the subspace of  $V$ , generated by the first  $k$  eigenvectors of  $-\Delta$ .

Let

$$u_k(t) \equiv \sum_{j=1}^k g_{jk}(t) w_j \in [w_1, \dots, w_k]$$

be a solution of the system:

$$(u_k'' - \Delta u_k - M(|u_k|_1^2) \Delta u_k'', w) = (f, w) \text{ for all } w \in [w_1, \dots, w_k] \quad (3.1)$$

$$u_k(0) = u_{0k} \rightarrow u_0 \text{ strongly in } V_m \text{ as } k \rightarrow \infty, \quad (3.2)$$

$$u_k'(0) = u_{1k} \rightarrow u_1 \text{ strongly in } V_m \text{ as } k \rightarrow \infty, \quad (3.3)$$

$$\text{where } u_{0k} = \sum_{j=1}^k (u_0, w_j) w_j, \quad u_{1k} = \sum_{j=1}^k (u_1, w_j) w_j.$$

Then we see that the solution  $u_k(t)$  exists on an interval  $[0, T_k]$ ,  $0 < T_k < T$ . A priori estimates will permit us to extend  $u_k(t)$  to all interval  $[0, T]$ .

b) A PRIORI ESTIMATES

I) Putting  $w = u_k''(t)$  in (3.1),

we have

$$|u_k''|_0^2 + (u_k, u_k'')_1 + M(|u_k|_1^2) |u_k''|_1^2 = (f, u_k'').$$

Thus by (A.1),

$$\begin{aligned} \frac{|u_k''|_0^2}{M(|u_k|_1^2)} + |u_k''|_1^2 &< \frac{|u_k|_1 |u_k''|_1}{\alpha |u_k|_1} + \frac{|f|_0 |u_k''|_0}{\rho} \\ &< \frac{1}{\alpha^2} + \frac{1}{4} |u_k''|_1^2 + \frac{c^2 |f|_0^2}{\rho^2} + \frac{1}{4} |u_k''|_1^2 \end{aligned}$$

where  $C$  is a positive constant such that  $|u_k''|_0 < c |u_k''|_1$ . Then from this and (2.2),

$$|u_k''|_1^2 < 2 \left( \frac{1}{\alpha^2} + \frac{c^2 |f|_0^2}{\rho^2} \right) < c_1$$

where  $c_1 > 0$  is a constant independent of  $t$  and  $k$ .

Thus

$$(u_k'') \text{ is bounded in } L^\infty(0, T_k; V_1). \quad (3.4)$$

By Fundamental Theorem of Calculus, we have

$$(u_k') \text{ is bounded in } L^\infty(0, T_k; V_1) \quad (3.5)$$

and

$$(u_k) \text{ is bounded in } L^\infty(0, T_k; V_1). \quad (3.6)$$

The above estimates permit us to extend  $u_k(t)$  to all interval  $[0, T]$ .

II) Making  $w = (-\Delta)^{m-1} u_k'$ , in (3.1)

we have:

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} [ |u_k'|_{m-1}^2 + |u_k|_m^2 + M(|u_k|_1^2) |u_k'|_m^2 ] \\ = M'(|u_k|_1^2) (u_k, u_k')_1 |u_k'|_m^2 + (f, u_k')_{m-1}. \end{aligned}$$

By (A.2) and (3.6), we have

$$\begin{aligned} M'(|u_k|_1^2) (u_k, u_k')_1 |u_k'|_m^2 &< \beta (|u_k|_1^2) M(|u_k|_1^2) |u_k'|_1 |u_k'|_m^2 \\ &\leq \frac{C_2}{2} M(|u_k|_1^2) |u_k'|_m^2. \end{aligned}$$

Then integrating from 0 to  $t$  and using (2.1) and (2.2) we obtain:

$$\begin{aligned} |u_k|_{m-1}^2 + |u_k|_m^2 + M(|u_k|_1^2) |u_k'|_m^2 \\ \leq C_3 + C_4 \int_0^t [|u_k'|_{m-1}^2 + M(|u_k|_1^2) |u_k'|_m^2] ds \end{aligned}$$

where  $C_3 = C_3(u_0, u_1, T)$  and  $C_4 = \max\{1, C_2\}$ .

Thus, by Gronwall inequality and (A.1), we have:

$$|u_k|_{m-1}^2 + |u_k|_m^2 + \rho |u_k'|_m^2 \leq C_5$$

where  $C_5 > 0$  is a constant independent of  $t$  and  $k$ .

Whence,

$$(u_k) \text{ is bounded in } L^\infty(0, T; V_m), \quad (3.7)$$

$$(u_k') \text{ is bounded in } L^\infty(0, T; V_m). \quad (3.8)$$

(III) Taking the derivative of (3.1) with respective to  $t$ , we obtain:

$$\begin{aligned} (u_k^{(3)}) - \Delta u_k' - M(|u_k|_1^2) \Delta u_k^{(3)} \\ - 2M'(|u_k|_1^2) (u_k, u_k') \Delta u_k'' + (f', w) \text{ for all } w \in [w_1, \dots, w_k]. \end{aligned} \quad (3.9)$$

Putting in (3.9)  $w = (-\Delta)^{m-1} u_k''$ , we have:

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} [|u_k''|_{m-1}^2 + |u_k'|_m^2 + M(|u_k|_1^2) |u_k''|_m^2] \\ + M'(|u_k|_1^2) (u_k, u_k') |u_k''|_m^2 = (f', u_k'')_{m-1}. \end{aligned}$$

Thus by (3.2) and (3.6),

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} [|u_k''|_{m-1}^2 + |u_k'|_m^2 + M(|u_k|_1^2) |u_k''|_m^2] \\ \leq \frac{C_2}{2} M(|u_k|_1^2) |u_k''|_m^2 + \frac{1}{2} |f'|_{m-1}^2 + \frac{1}{2} |u_k''|_{m-1}^2. \end{aligned}$$

Integrating from 0 to  $t$ , we have:

$$|u_k''|_{m-1}^2 + |u_k'|_m^2 + M(|u_k|_1^2) |u_k''|_m^2$$

$$\begin{aligned}
& \leq \int_0^t \left| f'(s) \right|_{m-1}^2 ds + C_4 \int_0^T [\left| u_k''(s) \right|_{m-1}^2 + M(\left| u_k \right|_1^2) \left| u_k''(s) \right|_m^2] ds \\
& \quad + \left| u_k''(0) \right|_{m-1}^2 + \left| u_{1k} \right|_m^2 + M(\left| u_{0k} \right|_1^2) \left| u_k''(0) \right|_m^2. \tag{3.10}
\end{aligned}$$

We now estimate  $\left| u_k''(0) \right|_m$ .

Putting  $w = (-\Delta)^{m-1} u_k''(0)$  in (3.1) and tending  $t \rightarrow 0$ , we obtain:

$$\begin{aligned}
& \left| u_k''(0) \right|_{m-1}^2 + (u_k(0), u_k''(0))_m + M(\left| u_{0k} \right|_1^2) \left| u_k''(0) \right|_m^2 \\
& = (f(0), u_k''(0))_{m-1}.
\end{aligned}$$

Thus,

$$\begin{aligned}
& \left| u_k''(0) \right|_{m-1}^2 + M(\left| u_{0k} \right|_1^2) \left| u_k''(0) \right|_m^2 \leq \frac{1}{\gamma} \left| u_{0k} \right|_m^2 + \gamma \left| u_k''(0) \right|_m^2 \\
& \quad + \frac{1}{2} \left| f(0) \right|_{m-1}^2 + \frac{1}{2} \left| u_k''(0) \right|_{m-1}^2
\end{aligned}$$

for  $\gamma > 0$ .

Then, by (A.1), (2.2) and (3.2), we have

$$\left| u_k''(0) \right|_{m-1}^2 + 2(\rho - \gamma) \left| u_k''(0) \right|_m^2 \leq C_6$$

where  $C_6 > 0$  is a constant independent of  $t$  and  $k$ .

Whence, taking  $\gamma$  as  $0 < \gamma < \rho$ , we obtain:

$$\left| u_k''(0) \right|_m^2 \leq C. \tag{3.11}$$

Thus, by (3.10) and (3.11), we have:

$$\begin{aligned}
& \left| u_k'' \right|_{m-1}^2 + \left| u_k' \right|_m^2 + M(\left| u_k \right|_1^2) \left| u_k'' \right|_m^2 \\
& \leq C_7 + C_4 \int_0^t [\left| u_k'' \right|_{m-1}^2 + M(\left| u_k \right|_1^2) \left| u_k'' \right|_m^2] ds.
\end{aligned}$$

And by Gronwall lemma and (A.1), we obtain:

$$\left| u_k'' \right|_{m-1}^2 + \left| u_k' \right|_m^2 + \rho \left| u_k'' \right|_m^2 \leq C_8$$

where  $C_8 > 0$  is a constant independent of  $t$  and  $k$ .

Whence,

$$(u_k'') \text{ is bounded in } L^\infty(0, T; V_m). \quad (3.12)$$

IV) Putting in (3.9)  $w = (-\Delta)^{m-1} u_k^{(3)}$ ,  
we obtain:

$$\begin{aligned} & |u_k^{(3)}|_{m-1}^2 + (u_k', u_k^{(3)})_m + M(|u_k|_1^2) |u_k^{(3)}|_m^2 \\ & + 2M'(|u_k|_1^2) (u_k', u_k')_1 (u_k'', u_k^{(3)})_m = (f', u_k^{(3)})_{m-1}. \end{aligned}$$

Thus by (A.1), (A.2), (3.7), (3.8) and (3.12), we have:

$$\begin{aligned} & |u_k^{(3)}|_{m-1}^2 + \rho |u_k^{(3)}|_m^2 \leq \frac{C_9}{\gamma} + \gamma |u_k^{(3)}|_m^2 + \frac{1}{2} |f'|_{m-1}^2 \\ & + \frac{1}{2} |u_k^{(3)}|_{m-1}^2. \end{aligned}$$

Then

$$|u_k^{(3)}|_{m-1}^2 + 2(\rho - \gamma) |u_k^{(3)}|_m^2 \leq C_{10}$$

where  $C_{10} > 0$  is a constant independent of  $t$  and  $k$ . Whence we can assert that

$$(u_k^{(3)}) \text{ is bounded in } L^\infty(0, T; V_m), \quad (3.13)$$

c) PASSAGE TO THE LIMIT.

By estimates (3.7), (3.8), (3.12) and (3.13) there is a subsequence of  $(u_k)_{k \in \mathbb{N}}$  what we denote by  $(u_k)_{k \in \mathbb{N}}$  and there exists a function  $u$ , such that:

$$u_k \xrightarrow{*} u \text{ weak star in } L^\infty(0, T; V_m) \quad (3.14)$$

$$u_k' \xrightarrow{*} u' \text{ weak star in } L^\infty(0, T; V_m) \quad (3.15)$$

$$u_k'' \xrightarrow{*} u'' \text{ weak star in } L^\infty(0, T; V_m) \quad (3.16)$$

$$u_k^{(3)} \xrightarrow{*} u^{(3)} \text{ weak star in } L^\infty(0, T; V_m) \quad (3.17)$$

By (3.14) and (3.15), for  $m=2$ , and since the embedding of  $V_2$  is compact in  $V_1$ , it follows from Aubin-Lions Theorem [10],

$$u_k \rightarrow u \text{ strongly in } L^2(0, T; V_1)$$

whence

$$M(|u_k|_1^2) \rightarrow M(|u|_1^2) \text{ strongly in } L^2(0, T). \quad (3.18)$$

Now by (3.16),

$$\Delta u_k'' + \Delta u'' \text{ weak in } L^2(0, T; V_{m-2}). \quad (3.19)$$

Thus, by (3.18) and (3.19) we have

$$M(|u_k|_1^2) \Delta u_k'' + M(|u|_1^2) \Delta u'' \text{ weak in } L^2(0, T; V_{m-2}) \quad (3.20)$$

The above convergences permit us to pass to the limit in the approximate equation (3.1), as  $k \rightarrow \infty$ . We then get:

$$(u'' - \Delta u - M(|u|_1^2) \Delta u'', w) = (f, w)$$

for each  $w \in V$  in the sense of  $L^2(0, T)$ .

REMARK 1. Since the solution  $u$  of (2.4) is in  $C^2(0, T; V_m)$ , and by Sobolev's theorem

$$H^k(\Omega) \subset C^k(\bar{\Omega})$$

with  $k < \ell - \frac{n}{2} < k + 1$ ,  $k \geq 0$  integer,

then  $u$  satisfies (2.4) in the classic sense if we choose  $m$  large enough.

#### d) UNIQUENESS

Let  $u, v$  be solutions of (1.1) in the class of Theorem 1. Then  $\omega = u - v$  satisfies:

$$\omega'' - \Delta \omega - M(|u|_1^2) \Delta \omega'' - [M(|v|_1^2) - M(|u|_1^2)] \Delta v'' = 0 \quad (3.21)$$

$$\omega(0) = 0 \quad (3.22)$$

$$\omega'(0) = 0 \quad (3.23)$$

Taking of scalar product in  $L^2(\Omega)$  of (3.21) by  $\omega'$ , we obtain:

$$\frac{1}{2} \frac{d}{dt} [|\omega'|_0^2 + |\omega|_1^2 + M(|u|_1^2) |\omega'|_1^2] - M'(|u_1|^2) (u, u')_1 |\omega'|_1^2$$

$$+ [M(|v|_1^2) - M(|u|_1^2)] (v'', \omega')_1 = 0$$

Now, by (A.1) and (A.2), we have

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} [|\omega'|_0^2 + |\omega|_1^2 + M(|u|_1^2) |\omega'|_1^2] \\ & \leq \frac{C_1}{2} M(|u|_1^2) |\omega'|_1^2 + M'(\xi) (|u|_1 + |v|_1) |\omega|_1 |\omega'|_1, \end{aligned}$$

where  $\xi = (1 - \theta) |u|_1^2 + \theta |v|_1^2$ ,  $0 < \theta < 1$ .

Then,

$$\frac{1}{2} \frac{d}{dt} [|\omega'|_0^2 + |\omega|_1^2 + M(|u|_1^2)|\omega'|_1^2] \leq \frac{C}{2} (|\omega|_1^2 + |\omega'|_1^2),$$

whence by (A.1),

$$|\omega'|_0^2 + |\omega|_1^2 + \rho |\omega'|_1^2 \leq C \int_0^t (|\omega|_1^2 + |\omega'|_1^2) ds.$$

Thus, we have

$$\omega \equiv 0 \text{ in } [0, T]$$

REMARK 2. In the forthcoming work we will try to study the equation (1.1) when  $M(\lambda)$  has zero points, that is, degenerate case.

#### REFERENCES

1. LOVE, A.H. A treatise on the mathematical theory of elasticity. Dover New York 1944.
2. PEREIRA, D.C. and IZAGUIRRE, R.M. Sobre uma equacao de vibracoes nao lineares. Proceedings of 23o seminario Brasileiro de Analise, Campinas, Sao Paulo (1986), 155-158.
3. PEREIRA, D.C. Problema misto para uma equacao de vibracoes nao lineares. To appear.
4. POHOZAEV, S.I. On a class of quasilinear hyperbolic equations. Mat USSR Sbornik 25,1 (1975), 145-148.
5. NISHIHARA, K. On a global solution of some quasilinear hyperbolic equation. Tokyo Journal of Math. 7 (1984).
6. LIONS, J.L. On some questions in boundary value problems of mathematical physics. IM-UFRJ (1978).
7. AROSIO, A. and SPAGNOLLO, S. Global solutions of the Cauchy Problem for a nonlinear hyperbolic equation. Universita de Pisa, Dipartimento di Mathematica, Roma (1982).
8. EBIHARA, Y., MEDEIROS, L.A. and MILLA MIRANDA, M. Local solutions for a nonlinear degenerate hyperbolic equation. Nonlinear Analysis, Theory Methods and Applications, 10 (1986), 27-40.
9. YAMADA, Y. Some nonlinear degenerate wave equation. To appear in Nonlinear Analysis, Theory, Methods and Applications.
10. LIONS, J.L. Quelques methodes de resolution des problemes aux limites non lineaires. Dunod, Paris, (1969).

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)