

SOME REMARKS ON INEQUALITIES THAT CHARACTERIZE INNER PRODUCT SPACES

JAVIER ALONSO

Departamento de Matemáticas
Universidad de Extremadura
06071 Badajoz, Spain

(Received November 28, 1990 and in revised form March 11, 1991)

ABSTRACT. The purpose of this paper is to obtain some new characterizations of inner product spaces in the line of the Jordan-von Neumann identity and to make some remarks on similar known characterizations of Day, Delbosco, Rassias and Senechalle, among others.

KEY WORDS AND PHRASES. Inner product spaces, characterizations, inequalities.

1980 AMS SUBJECT CLASSIFICATION CODE. 46B20.

Let E be a real normed linear space. It is well known that E is an inner product space (i.p.s.) if and only if the "parallelogram equality" of P. Jordan and J. von Neumann [1]

$$\|x+y\|^2 + \|x-y\|^2 = 2(\|x\|^2 + \|y\|^2)$$

holds for every $x, y \in E$. The aim of this paper is to give an account of some characterizations of inner product spaces weaker than the one above in order to make some remarks about them and to obtain some new characterizations of the same type.

We shall use S to denote the unit sphere of E , and $H(x, y)$ the function

$$H(x, y) = \|x+y\|^2 + \|x-y\|^2 - 2(\|x\|^2 + \|y\|^2) \quad (x, y \in E).$$

The first weakening of the Jordan-von Neumann condition was given by M.M. Day [2] by proving that it can be restricted to unit vectors (i.e. the "rhombus equality"):

$$E \text{ is an i.p.s.} \Leftrightarrow H(x, y) = 0 \text{ for every } x, y \in S.$$

Then I.J. Schoenberg [3] proved that the rhombus equality can be replaced by the "rhombus inequality":

$$E \text{ is an i.p.s.} \Leftrightarrow H(x, y) \sim 0 \text{ for every } x, y \in S,$$

where \sim denotes either \geq or \leq .

The parallelogram equality can also be changed into the "rectangle inequality" of C. Benítez and M. del Río [4]:

$$E \text{ is an i.p.s.} \Leftrightarrow H(x, y) \sim 0 \text{ for every } x, y \in E, x \perp y,$$

where $x \perp y$ means "orthogonality" in the G. Birkhoff [5] sense (i.e. $\|x+\lambda y\| \geq \|x\|$ for every $\lambda \in \mathbb{R}$).

As D. Amir pointed out [6], implicit in the proof of the above result is that it remains valid if Birkhoff-orthogonality is replaced by any other relation R in E that likewise "admits diagonals" (i.e. such that for every $x, y \in E \setminus \{0\}$ there exists $\alpha > 0$ such that $x+\alpha y R x-\alpha y$) [7]. This happens, for example, if \perp denotes any of the orthogonalities Carlsson [8,9], Diminnie [10,11] and Area [12,13]. (Recall that an

orthogonality in a normed linear space is a relation that coincides with the usual orthogonality when the norm is induced by an inner product.)

It is known [5, 2, 14, 15] that if $\dim E \geq 3$ the symmetry of Birkhoff-orthogonality is an inner product space characteristic, and that there exist two-dimensional non inner product spaces (characterized by M.M. Day [2]) with this property satisfied. Nevertheless, it was proved also in [4] that if E is a two-dimensional normed linear space with symmetric B-orthogonality that satisfies a "square inequality" then E is an inner product space. This can be expressed in the following way:

$$E \text{ is an i.p.s.} \Leftrightarrow [x, y \in S, x \perp B y \Rightarrow y \perp B x \text{ and } H(x, y) \leq 0].$$

However, there exist non-inner product spaces with symmetric B-orthogonality and $H(x, y) \geq 0$ for every $x, y \in S$, $x \perp B y$ (i.e. \mathbb{R}^2 endowed with a norm whose spheres are regular hexagons [4]). It is an open question whether the square inequality (\leq) alone is characteristic of inner product spaces.

As a particular case of a more general result, D.A. Senechal [16] proved that if $\epsilon > 0$ then

$$E \text{ is an i.p.s.} \Leftrightarrow [x, y \in S, \|x - y\| < \epsilon \Rightarrow H(x, y) \sim 0].$$

Pursuing this line further, and taking into account that the relation " $x R y$ iff $\|x - y\| = \epsilon$ " admits diagonals, we can say that if $\epsilon > 0$ then

$$E \text{ is an i.p.s.} \Leftrightarrow [x, y \in S, \|x - y\| = \epsilon \Rightarrow H(x, y) \sim 0].$$

In the next proposition we shall see that, for particular values of ϵ , these preceding two characterizations can be improved.

PROPOSITION. Let $D = \{2\cos \frac{k\pi}{2n} : n=2, 3, \dots, k=1, 2, \dots, n-1\} \subset [0, 2]$. If $\epsilon \in (0, 2) \setminus D$ then

$$E \text{ is an i.p.s.} \Leftrightarrow [x, y \in S, \|x - y\| = \epsilon \Rightarrow H(x, y) \sim 0],$$

where \sim denotes either \geq or \leq .

PROOF. It is proved in [17] that if $\epsilon \in (0, 2) \setminus D$ then only the inner product spaces satisfy the property

$$Q_\epsilon: \quad x, y \in S, \|x - y\| = \epsilon \Rightarrow H(x, y) = 0.$$

On the other hand, G. Nordlander [18] proved that if $S_{\mathbb{R}^2}$ is the unit sphere of a norm in \mathbb{R}^2 and $0 < \epsilon \leq 2$ then the set $\{x + y : x, y \in S_{\mathbb{R}^2}, \|x - y\| = \epsilon\}$ is a simple closed curve that bounds an area $(4 - \epsilon^2)$ -times the area bounded by $S_{\mathbb{R}^2}$. Thus, any normed linear space that satisfies the property

$$x, y \in S, \|x - y\| = \epsilon \Rightarrow H(x, y) \sim 0$$

must also satisfy Q_ϵ .

However, for $\epsilon \in D$ there exist non-inner product spaces that satisfy Q_ϵ (i.e. if E is the vector space \mathbb{R}^2 endowed with a norm whose spheres are $4n$ -gons then it satisfies Q_ϵ for $\epsilon = 2\cos \frac{k\pi}{2n}$, $k=1, 2, \dots, n-1$) [17].

In an extensive paper J. Oman [19] proved that, for two-dimensional normed linear spaces, one of the vectors in the parallelogram inequality (\leq) can be fixed. That is, if $\dim E = 2$ then

$$E \text{ is an i.p.s.} \Leftrightarrow \text{there exists an } x \in E \text{ such that } H(x, y) \leq 0 \text{ for every } y \in E.$$

However, this result is not true for (\geq). If E is \mathbb{R}^2 endowed with the norm $\|(x_1, x_2)\| = |x_1| + |x_2|$ and $x = (1, 0)$ then $H(x, y) \geq 0$ for every $y \in E$ [19].

It is essential for the Oman characterization that $\dim E=2$. If, for example, we take E to be the vector space \mathbb{R}^3 endowed with the norm whose unit ball is the set of points (x_1, x_2, x_3) defined by the intersection of the two circular cylinders $x_1^2+x_2^2\leq 1$, $x_2^2+x_3^2\leq 1$, then any plane containing the origin and the point $x=(0,0,1)$ intersects the unit sphere in an ellipse. Therefore, $H(x,y)=0$ for every $y\in E$. (Similar results for dimensions greater than two can be found in [19].)

It is an open question whether one vector can also be fixed in the rhombus inequality (\leq).

With regard to norm inequalities of higher degree, Yang Cong-Ren [20] proved that if $p>2$ then

$$E \text{ is an i.p.s.} \Leftrightarrow \|x+y\|^p + \|x-y\|^p \geq 2(\|x\|^p + \|y\|^p) \text{ for every } x, y \in E.$$

(This inequality does not hold for $p<2$. Similar inequalities for this case have been obtained in [20].)

D. Delbosco [21] proved that if there exists $p>0$ such that $\|x+y\|^p + \|x-y\|^p = 2^p$ for every $x, y \in S$, then E must be an inner product space. Therefore, taking into account that if $x, y \in S$ are orthogonal then $\|x \pm y\| = 2^{1/2}$, it follows that $p=2$. Moreover, taking $\epsilon = 2^{1/2}$ in the Nordlander sets defined above, it follows that in any two-dimensional subspace of any normed linear space there always exist $x, y \in S$ such that $\|x+y\| = \|x-y\| = 2^{1/2}$. Therefore, Delbosco's identity holds only for $p=2$.

Finally, we shall consider the T.M. Rassias [22] characterization:

$$E \text{ is an i.p.s.} \Leftrightarrow \|x+y\|^p + \|x-y\|^p \leq 2^{p-1}(\|x\|^p + \|y\|^p) \text{ for every } x, y \in E \text{ and } p>2.$$

Rassias's proof of this result is based on making p tend to 2 in order to obtain the rhombus inequality. Some observations are suggested by this characterization:

Firstly, from the arguments given above, Rassias's inequality does not hold if we change " \leq " into " \geq ".

Secondly, we cannot fix p . For any $p>2$ the spaces ℓ^p and L^p both satisfy the inequality

$$\|x+y\|_p^p + \|x-y\|_p^p \leq 2^{p-1}(\|x\|_p^p + \|y\|_p^p)$$

(see e.g. Köthe [23], p.356). Furthermore, we shall see that the Rassias characterization does not hold if the inequality holds for every $p>r$ for some $r>2$. This proves that the hypothesis on p in the Rassias characterization of inner product spaces cannot be weakened.

Taking into account that, for every $c>0$, the functions $f(x) = (1+cx)^{1/x}$ and $g(x) = \left[\frac{1+c^x}{2}\right]^{1/x}$, $x>0$, are, respectively, monotone decreasing and monotone increasing, it is easy to see that for every $a>0$, $b>0$, $0<r\leq p$, the following two inequalities hold:

$$(a^p + b^p)^{1/p} \leq (a^r + b^r)^{1/r}$$

$$\left[\frac{a^p + b^p}{2}\right]^{1/p} \geq \left[\frac{a^r + b^r}{2}\right]^{1/r}.$$

Bearing this in mind, we have therefore proved that for $p>r>2$ and every x, y in ℓ^r or L^r , according to the case,

$$\begin{aligned} (\|x+y\|_r^p + \|x-y\|_r^p)^{1/p} &\leq (\|x+y\|_r^r + \|x-y\|_r^r)^{1/r} \leq \\ &\leq 2^{1-1/r} (\|x\|_r^r + \|y\|_r^r)^{1/r} \leq 2^{1-1/p} (\|x\|_r^p + \|y\|_r^p)^{1/p}. \end{aligned}$$

Hence

$$\|x+y\|_r^p + \|x-y\|_r^p \leq 2^{p-1} (\|x\|_r^p + \|y\|_r^p).$$

ACKNOWLEDGEMENT. I am grateful to T.M. Rassias for his valuable suggestions.

REFERENCES

1. JORDAN, P., VON NEUMANN, J. On inner products in linear metric spaces, Ann. of Math. **36** (1935), 719–723.
2. DAY, M.M. Some characterizations of inner product spaces, Trans. Amer. Math. Soc. **62** (1947), 320–337.
3. SCHOENBERG, I.J. A remark on M.M. Day's characterization of inner-product spaces and a conjecture of L.M. Blumenthal, Proc. Amer. Math. Soc. **3** (1952), 961–964.
4. BENITEZ, C., DEL RIO, M. Characterization of inner product spaces through rectangle and square inequalities, Rev. Roumaine Math. Pures Appli. **29** (1984), 543–546.
5. BIRKHOFF, G. Orthogonality in linear metric spaces, Duke Math. J. **1** (1935), 169–172.
6. AMIR, D. Characterizations of Inner Product Spaces, Birkhäuser, 1986.
7. BENITEZ, C. Una propiedad de la ortogonalidad de Birkhoff y una caracterización de espacios prehilbertianos, Collect. Math. **26** (1975), 211–218.
8. CARLSSON, S.O. Orthogonality in normed linear spaces, Ark. Mat. **4** (1962), 297–318.
9. DEL RIO, M. Ortogonalidad en espacios normados y caracterización de espacios prehilbertianos, Dept. Anal. Matem. Univ. de Santiago de Compostela, Spain, serie B, 14, 1975.
10. DIMINNIE, C.R. A new orthogonality relation in normed linear spaces, Math. Nachr. **114** (1983), 197–203.
11. ALONSO, J., BENITEZ, C. Complements on Diminnie orthogonality in normed linear spaces, (preprint).
12. ALONSO, J. Ortogonalidad en Espacios Normados, Ph.D. Thesis, Univ. Extremadura, Badajoz, Spain, 1984.
13. ALONSO, J., BENITEZ, C. Area orthogonality in normed linear spaces, (preprint).
14. JAMES, R.C. Orthogonality and linear functionals in normed linear spaces, Trans. Amer. Math. Soc. **61** (1947), 265–292.
15. JAMES, R.C. Inner products in normed linear spaces, Bull Amer. Math. Soc. **53** (1947), 559–566.
16. SENECHALLE, D.A. Euclidean and non-Euclidean norms in a plane, Illinois Jour. Math. **18** (1971), 281–289.
17. ALONSO, J., BENITEZ, C. Some characteristic and non-characteristic properties of inner product spaces, J. Approx. Theory **55** (1988), 318–325.
18. NORDLANDER, G. The modulus of convexity in normed linear spaces, Arkiv für Matem. **4** (1958), 15–17.
19. OMAN, J. Norm identities which characterize inner product spaces, In The Geometry of Metric and Linear Spaces (Ed. A. Dold and B. Eckmann), Springer Lecture Notes no. 490, 116–133, Michigan 1974.
20. YANG CONG-REN Characterization of inner product spaces, Shuxue Jinzhan **6** (1963), 197–200.
21. DELBOSCO, D. Spazi normati con prodotto interno, Rend. Sem. Mat. Univ. Torino **36** (1977/78), 415–419.
22. RASSIAS, T.M. New characterizations of inner product spaces, Bull Sci. Math. (2) **108** (1984), 95–99.
23. KÖTHE, G. Topological Vector Spaces I (2nd ed.), Springer-Verlag, Berlin, Heidelberg, New York, 1983.

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru