

TRANSIENT DEVELOPMENT OF GRAVITY WAVES FOR TWO LAYERED FLUIDS

A.H. ESSAWY and M.S. FALTAS

Department of Mathematics
University of Bahrain
Isa Town, P.O. Box 32038
State of Bahrain

(Received December 11, 1989)

ABSTRACT. The transient gravity waves generated by a harmonically oscillating wave maker immersed in two incompressible fluids, the upper fluid having a free surface, is considered. The resulting linearized initial value problem is solved using the method of generalized functions, and asymptotic analysis for large time and distance are given for the elevation.

KEY WORDS AND PHRASES. Internal waves, transient waves.

1991 AMS SUBJECT CLASSIFICATION CODES. 76B15, 76C10.

1. INTRODUCTION.

The two-dimensional problem of gravity waves generated by moving oscillating surface pressure distributions in a fluid which is unbounded in both horizontal directions has been studied by Kaplan [1] and Debnath and Rosenblat [2] in infinite depth and in finite depth respectively. Pramanik [3] considered the initial value problem of waves generated by a moving oscillating surface pressure against a vertical cliff and a uniform asymptotic analysis was given for the unsteady case. Debnath and Basu [4] treated the same problem taking into account the effect of surface tension. Faltas [5] investigated the initial value problem of surface waves generated by a harmonically oscillating vertical wave maker immersed in an infinite incompressible fluid of finite constant depth. It is the purpose of this paper to discuss the transient development of two-dimensional linearized waves at the free surface and at the interface between two fluids. The waves are produced by a harmonically oscillating wave maker immersed vertically in both fluids. The integral representations of free and interface elevations are obtained through an application of the Laplace and the generalized cosine Fourier transforms of the equations of motion. Then the application of the stationary phase method combined with the contour integration method leads to the asymptotic waves valid for large time and distance.

2. FORMULATION AND SOLUTION OF THE PROBLEMS.

We are concerned with the transient development of two dimensional infinitesimal wave motion of two superimposed immiscible non-viscous and incompressible fluids separated by a common interface, where the upper fluid has a free surface. The waves are generated by a harmonically oscillating wave maker immersed vertically in the two fluids.

Take the origin O at the mean level to the interface and the axis Oy to be vertically downwards along the wave maker. The upper fluid is of finite constant height with mean level at

$y = -h$, while the lower fluid has infinite depth. If the motion is generated originally from rest by the oscillations of the wavemaker, it will be irrotational throughout all time and we may describe the motion by velocity $0 < y < \infty$, and $0 < x < \infty$, $-h < y < 0$ of the lower and upper fluids respectively. The unsteady motion is produced in the two fluids by the continuous oscillations of the wave maker. Let it oscillate horizontally with velocity $U(y, t)$ given by

$$U(y, t) = u(y)e^{i\omega t}H(t) \quad (2.1)$$

where $u(y)$ is an arbitrary function of y , ω is the frequency, and $H(t)$ is the unit step function. The functions ϕ_j satisfy an initial boundary value problem in which

$$\nabla^2 \phi_j = 0. \quad (2.2)$$

Neglecting surface tension, the linearized pressure and kinematical boundary conditions at the interface of the two fluids are respectively

$$\left. \begin{array}{l} \frac{\partial \phi_1}{\partial t} - s \frac{\partial \phi_2}{\partial t} = (1-s)g\eta_2 \\ \frac{\partial \eta_1}{\partial t} = \frac{\partial \phi_1}{\partial y} = \frac{\partial \phi_2}{\partial t} \end{array} \right\} y = 0 \quad (2.3)$$

while the corresponding conditions at the free surface of the upper fluid are

$$\left. \begin{array}{l} \frac{\partial \phi_2}{\partial t} = g\eta_2 \\ \frac{\partial \eta_2}{\partial t} = \frac{\partial \phi_2}{\partial y} \end{array} \right\} y = -h \quad (2.4)$$

where $s(0 < s < 1)$ is the ratio of the densities of the upper and lower fluids and $n_j = n_j(x, t)$ are the wave elevations associated with the lower and upper fluids. Also,

$$\frac{\partial \phi_1}{\partial y} \rightarrow 0 \quad \text{as } y \rightarrow \infty. \quad (2.5)$$

At the wave maker

$$\frac{\partial \phi_j}{\partial y} = U(y, t) \quad \text{on } x = 0 \quad (2.6)$$

and the initial conditions are

$$\phi_j = \eta_j = 0 \quad \text{when } t = 0. \quad (2.7)$$

We suppose also that ϕ_j, η_j are treated as the generalized function in the sense of Lighthill [6].

We introduce the Fourier cosine transform with respect to x and the Laplace transform with respect to t as

$$\bar{F}_c(k, y, r) = \sqrt{\frac{2}{\pi}} \int_0^\infty \cos kx dx \int_0^\infty e^{-rt} F(x, y, t) dt \quad (2.8)$$

where the suffix c and the bar in the transformed function refer to the cosine Fourier and Laplace transform respectively. Application of (2.8) to the system (2.2) - (2.7) gives

$$\frac{d^2}{dy^2} \bar{\phi}_{jc} - k^2 \bar{\phi}_{jc} = \sqrt{\frac{2}{\pi}} \bar{U}(y, t), \quad r > 0 \quad (2.9a, b)$$

$$\left. \begin{array}{l} \bar{\phi}_{1c} - s \bar{\phi}_{2c} = \frac{g}{r} (1-s) \bar{\eta}_{1c} \\ \frac{d}{dy} \bar{\phi}_{1c} = \frac{d}{y} \bar{\phi}_{2c} = r \bar{\eta}_{1c} \end{array} \right\} \text{on } y = 0, r > 0 \quad (2.10)$$

$$\left. \begin{array}{l} \bar{\phi}_{2c} = \frac{g}{r} \bar{\eta}_{2c} \\ \frac{d}{dy} \bar{\phi}_{2c} = r \bar{\eta}_{2c} \end{array} \right\} \text{on } y = -h, r > 0 \quad (2.11)$$

$$\frac{d}{dy} \phi_{1c} \rightarrow 0 \quad \text{as } y \rightarrow \infty \quad (2.12)$$

The solutions of (2.9a) and (2.9b) satisfying condition (2.1) are respectively

$$\bar{\phi}_{1c} = A(k, r)e^{-ky} - \frac{1}{2} \sqrt{\frac{2}{\pi}} \int_0^\infty k^{-1} e^{k(y-z)} \bar{U}(z, r) dz + \frac{2}{\pi} \int_0^\infty k^{-1} \sinh k(y-z) \bar{U}(z, r) dz, \quad (2.13)$$

$$\bar{\phi}_{2c} = B(k, r)e^{ky} + C(k, r)e^{-ky} + \sqrt{\frac{2}{\pi}} \int_0^y k^{-1} \sinh k(y-z) \bar{U}(z, r) dz, \quad (2.14)$$

where $A(k, r)$, $B(k, r)$ and $C(k, r)$ are functions to be determined. The transformed boundary conditions (2.10) are satisfied if

$$\begin{aligned} A &= -\frac{1}{2} \sqrt{\frac{2}{\pi}} \int_0^\infty \frac{e^{-kz}}{k} \bar{U} dz - \frac{r}{k} \bar{\eta}_{1c}, \\ B &= -\frac{1}{2s} \sqrt{\frac{2}{\pi}} \int_0^\infty \frac{e^{-kz}}{k} \bar{U} dz - \frac{(1-s)(r^2 + gk)}{2skr} \bar{\eta}_{1c}, \\ C &= -\frac{1}{2s} \sqrt{\frac{2}{\pi}} \int_0^\infty \frac{e^{-kz}}{k} \bar{U} dz - \frac{(1+s)r^2 + gk(1-s)}{2skr} \bar{\eta}_{1c} \end{aligned} \quad (2.15)$$

We are interested in the waves after a large time and large distance. To investigate the principal feature of the wave motion it suffices to work only with the elevation η_j . From (2.11), (2.13) - (2.15) we get

$$\begin{aligned} \sqrt{\frac{\pi}{2}} \bar{\eta}_{1c} &= \frac{-rc}{(r^2 + m_1^2)(r^2 + m_2^2)} \left\{ r^2 \left[\cosh kh \int_0^\infty e^{-kz} \bar{U} dz + s \int_0^{-h} \sinh k(h+z) \bar{U} dz \right] \right. \\ &\quad \left. + gk \left[\sinh k \int_0^\infty e^{-kz} \bar{U} dz + s \int_0^{-h} \cosh kh(h+z) \bar{U} dz \right] \right\} \end{aligned} \quad (2.16)$$

$$\begin{aligned} \sqrt{\frac{\pi}{2}} \bar{\eta}_{2c} &= \frac{-rc}{(r^2 + m_1^2)(r^2 + m_2^2)} \left\{ r^2 \left[\int_0^\infty e^{-kz} \bar{U} dz - \int_0^{-h} \cosh kz + s \sinh kz \bar{U} dz \right] \right. \\ &\quad \left. - gk(1-s) \int_0^{-h} \cosh kz \bar{U} dz \right\}, \end{aligned} \quad (2.17)$$

where $1/c = \cosh kh + s \sinh k$ and $m_1^2(k) = gk$, $m_2^2(k) = \frac{(1-s)gk}{s + \coth kh}$. Assuming the particular form of $U(y, t)$ as given by (2.1), the inverse of Laplace and cosine Fourier transforms with the convolution theorem for Laplace transform give

$$\eta_1(x, t) = -\frac{2}{\pi} \sum_{p=1}^2 \int_0^\infty \beta_p(k) \cos kx dk \int_0^t e^{i\omega t} \cos m_p(t-\tau) d\tau, \quad (2.18)$$

$$\eta_2(x, t) = -\frac{2}{\pi} \sum_{p=1}^2 \int_0^\infty \gamma_p(k) \cos kx dk \int_0^t e^{i\omega t} \cos m_p(t-\tau) d\tau, \quad (2.19)$$

where

$$\begin{aligned}
\beta_1(k) &= \frac{c m_1^2}{(m_2^2 - m_1^2)} \left[s \int_0^{-h} e^{-k(h+z)} u(z) dz - \int_0^{\infty} e^{-k(h+z)} u(z) dz \right], \\
\beta_2(k) &= \frac{c}{(m_2^2 - m_1^2)} \left[s \int_0^{-h} (m_2^2 \sinh k(h+z) - m_1^2 \cosh k(h+z)) u(z) dz \right. \\
&\quad \left. + (m_2^2 \cosh kh - m_1^2 \sinh kh) \int_0^{\infty} e^{-kz} u(z) dz \right], \\
\gamma_1(k) &= \frac{c m_1^2}{(m_2^2 - m_1^2)} \left[s \int_0^{-h} e^{kz} u(z) dz - \int_0^{\infty} e^{-kz} u(z) dz \right], \\
\gamma_2(k) &= \frac{-c}{(m_2^2 - m_1^2)} \left[s \int_0^{-h} (m_2^2 \sinh kz + m_1^2 \cosh kz + (m_2^2 - m_1^2) \cosh kz) u(z) dz - m_2^2 \int_0^{\infty} e^{-kz} u(z) dz \right].
\end{aligned}$$

Carrying out the integral in (2.18) or (2.19), we get,

$$\eta_1(x, t) = \frac{2}{\pi} \sum_{p=1}^2 \int_0^{\infty} \frac{\beta_p(k)}{m_p^2 - \omega^2} [i\omega \cos m_p t - m \sin m_p t - i\omega e^{i\omega t}] \cos kx dk, \quad (2.20)$$

and a similar expression for $\eta_2(x, t)$.

3. ASYMPTOTIC ANALYSIS OF SOLUTION.

To evaluate the integral (2.20) or the corresponding one for η_2 for large values of x and t we shall use formulae developed by Lighthill [6] and Jones [7]. Write $\eta_1 = I + I' + J + J'$, where

$$I = \frac{-2}{\pi} i\omega e^{i\omega t} \int_0^{\infty} \frac{\beta_1(k)}{m_1^2 - \omega^2} \cos kx dk, \quad I' = \frac{-2i\omega}{\pi} e^{i\omega t} \int_0^{\infty} \frac{\beta_2(k)}{m_2^2 - \omega^2} \cos kx dk \quad (3.1)$$

$$\left. \begin{aligned}
J &= \frac{2}{\pi} \int_0^{\infty} \frac{\beta_1(k)}{m_1^2 - \omega^2} (i\omega \cos m_1 t - m_1 \sin m_1 t) \cos kx dk, \\
J' &= \frac{2}{\pi} \int_0^{\infty} \frac{\beta_1(k)}{m_2^2 - \omega^2} (i\omega \cos m_2 t - m_2 \sin m_2 t) \cos kx dk
\end{aligned} \right\} \quad (3.2)$$

The first two integrals (3.1), represent the steady state solution while the second two (3.2) represent the transient solution. It is convenient to rewrite (3.1), (3.2) as follows

$$I = \frac{i}{2\pi} e^{i\omega t} \sum_{n=1}^2 I_n, \quad I' = \frac{i}{2\pi} e^{i\omega t} \sum_{n=1}^2 I'_n, \quad J = \frac{i}{2\pi} \sum_{n=1}^4 J_n, \quad J' = \frac{i}{2\pi} \sum_{n=1}^4 J'_n$$

where

$$I_1, I_2 = \pm \int_0^{\infty} \frac{\beta_1(k)}{m_1 \mp \omega} (e^{ikx} + e^{-ikx}) dk, \quad J_1, J_2 = \int_0^{\infty} \frac{\beta_1(k)}{m_1 \mp \omega} e^{i(\omega t \pm kx)} dk$$

$$J_3, J_4 = - \int_0^{\infty} \frac{\beta_1(k)}{m_1 + \omega} e^{i(\omega t \mp kx)} dk, \quad I_1, I'_2 = \mp \int_0^{\infty} \frac{\beta_2(k)}{m_2 \mp \omega} (e^{ikx} + e^{-ikx}) dk,$$

$$J'_1, J'_2 = \int_0^{\infty} \frac{\beta_2(k)}{m_2 \pm \omega} e^{i(\omega t \pm kx)} dk, \quad J'_3, J'_4 = - \int_0^{\infty} \frac{\beta_2(k)}{m_2 + \omega} e^{i(\omega t \mp kx)} dk.$$

We follow the method of Debnath and Rosenblat [2] to evaluate these wave integrals. The main contribution to the asymptotic value of the above integrals comes from the poles and stationary points of the integrals. It is noted that each I_1, J_1 , and J_2 contains one pole at $k = k_0$ where $k_0 = \omega^2/g$, and each of I'_1, J'_2, J'_2 contains one pole at $k = k'_0$, where k'_0 is the only real positive root of the equation

$$\sqrt{\frac{(1-s)gk}{s + \coth kh}} = \omega.$$

In addition, the integrals J_2, J_3 contain one stationary point at $k = k_1$, which is the root of the equations

$$\frac{dm_1}{dk} = \frac{x}{t} \quad \text{i.e., } k_1 = \frac{gt^2}{4x^2},$$

also, the integrals J'_2, J'_1 contain one stationary point at $k = k'_1$ which is the root of the equation

$$\frac{dm_2}{dk} = \frac{x}{t}. \quad (3.3)$$

We note that

$$\begin{aligned} \frac{d^2m}{dk^2} = \frac{m(k)}{4k^2} & (s \sinh 2kh + \cosh 2kh - 1)^{-1} [(4h^2k^2 - \sinh^2 2kh) \\ & + 4kh(\sinh 2kh - 2kh \cosh 2kh - s^2(\cosh 2kh - 1)^2) \\ & + s(-8h^2k^2 \sinh 2kh + 2(2kh - \sinh^2 2kh)(\cosh 2kh - 1))] < 0. \end{aligned}$$

Therefore dm/dk decreases monotonically from $\sqrt{gh(1-s)}$ to 0 as k varies from 0 to ∞ . Hence equation (3.3) has only one real root k'_1 . On the other hand, the integrals I_2, J_4, I'_2 and I'_4 contains neither poles nor stationary points in the range of integration.

Now the contribution from the poles k_0, k'_0 can be evaluated using the formula (24) for the asymptotic development as stated by Debnath and Rosenblat [2]. It then follows that as $x \rightarrow \infty$,

$$I \sim \frac{\beta_1(k_0)}{2m'_1(k_0)} e^{i\omega t} (e^{ik_0 x} - e^{-ik_0 x}), \quad I' \sim \frac{\beta_2(k'_0)}{2m'_2(k'_0)} e^{i\omega t} (e^{ik'_0 x} - e^{-ik'_0 x}) \quad (3.4ab)$$

where $m'_1(k_0), m'_2(k'_0)$ are the derivatives of $m_1(k)$ at $k = k'_0$ and $m_2(k)$ at $k = k'_0$ respectively.

The method of stationary phase (Jones [7]) can be used to evaluate the transient component of J (that is the contribution from the stationary points)

$$J_{tr} \sim \frac{i}{2\pi} \beta_1(k_1) \sqrt{\frac{2\pi}{t |m''_1(k_1)|}} \left[\frac{\exp[i\{tm_1(k_1) - k_1 x - \frac{\pi}{4}\}]}{m_1(k_1) + \omega} - \frac{\exp[-i\{tm_1(k_1) - k_1 x - \frac{\pi}{4}\}]}{m_1(k_1) + \omega} \right] \quad (3.5)$$

$$J'_{tr} \sim \frac{i}{2\pi} \beta_1(k'_1) \sqrt{\frac{2\pi}{t |m''_2(k'_1)|}} \left[\frac{\exp[i\{tm_2(k'_1) - k'_1 x - \pi/4\}]}{m_2(k'_1) - \omega} - \frac{\exp[i\{tm_2(k'_1) - k'_1 x - \pi/4\}]}{m_2(k'_1) + \omega} \right] + 0(1/t), \quad (3.6)$$

where J_{tr}, J'_{tr} denote the transient parts of J and J' respectively for large t .

Finally we calculate the contribution to J and J' from their polar singularity. This can be easily estimated by formula (24), as stated in Debnath and Rosenblat [2].

$$J_{\text{polar}} \sim -\frac{\beta_1(k_0)}{2m'_1(k_0)} e^{i\omega t} (e^{ik_0 x} + e^{-ik_0 x}), \quad J'_{\text{polar}} \sim -\frac{\beta_2(k_0)}{2m'_2(k'_0)} e^{i\omega t} (e^{ik'_0 x} + e^{-ik'_0 x}). \quad (3.7ab)$$

We write $\eta_1 = \eta_{st} + \eta'_{st} + \eta_{tr} + \eta'_{tr}$ where η_{st}, η'_{st} are the steady state components of η_1 and η_{tr}, η'_{tr} are the transient components. The first term in η_1 , is the polar contribution to I and J and the

second term is the polar contribution to I' and J' which are given by

$$\eta_{st} = -\frac{\beta_1(k_0)}{m'_1(k_0)} e^{i(\omega t - k_0 x)} + 0\left(\frac{1}{x}\right), \quad \eta'_{st} = -\frac{\beta_2(k'_0)}{m'_2(k'_0)} e^{i(\omega t - k'_0 x)} + 0\left(\frac{1}{x}\right) \quad (3.8ab)$$

and the transient components η_{tr} , η'_{tr} are given respectively by (3.5) and (3.6).

So far the entire analysis of the asymptotic behavior has been carried out for $\eta_1(x, t)$. A similar asymptotic analysis can be obtained for $\eta_2(x, t)$. It is clear that there are two modes of waves spreading at each of the free surface of the upper fluid and in the interface of the two fluids and of course one of them will dominate on the other. The above analysis reveals the fact that the transient solution decays rapidly to zero as time $t \rightarrow \infty$. The ultimate steady state is established in the limit. Solutions (3.8ab) represent outgoing waves propagating with phase velocity ω/k_0 and ω/k'_0 respectively. These results justify the use by previous authors of the condition at infinity known as the Sommerfield radiation condition when investigating steady-state harmonic surface waves problem. The application of this condition instead of the boundedness condition at infinity was necessary to render the solution unique.

REFERENCES

1. KAPLAN, P., The waves generated by the forward motion of oscillatory pressure distributions, Prof. 5th Midwest Conf. Fluid Mech. (1957), 316.
2. DEBNATH, L. and ROSENBLAT, S., The ultimate approach to the steady states in the generation of waves on a running stream, Quart. J. Mech. Appl. Math. **22** (1969), 221.
3. PRAMANIK, A.K., Waves created against a verticae cliff a uniform asymptotic solution, Math. Proc. Camb. Phil. Soc. **83** (1978), 321.
4. DEBNATH, K. and BASU, U., Capillary-gravity waves against a vertical cliff, Ind. J. Math. **26** (1984), 49.
5. FALTAS, M.S., Asymptotic analysis of surface waves due to oscillatory wave maker, Quart. Appl. Math. **46** (1988), No. 3, 489.
6. LIGHTHILL, M.J., Fourier Analysis and Generalized Functions, Cambridge University Press, Cambridge (1962).
7. JONES, D.S., Generalized Functions, McGraw-Hill, 1966.

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru