

**ANALYTIC SOLUTION OF RICCATI EQUATIONS OCCURRING  
IN OPEN-LOOP NASH MULTIPLAYER DIFFERENTIAL GAMES**

L. JÓDAR and E. NAVARRO

Departamento de Matemática Aplicada  
Universidad Politécnica de Valencia  
E.T.S.I.I., P.O. Box 22.012  
Valencia, Spain

(Received January 2, 1991)

**ABSTRACT.** In this paper we present explicit analytic solutions of coupled Riccati matrix differential systems appearing in open-loop Nash games. Two different cases are considered. Firstly, by means of appropriate algebraic transformations the problem is decoupled so that an explicit solution of the problem is available. The second is based on the existence of a solution of a rectangular Riccati type algebraic matrix equation associated with the problem.

**KEY WORDS AND PHRASES.** Riccati equations, differential games, optimization problem.

**1991 AMS SUBJECT CLASSIFICATION CODE.** 34A34

**1. INTRODUCTION.** When noncooperative problems are tackled, a game theoretic approach is necessary: each control agent (decision maker or player) tries to optimize his own cost function which conflicts more or less with others. An equilibrium solution must be sought, and the Nash strategy is a natural choice. In this case, a player cannot improve his payoff by deviating unilaterally from his Nash strategy. Due to this noncooperation, the optimization problem of various players are strongly coupled and necessary conditions for open-loop Nash strategy lead to complex two-point boundary value problems.

Consider a  $N$ -players linear quadratic differential game defined by

$$x' = Ax + \sum_{i=1}^N B_i u_i; \quad x(0) = x_0 \quad (1.1)$$

with the cost functionals associated with the players

$$J_i = \frac{1}{2} \left\{ x_f^T K_{if} x_f + \int_0^{t_f} (x^T Q_i x + \sum_{j=1}^N u_j^T R_{ij} u_j) dt \right\}; \quad x(t_f) = x_f \quad (1.2)$$

where all matrices are  $n \times n$  symmetric with  $R_{ii}$ , for  $1 \leq i \leq N$ , positive definite. It is well known that the open-loop Nash control must satisfy [12]:

$$u_i = -R_{ii}^{-1} B_i^T \Psi_i; \quad \Psi_i' = -Q_i x - A^T \Psi_i, \quad \Psi_i(t_f) = K_{if} x_f, \quad 1 \leq i \leq N \quad (1.3)$$

where  $\Psi_i$  is the costate vector associated with player "i". When the transformation  $\Psi_i = K_i x$ , is introduced, for  $1 \leq i \leq N$ , the open-loop Nash strategy  $(u_i^*)_{i=1}^N$  is given by

$$u_i^* = -R_{ii}^{-1} B_i^T K_i(t) \Phi(t, 0) x_0, \quad 1 \leq i \leq N \quad (1.4)$$

where

$$K'_i = -A^T K_i - K_i A - Q_i + K_i \sum_{j=1}^N S_j K_j; \quad K_i(t_f) = K_{if}, \quad 1 \leq i \leq N \quad (1.5)$$

with

$$S_i = B_i R_{ii}^{-1} B_i^T, \quad 1 \leq i \leq N \quad (1.6)$$

and  $\Phi(t, 0)$  is the system's transition matrix satisfying

$$\Phi'(t, 0) = (A - \sum_{j=1}^N S_j K_j) \Phi(t, 0); \quad \Phi(t, t) = I \quad (1.7)$$

Note that the matrices  $R_{ij}$  for  $i \neq j$  do not appear in the necessary conditions due to the fact that under open-loop strategy assumptions, each player optimizes his criterion knowing that  $\partial u_i / \partial x = 0$ , for  $1 \leq i \leq N$ . For the open-loop Nash strategy and under the existence of a solution of the coupled Riccati system, the optimization problem has only one solution [2].

The solution of system (1.5) is generally difficult to obtain due to the permanent coupling between the players' strategies. In [4] a series solution of system (1.5) is proposed but the coefficient are obtained solving several linear matrix equations. In [14] a numerical algorithm for the integration of (1.5) is given. A singular perturbation method for solving (1.5) is proposed in [10]. For the case  $N = 2$ , an iterative algorithm for solving (1.5) is given in [5]. For the case  $N = 2$  and  $Q_2 = \alpha Q_1$ , where  $\alpha$  is a scalar, an analytic solution of system (1.5) was pointed out in [1]. In this paper we obtain an explicit solution of system (1.5) for a case more general than the one proposed in [1]. Also, a different type condition expressed in terms of the existence of certain coupled algebraic Riccati matrix system is proposed.

## 2. ANALYTIC SOLUTION OF COUPLED RICCATI SYSTEM BASED ON ALGEBRAIC TRANSFORMATIONS.

For convenience, the necessary conditions to be satisfied (1.1), (1.3), are rewritten in a matrix form as

$$\begin{bmatrix} x' \\ \Psi'_1 \\ \Psi'_2 \\ \vdots \\ \Psi'_N \end{bmatrix} = \begin{bmatrix} A & -S_1 & -S_2 & \cdots & -S_N \\ -Q_1 & -A^T & 0 & \cdots & 0 \\ -Q_2 & 0 & -A^T & \cdots & 0 \\ \vdots & & & & \vdots \\ -Q_N & 0 & 0 & \cdots & -A^T \end{bmatrix} \begin{bmatrix} x \\ \Psi_1 \\ \Psi_2 \\ \vdots \\ \Psi_N \end{bmatrix} = M \begin{bmatrix} x \\ \Psi_1 \\ \Psi_2 \\ \vdots \\ \Psi_N \end{bmatrix} \quad (2.1)$$

$$x(0) = x_0, \quad \Psi_i(t_f) = K_{if} x_f, \quad 1 \leq i \leq N \quad (2.2)$$

Now, let us introduce the change of basis

$$\begin{bmatrix} x \\ \Psi_1 \\ \Psi_2 \\ \vdots \\ \Psi_N \end{bmatrix} = T \begin{bmatrix} x \\ \Psi_1 \\ w_2 \\ \vdots \\ w_N \end{bmatrix}; \quad T = \begin{bmatrix} I & 0 & 0 & \cdots & 0 \\ 0 & I & 0 & \cdots & 0 \\ 0 & L_2 & I & \cdots & 0 \\ \vdots & \vdots & & & \vdots \\ 0 & L_N & 0 & \cdots & I \end{bmatrix} \quad (2.3)$$

for appropriate matrices  $L_2, L_3, \dots, L_N$  in  $\mathbb{R}^{n \times n}$  to be determined. Thus problem (2.1), (2.2) is

equivalent to the following one

$$\begin{bmatrix} x' \\ \Psi'_1 \\ w'_2 \\ \vdots \\ w_N \end{bmatrix} = \begin{bmatrix} A & -S_1 - S_2 L_2 - \cdots - S_N L_N & -S_2 & \cdots & -S_N \\ -Q_1 & -A^T & 0 & \cdots & 0 \\ L_2 Q_1 - Q_2 & L_2 A^T - A^T L_2 & -A^T & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ L_N Q_1 - Q_N & L_N A^T - A^T L_N & 0 & \cdots & -A^T \end{bmatrix} \begin{bmatrix} x \\ \Psi_1 \\ w_2 \\ \vdots \\ w_N \end{bmatrix} \quad (2.4)$$

$$x(0) = x_0, \quad \Psi_1(t_f) = K_{1f} x_f, \quad w_j(t_f) = (K_{jf} - L_j K_{1f}) x_f, \quad 2 \leq j \leq N \quad (2.5)$$

The purpose of this transformation is to find under what conditions the player's optimization problem can be decoupled. In fact, note that if  $L_2, L_3, \dots, L_N$  satisfy the system

$$L_j Q_1 = Q_j, \quad L_j A^T = A^T L_j, \quad 2 \leq j \leq N \quad (2.6)$$

the matrix  $T^{-1}MT$  is reduced to a block triangular form and the costate vectors  $\Psi_1, w_2, \dots, w_N$ , are coupled only via the terminal conditions (2.2).

Note that for the case  $N = 2$  and the matrices  $Q_1$  and  $Q_2$  are proportional, i.e.,  $Q_2 = \alpha Q_1$ , for some scalar  $\alpha$ , taking  $L = \alpha I$ , one gets solutions of system (2.6) for  $N = 2$ . Thus the case studied in [1] is a particular case of (2.6).

In order to characterize the existence of solutions for the algebraic system (2.6), we recall the concept of tensor product of matrices. If  $A, B$  are matrices in  $\mathbb{R}^{m \times n}$  and  $\mathbb{R}^{k \times s}$ , respectively, then the tensor product of  $A$  and  $B$ , denoted  $A \otimes B$ , is defined as the partitioned matrix

$$A \otimes B = \begin{bmatrix} a_{11}B & a_{12}B & \cdots & a_{1n}B \\ \vdots & & & \vdots \\ a_{m1}B & a_{m2}B & \cdots & a_{mn}B \end{bmatrix}$$

If  $A \in \mathbb{R}^{m \times n}$ , we denote

$$A_{\cdot j} = \begin{bmatrix} a_{1j} \\ \vdots \\ a_{mj} \end{bmatrix}, \quad 1 \leq j \leq n; \quad \text{vec } M = \begin{bmatrix} M_{\cdot 1} \\ \vdots \\ M_{\cdot n} \end{bmatrix}$$

If  $M, N$  and  $P$  are matrices of suitable dimensions, then using the column lemma [7, p. 410], we get

$$\text{vec}(MNP) = (P^T \otimes M) \text{vec } N \quad (2.7)$$

Taking into account (2.7), if we apply tensor products in each equation of system (2.6), it may be written in the equivalent form

$$C \text{ vec } L_j = \text{vec}[0, Q_j] \quad (2.8)$$

where  $2 \leq j \leq N$  and

$$C = \begin{bmatrix} I \otimes A^T - A \otimes I \\ Q_1 \otimes I \end{bmatrix} \quad (2.9)$$

If we denote by  $C^+$  the Moore-Penrose pseudoinverse of  $C$ , then from Theorem 2.3.2 of [11, p.24],

the system (2.9) is compatible if and only if

$$CC^+ \text{vec}[0, Q_j] = \text{vec}[0, Q_j] \quad (2.10)$$

Furthermore, under condition (2.10), the general solution of (2.9) for  $L_j$  is given by

$$\text{vec } L_j = C^+ \text{vec}[0, Q_2] + (I - C^+ C)z \quad (2.11)$$

where  $I$  denotes the identity matrix in  $\mathbb{R}^{n^2} \times n^2$  and  $z$  is an arbitrary vector in  $\mathbb{R}^{n^2}$ . We recall that  $C^+$  may be computed by using MATLAB [8].

Let us assume the existence of solutions  $L_j$  of (2.8) for  $2 \leq j \leq N$ , then from (2.4), (2.5) it follows that

$$\begin{bmatrix} x' \\ \Psi'_1 \\ w'_2 \\ \vdots \\ w'_N \end{bmatrix} = \begin{bmatrix} V & | & -S_2 & \cdots & -S_N \\ \hline & | & 0 & \cdots & 0 \\ \hline & | & -A^T & \cdots & 0 \\ \hline 0 & | & \vdots & \ddots & \vdots \\ \hline & | & 0 & \cdots & -A^T \end{bmatrix} \begin{bmatrix} x \\ \Psi'_1 \\ w'_2 \\ \vdots \\ w'_N \end{bmatrix}; \quad (2.12)$$

$$x(0) = x_0 \quad \Psi'_1(t_f) = K_{1f}x_f \quad w_j(t_f) = (K_{jf} - L_j K_{1f})x_f \quad 2 \leq j \leq N$$

where

$$V = \begin{bmatrix} A & -S_1 - S_2 L_2 - \cdots - S_N L_N \\ -Q_1 & -A^T \end{bmatrix} \quad (2.13)$$

Let us consider the change  $t = t(s) = t_f - s$ ,  $0 \leq s \leq t_f$ , and let

$$\begin{aligned} \hat{x}(s) &= x(t_f - s) = x(t) \\ \hat{\Psi}_1(s) &= \Psi'_1(t_f - s) = \Psi_1(t) \\ \hat{w}_j(s) &= w_j(t_f - s) = w_j(t), \quad 2 \leq j \leq N \end{aligned} \quad (2.14)$$

Hence problem (2.11) may be written in the form

$$(d/ds) \begin{bmatrix} \hat{x} \\ \hat{\Psi}_1 \\ \hat{w}_2 \\ \vdots \\ \hat{w}_N \end{bmatrix} = \begin{bmatrix} -V & | & -S_2 & \cdots & -S_N \\ \hline & | & 0 & \cdots & 0 \\ \hline & | & -A^T & \cdots & 0 \\ \hline 0 & | & \vdots & \ddots & \vdots \\ \hline & | & 0 & \cdots & -A^T \end{bmatrix} \begin{bmatrix} \hat{x} \\ \hat{\Psi}_1 \\ \hat{w}_2 \\ \vdots \\ \hat{w}_N \end{bmatrix}$$

(2.15)

$$\hat{x}(t_f) = x_o, \quad \hat{\Psi}_1(0) = K_{1f} \hat{x}(0), \quad \hat{w}_j(0) = (K_{jf} - L_j K_{1f}) \hat{x}(0), \quad 2 \leq j \leq N$$

Solving (2.15) we obtain

$$\hat{w}_j(s) = \exp(sA^T) \hat{w}_j(0), \quad 2 \leq j \leq N \quad (2.16)$$

$$\begin{bmatrix} \hat{x}(s) \\ \hat{\Psi}_1(s) \end{bmatrix} = \exp(-sV) \left\{ \begin{bmatrix} \hat{x}(0) \\ \hat{\Psi}_1(0) \end{bmatrix} + \sum_{j=2}^N \int_0^s \exp(uV) \begin{bmatrix} S_j \\ 0 \end{bmatrix} \exp(uA^T) \hat{w}_j(0) du \right\} \quad (2.17)$$

From (2.14) and (2.16) we have

$$\begin{bmatrix} \hat{x}(s) \\ \hat{\Psi}_1(s) \end{bmatrix} = G(s) \hat{x}(0) \quad (2.18)$$

where

$$G(s) = \exp(-sV) \left\{ \begin{bmatrix} I \\ K_{1f} \end{bmatrix} + \sum_{j=2}^N \int_0^s \exp(uV) \begin{bmatrix} S_j \\ 0 \end{bmatrix} \exp(uA^T) du (K_{jf} - L_j K_{1f}) \right\} \quad (2.19)$$

Thus we have

$$\hat{x}(s) = [I, 0] G(s) \hat{x}(0); \quad \hat{\Psi}_1(s) = [0, I] G(s) \hat{x}(0) \quad (2.20)$$

Note that  $[I, 0]G(0) = I$ , and from the continuity of  $G$ , there exists an interval  $0 \leq s \leq \delta$ , such that

$$[I, 0]G(s) \quad \text{is invertible for all } s \in [0, \delta] \quad (2.21)$$

From (2.19) and (2.20) we obtain

$$\begin{aligned} \hat{x}(0) &= \{[I, 0]G(s)\}^{-1} \hat{x}(s) \\ \hat{\Psi}_j(s) &= [0, I]G(s) \{[I, 0]G(s)\}^{-1} \hat{x}(s), \quad 0 \leq s \leq \delta \end{aligned} \quad (2.22)$$

Now, from (2.3) and (2.13), it follows that

$$\hat{\Psi}_j(s) = L_j \hat{\Psi}_1(s) + \hat{w}_j(s), \quad 2 \leq j \leq N$$

and from (2.15) and (2.19)

$$\hat{\Psi}_j(s) = \{\exp(sA^T)(K_{jf} - L_j K_{1f}) + L_j [0, I]G(s)\} \{[I, 0]G(s)\}^{-1} \hat{x}(s) \quad (2.23)$$

for  $0 \leq s \leq \delta$ .

From (2.13) and the relations  $\Psi_j(t) = K_j(t)x(t)$ ,  $1 \leq j \leq N$ , it follows that

$$K_1(t) = [0, I]G(t_f - t) \{[I, 0]G(t_f - t)\}^{-1} \quad (2.24)$$

$$\begin{aligned} K_j(t) &= \{\exp(A^T(t_f - t))(K_{jf} - L_j K_{1f}) + L_j [0, I]G(t_f - t)\} \{[I, 0]G(t_f - t)\}^{-1}, \\ t_f - \delta \leq t \leq t_f; \quad 2 \leq j \leq N \end{aligned} \quad (2.25)$$

where  $G$  is defined by (2.18). Thus the following result has been proved:

**THEOREM 1.** Let us assume that matrices  $A$  and  $Q_i$  for  $1 \leq i \leq N$ , satisfy the condition (2.9) where  $C$  is defined by (2.8), and let  $L_j$  be the solution of (2.8) for  $2 \leq j \leq N$ . Then there exists a positive number  $\delta$  such that on the interval  $[t_f - \delta, t_f]$  the unique solution of the coupled Riccati system (1.5) is given by (2.23), (2.24).

REMARK 1. Note that the case  $N = 2$ ,  $Q_2 = \alpha Q_1$ , where  $\alpha$  is a scalar, is a particular case of the previous theorem taking  $L_2 = \alpha I$ . It is important to note that from (2.23) and (2.24), we have the following relation between  $K_1(t)$  and  $K_2(t)$ :

$$K_2(t) = L_2 K_1(t) + \exp(A^T(t_f - t)) [K_{2f} - L_2 K_{1f}] \{[I, 0] G(t_f - t)\}^{-1}$$

and since the function  $\{[I, 0] G(t_f - t)\}^{-1}$  is involved in the computation of  $K_1(t)$ , the computational cost is reduced because  $K_2(t)$  is expressed in terms of  $K_1(t)$ . Finally we recall that efficient methods for computing matrix exponentials and integrals involving them and that appear in the expression of  $G(s)$ , may be found in [13]. These procedures are extremely easy to implement and yield an estimation of the approximation error.

### 3. ANALYTIC SOLUTION OF COUPLED RICCATI DIFFERENTIAL SYSTEM BASED ON THE EXISTENCE OF A SOLUTION OF A COUPLED ALGEBRAIC RICCATI SYSTEM.

Riccati type matrix equations with rectangular coefficients arise for instance in the problem of finding a state estimate feedback controller [3] and in the transformation of ill-conditioned linear systems to a block diagonal form [6,9]. An efficient method to find solutions of such equations may be found in [15]. The aim of this section is to propose another way to find an analytic solution of the coupled Riccati differential system (1.5). Note that system (1.5) may be written in the following compact form

$$K'(t) = -Q - K(t)A - BK(t) + K(t)SK(t); \quad K(t_f) = K_f \quad (3.1)$$

where

$$K = \begin{bmatrix} K_1 \\ \vdots \\ K_N \end{bmatrix}; \quad Q = \begin{bmatrix} Q_1 \\ \vdots \\ Q_N \end{bmatrix}; \quad B = \text{Diag}(A^T, A^T, \dots, A^T), \quad K_f = \begin{bmatrix} K_{1f} \\ \vdots \\ K_{Nf} \end{bmatrix} \quad (3.2)$$

and

$$S = [S_1, S_2, \dots, S_N] \quad (3.3)$$

Let us assume that the rectangular algebraic Riccati equation

$$-Q - XA - BX + XSX = 0 \quad (3.4)$$

admits a solution  $X_o \in \mathbb{C}^{Nn \times n}$  and let us consider the change

$$U(t) = K(t) \equiv X_o \quad (3.5)$$

Then problem (3.1) is equivalent to the following one

$$U'(t) = B_o U(t) - U(t)A_o + U(t)S U(t); \quad U(t_f) = U_f \quad (3.6)$$

where

$$B_o = X_o S - B; \quad A_o = A - S X_o; \quad U_f = K_f - X_o \quad (3.7)$$

Now, let us consider the extended linear system

$$(d/dt) \begin{bmatrix} V(t) \\ Z(t) \end{bmatrix} = \begin{bmatrix} A_o & -S \\ 0 & B_o \end{bmatrix} \begin{bmatrix} V(t) \\ Z(t) \end{bmatrix}; \quad \begin{bmatrix} V(t_f) \\ Z(t_f) \end{bmatrix} = \begin{bmatrix} I \\ U_f \end{bmatrix} \quad (3.8)$$

where  $V(t) \in \mathbb{C}^{n \times n}$  and  $Z(t) \in \mathbb{C}^{Nn \times n}$ . If we define the matrix function

$$S(t, s) = \begin{bmatrix} \exp((t-s)A_o) & - \int_{t_f}^t \exp((t-v)A_o) S \exp((v-x)B_o) dv \\ 0 & \exp((t-s)B_o) \end{bmatrix} \quad (3.9)$$

then an easy computation yields

$$(\partial/\partial t)S(t, s) = \begin{bmatrix} A_o & -S \\ 0 & B_o \end{bmatrix} S(t, s) \quad (3.10)$$

and thus  $S(t, s)$  is a fundamental matrix of (3.8) and the unique local solution of (3.8) in a neighborhood of  $t_f$  is given by

$$\begin{bmatrix} V(t) \\ Z(t) \end{bmatrix} = S(t, t_f) \begin{bmatrix} I \\ U_f \end{bmatrix}; \quad S(t_f, t_f) = I_{(N+1)n}$$

Note that  $V(t_f) = I_n$  and thus in a neighbourhood  $J$  containing  $t_f$ ,  $V(t)$  is invertible. Now, let us define the  $\mathbb{C}^{Nn \times n}$ -values matrix function  $U(t) = Z(t)(V(t))^{-1}$  for  $t \in J$ . Note that from (3.8) it follows that

$$V'(t) = A_o V(t) - S Z(t) \quad \text{and} \quad Z'(t) = B_o Z(t)$$

Computing it follows that

$$U'(t) = Z'(t)(V(t))^{-1} - Z(t)(V(t))^{-1} V'(t)(V(t))^{-1} = B_o U(t) - U(t)A_o + U(t)S U(t)$$

for all  $t \in J$ . Hence  $K(t) = U(t) + X_o$  is the solution of (3.1), defined on the interval  $J$ . From (3.10) it follows that

$$\begin{aligned} V(t) &= \exp((t-t_f)A_o) - \int_{t_f}^t \exp((t-v)A_o) S \exp((v-t_f)B_o) dv U_f \\ &= \exp((t-t_f)A_o) \left\{ I - \int_{t_f}^t \exp((t_f-v)A_o) S \exp(vB_o) dv \exp(-t_f B_o) U_f \right\} \\ Z(t) &= \exp((t-t_f)B_o) U_f \\ K(t) &= X_o + \exp((t-t_f)B_o) U_f \left\{ I - \int_{t_f}^t \exp((t_f-v)A_o) S \exp(vB_o) dv \exp(-t_f B_o) U_f \right\}^{-1} \exp((t_f-t)A_o) \end{aligned} \quad (3.11)$$

Thus if the algebraic Riccati equation (3.4) has a solution  $X_o$ , then the solution  $K(t)$  of (3.1) is defined by (3.11) where  $U_f$ ,  $A_o$  and  $B_o$  are given by (3.7). The solution is defined in the neighbourhood of  $t_f$  where  $V(t)$  is invertible.

ACKNOWLEDGEMENT. This paper has been partially supported by the D.G.I.C.Y.T. grant PS87-0064 and the NATO grant 900040.

#### REFERENCES

1. ABOU-KANDIL, H. and BERTRAND, P., Analytic solution for a class of linear quadratic open-loop Nash games, *Internat. J. Control.* **43** (1986), 997-1002.

2. BASAR, T., On the uniqueness of the Nash solution in linear quadratic differential games, Internat. J. of Game Theory, vol. 5 (1973), 65-90.
3. ANDERSON, B.D.O., The testing of optimality linear systems, Internat. J. Control 4 No. 1 (1966), 29-40.
4. CRUZ, J.B., JR. and CHEN, C.I., Series Nash solution of two-person nonzero sum linear quadratic games, J. Opt. Theory and Appl. 7(1971), 240-257.
5. JÓDAR, L. and ABOU-KANDIL, H., A resolution method for Riccati differential systems coupled in their quadratic terms, SIAM J. Math. Anal. 19 (1988), 1225-1230.
6. KOKOTOVIC, P.V., A Riccati equation for block-diagonalization of ill-conditioned systems, IEEE Trans. Autom. Control AC-21(1975), 812-814.
7. LANCASTER, P. and TISMENETSKY, M., The theory of matrices, Second ed., Academic Press, New York, 1985.
8. MOLER, C.B., MATLAB User's guide, Technical Report CS81-1(1980) Department of Computer Science, Univ. of New Mexico, Alburquerque, New Mexico, 87131.
9. NARASIMHAMURTHI, N. and WU, F.F., On the Riccati equation arising from the study of singularly perturbed systems, Proc. of the 1977 Joint Autom. Control Conf. 22-24 June, San Francisco, Vol. V, 1244-1247.
10. OZGUNER, U. and PERKINS, W.R., A series solution to the Nash strategy for large scale interconnected systems, Automatica, Vol. 13(1977), 313-315.
11. RAO, C.R. and MITRA, S.K., Generalized Inverse of Matrices and its Applications, John-Wiley, New York, 1971.
12. STARR, A.W. and HO, Y.C., Non-zero sum differential games, J. Optimization Theory and Appl., 3(1969), 179-197.
13. VAN LOAN, C.F., Computing integrals involving the matrix exponential, IEEE Trans. Aut. Control, AC-23 (1978), 395-404.
14. SIMAAN, M. and CRUZ, J.B., JR., On the solution of open-loop Nash Riccati equations in linear quadratic differential games, Internat. J. Control, Vol. 18 (1973), 57-63.
15. MEYER, H.B., The matrix equation  $AZ + B - ZCZ - ZD = 0$ , SIAM J. Appl. Math. , Vol. 30, No. 1 (1976), 136-142.

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)