

## SUBCLASSES OF UNIFORMLY STARLIKE FUNCTIONS

ED MERKES

Department of Mathematical Sciences  
University of Cincinnati  
Cincinnati, Ohio 45221 U.S.A.

and

MOHAMMAD SALMASSI

Department of Mathematics  
College of the Holy Cross  
Worcester, MA 01610 U.S.A.

(Received October 23, 1990 and in revised form June 18, 1991)

**ABSTRACT.** We study subclasses of the class of uniformly starlike functions which were recently introduced by A.W. Goodman. One new subclass is defined and it is shown that it shares many properties of the parent class.

**KEY WORDS AND PHRASES.** Uniformly starlike functions, univalent functions, prestarlike functions, convex functions.

**1991 AMS SUBJECT CLASSIFICATION CODE.** 30C45.

### 1. INTRODUCTION.

Let  $ST$  be the class of analytic univalent functions  $f$  in the unit disk  $D = \{ |z| < 1 \}$  that are normalized by  $f(0) = 0, f'(0) = 1$ , and that are starlike with respect to the origin. The subclass of uniformly starlike functions (UST) consists of normalized analytic function  $f$  in  $D$  such that for each  $\zeta$  in  $D$  and any arc  $\gamma$  in  $D$  of a circle with center at  $\zeta$  is mapped by  $f$  onto an arc  $f(\gamma)$  that is starlike with respect to  $f(\zeta)$ . This class was recently introduced and studied by A.W. Goodman [1] who proved in particular the following analytic characterization.

**THEOREM A.** Let  $f$  be analytic in  $D$ ,  $f(0) = f'(0) - 1 = 0$ . Then  $f$  is in UST if and only if

$$Q(z, \zeta) = \frac{f(z) - f'(\zeta)}{(z - \zeta)f'(z)} \quad (z \neq \zeta) \text{ and } Q(z, z) = 1 \quad (1.1)$$

has positive real part for all  $z$  and  $\zeta$  in  $D$ .

Properties of the class UST are difficult to establish. One reason is that the usual transformations of univalent function theory generally do not preserve the UST class. The only known exceptions are rotations,  $e^{-i\alpha}f(e^{i\alpha}z)$  for some real  $\alpha$ , and the transformation  $t^{-1}f(tz)$ ,  $0 < t \leq 1$ .

In order to obtain a coefficient bound  $|a_n| \leq 2/n$  ( $n = 2, 3, \dots$ ) for  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$  in UST, Goodman proves that class UST is embedded in a larger subclass of starlike functions UST\*. A

function  $f \in ST$  is in  $UST^*$  if there is a real  $\alpha$  such that  $\operatorname{Re}\{e^{i\alpha}f'(z)\} \geq 0$  for  $z \in D$ . (Goodman credits the result to Charles Horowitz.) This suggests that information about the  $UST$  class might be generated through the study of subclasses of  $UST$  as well. In this paper we study one such class that shares and extends some known properties of  $UST$ .

## 2. SOME SUBCLASSES OF $UST$ .

If  $f(z) = \sum_{j=0}^{\infty} a_j z^j$  and  $g(z) = \sum_{j=0}^{\infty} b_j z^j$  are analytic in  $D$ , the Hadamard product  $(f*g)(z) = \sum_{j=0}^{\infty} a_j b_j z^j$  is also analytic in  $D$ . In particular, if  $f$  is a normalized analytic function in  $D$ , then for all complex numbers  $\alpha, \beta, 0 < |\alpha| \leq 1, \alpha \neq \beta$  we have

$$\frac{1}{\alpha} f(\alpha z) = f * \frac{z}{1 - \alpha z}, \frac{f(\alpha z) - f(\beta z)}{\alpha - \beta} = f * \frac{z}{(1 - \alpha z)(a - \beta z)}, z f'(\alpha z) = f * \frac{z}{(1 - \alpha z)^2}. \quad (2.1)$$

These identities lead to an equivalent form for Theorem A.

**THEOREM 1.** Let  $f$  be a normalized analytic function in  $D$ . Then  $f \in UST$  if and only if for all complex numbers  $\alpha, \beta, |\alpha| \leq 1, |\beta| \leq 1$ , and for all  $z \in D$  we have

$$\operatorname{Re} \left\{ \frac{f * \frac{z}{(1 - \alpha z)(1 - \beta z)}}{f * \frac{z}{(1 - \alpha z)^2}} \right\} \geq 0. \quad (2.2)$$

In this form we can appeal to the extensive work on Hadamard products initiated by the proof of the Polya-Schoenberg conjecture by Ruscheweyh and Sheil-Small [4]. The fundamental result in this proof was the following theorem [4].

**THEOREM B.** If  $\varphi$  is a normalized convex univalent function in  $D$  and  $g \in ST$ , then for all  $z \in D$

$$\operatorname{Re} \left\{ \frac{\varphi * (Fg)}{\varphi * g} \right\} \geq 0. \quad (2.3)$$

whenever  $F$  is an analytic function with positive real part in  $D$ .

These results, along with the following elementary observation about certain linear fractional transformations enable us to generate functions in  $UST$  from functions in the convex subclass  $K$  of  $ST$ .

**LEMMA.** Let  $\rho > 0$  and let  $\bar{D} = \{ |z| \leq 1 \}$ . Then  $\operatorname{Re}\{(1 - \alpha\rho z)/(1 - \beta\rho z)\} > 0$  for  $z \in D$  and for all  $\alpha, \beta$  in  $\bar{D}$  if and only if  $\rho \leq 1/\sqrt{2}$ .

The proof follows by considering the image of  $|z| = 1$  under the transformation

$$T = \frac{(1 - \alpha\rho z)}{(1 - \beta\rho z)}.$$

**THEOREM 2.** If  $f \in K$ , then  $\sqrt{2}f(z/\sqrt{2}) \in UST$ . The radius  $\rho = 1/\sqrt{2}$  is best possible.

**PROOF.** Let  $f \in K$ . Since  $z/(1 - \alpha\rho z)^2$  is in  $ST$  when  $|\alpha| \leq 1, \rho \in (0, 1)$ , we conclude from Theorem B and the Lemma that for  $z \in D$

$$\operatorname{Re} \left\{ \frac{f * \frac{(1 - \alpha\rho z)}{(1 - \beta\rho z)} \frac{z}{(1 - \alpha\rho z)^2}}{f * \frac{z}{(1 - \alpha\rho z)^2}} \right\} \geq 0$$

for all  $\alpha, \beta \in \bar{D}, 0 < \rho \leq 1/\sqrt{2}$ . But  $f*g(\rho z)/\rho = g*f(\rho z)/\rho$ . Hence, the expression above can be

rewritten as

$$\operatorname{Re} \left\{ \frac{\frac{1}{\rho} f(\rho z)^* \frac{z}{(1-\beta z)(1-\alpha z)}}{\frac{1}{\rho} f(\rho z)^* \frac{z}{(1-\alpha z)^2}} \right\} \geq 0 \quad z \in D, |\alpha| \leq 1, |\beta| \leq 1,$$

where  $0 < \rho \leq 1/\sqrt{2}$ . By Theorem 1 we have  $\frac{1}{\rho} f(\rho z) \in \text{UST}$  when  $0 < \rho \leq 1/\sqrt{2}$ . For the convex function  $f(z) = z/(1-z)$ , Theorem 1 states  $f(\rho z)/\rho = z/(1-\rho z) \in \text{UST}$  if and only if  $\operatorname{Re} \{(1-\alpha\rho z)/(1-\beta\rho z)\} > 0$  for  $z \in D$ ,  $|\alpha| \leq 1$ ,  $|\beta| \leq 1$ ,  $\rho > 0$ . This is the case by the Lemma only if  $\rho \leq 1/\sqrt{2}$ . The result is sharp. QED.

Goodman [1] proved  $z/(1-Az) \in \text{UST}$  whenever  $|A| \leq 1/\sqrt{2}$  which in particular establishes the sharpness of Theorem 2. This function, however, is also in  $\text{UST}^* \supset \text{UST}$ .

The function  $z/(1-\rho z)^2$  is starlike of order  $\alpha = (1-\rho)/(1+\rho)$ . Now a normalized analytic function  $f$  is said to be in the class  $R_\alpha$  of prestarlike functions if  $f^*z/(1-z)^{2-2\alpha}$  is in the class  $ST_\alpha$  of starlike functions of order  $\alpha$  when  $\alpha < 1$  or  $\operatorname{Re}\{f(z)/z\} > 1/2$  when  $\alpha = 1$ . Ruscheweyh [5, p. 54] proves a generalization of Theorem B to the case where  $\varphi \in R_\alpha$  and  $g \in ST_\alpha$ . By an argument that is similar to the proof of Theorem 2, we obtain the following result.

**THEOREM 3.** If  $f$  is in the class  $R_\alpha$  of prestarlike functions of order  $\alpha$ , then  $F(z) = f(\rho z)/\rho$  is in  $\text{UST}$  whenever  $\rho = (1-\alpha)/(1+\alpha)$  and  $(\sqrt{2}-1)/(\sqrt{2}+1) \leq \alpha < 1$ .

Except for the sharpness, Theorem 2 is a special case of Theorem 3 since  $R_0 = K \subset R_\alpha$  for  $0 < \alpha \leq 1$ . The link between the convex case and the fundamental Theorem B is our justification of first proving the less general result.

It is interesting to note that for  $\alpha > 1/2$ , the class  $R_\alpha$  contains functions that are not univalent in  $D$  [6]. The function  $F$  of Theorem 3 is, of course, univalent and starlike in  $D$ .

To obtain another subset of  $\text{UST}$ , notice that

$$\frac{f(z) - f(\zeta)}{z - \zeta} \frac{1}{f'(z)} = \frac{1}{f'(z)} \int_0^1 f'(tz + (1-t)\zeta) dt \quad (2.4)$$

**THEOREM 4.** Let  $f$  be a normalized analytic function in  $D$ . Then  $f \in \text{UST}$  if for all  $w, z$  in  $D$

$$\operatorname{Re} \left\{ \frac{f'(w)}{f'(z)} \right\} \geq 0.$$

If  $f \in \text{UST}$ , then for all  $w, z$  in  $D$

$$\operatorname{Re} \left\{ \frac{f'(w)}{f'(z)} \right\}^{1/2} \geq 0$$

and the  $1/2$  is best possible.

**PROOF.** The first part of the theorem follows from Theorem A and the real part of (2.4). For the second statement, we note that for  $f \in \text{UST}$

$$| \operatorname{arg} \left\{ \frac{f(w) - f(z)}{w - z} \frac{1}{f'(z)} \right\} | \leq \pi/2.$$

Hence,

$$| \operatorname{arg} \left\{ \frac{f'(w)}{f'(z)} \right\}^{1/2} | \leq \frac{1}{2} | \operatorname{arg} \frac{f(w) - f(z)}{w - z} \frac{1}{f'(z)} | + \frac{1}{2} | \operatorname{arg} \frac{w - z}{f(w) - f(z)} f'(w) | \leq \pi/2.$$

The function  $f(z) = z/(1-\rho z)$  with  $\rho = 1/\sqrt{2}$  proves the exponent  $1/2$  is best possible.

The condition  $\operatorname{Re} \{f'(w)/f'(z)\} \geq 0$  for  $z, w \in D$  implies  $f'(D)$  lies in a quarter plane that contains 1. Indeed,  $f'(0) = 1$  and  $f'(w) \neq itf'(z)$  for any real  $t$ . The condition  $\operatorname{Re} \{f'(w)/f'(z)\}^{1/2} \geq 0$  implies  $f'(D)$  lies in a half plane containing 1. This is the result of Goodman and Horowitz that proves  $\text{UST} \subset \text{UST}^*$ .

**COROLLARY 1.** [1] If  $f(z) = z + \sum_{j=2}^{\infty} a_j z^j$  as in UST, then there is an  $\alpha$ ,  $-\pi/2 < \alpha < \pi/2$  such that  $e^{i\alpha} f'(z)$  has positive real part in  $D$ . Furthermore,  $|a_n| \leq 2/n$  ( $n = 2, 3, \dots$ ).

The sharp bounds for the coefficients of  $f \in \text{UST}$  is an elusive open problem. Some information for this problem is contained in the next theorem.

**THEOREM 5.** For any integer  $n \geq 2$ , the function  $f(z) = z + Az^n$  is in UST if  $|A| \leq \sqrt{\frac{n+1}{2n^3}}$ .

**PROOF.** Since  $e^{-i\alpha} f(e^{i\alpha} z)$  is in UST whenever  $f$  is, we assume  $A > 0$ . For  $w \neq \xi$  nonzero in  $D$ , we have

$$\frac{f(w) - f(\xi)}{w - \xi} \frac{1}{f'(w)} = \frac{1 + A(w^{n-1} + \xi w^{n-2} + \dots + \xi^{n-1})}{1 + nA w^{n-1}}.$$

Replace  $w$ , and  $\xi$  respectively by  $z^{1/(n-1)}$ ,  $\alpha z^{1/(n-1)}$  where  $|\alpha| \leq 1$ ,  $\alpha \neq 1$ . The above expression becomes

$$\frac{1 + A(1 + \alpha + \alpha^2 + \dots + \alpha^{n-1})z}{1 + nAz} = \frac{1 + A\zeta z + Az}{1 + nAz},$$

where  $\zeta = \alpha + \alpha^2 + \dots + \alpha^{n-1}$ . The image of  $|z| = 1$  by this linear fractional transformation is a circle with center  $c$  and radius  $R$  given by

$$c = \frac{1 - nA^2(\zeta + 1)}{1 - n^2A^2}, R = \left| \frac{1 + A(\zeta + 1)}{1 + nA} - \frac{1 - nA^2(\zeta + 1)}{1 - n^2A^2} \right| = \frac{A}{|1 - nA^2|} |(n-1) - \zeta|.$$

The image is in the right half plane if

$$\frac{|(n-1) - \zeta|}{|1 - nA^2|} A \leq \left| \frac{1 - nA^2(\operatorname{Re} \zeta + 1)}{1 - n^2A^2} \right|.$$

Since  $|\zeta| = |\alpha + \alpha^2 + \dots + \alpha^{n-1}| \leq n-1$ , the inequality holds when

$$A^2[2(n-1)^2 - 2(n-1)x] \leq (1 - nA^2)^2 - 2nA^2(1 - nA^2)x + n^2A^2x^2$$

where  $x = \operatorname{Re} \zeta$ , that is,

$$n^2A^4x^2 + 2(n^2A^2 - 1)A^2x + 1 - 2(n^2 - n + 1)A^2 + n^2A^4 \geq 0.$$

The minimum of the function of  $x$  on the left of this inequality occurs when

$$x = -(n^2A^2 - 1)/n^2A^2.$$

If we substitute this for  $x$  and simplify we obtain

$$(n-1)(n+1 - 2n^3A^2) \geq 0 \text{ or } \frac{n+1}{2n^3} \geq A^2. \quad \text{QED.}$$

This improves the bound  $|A| \leq 1/(\sqrt{2}n)$  of Goodman [1]. It does not appear to be the best possible except when  $n = 2$ . Goodman [1] states that  $z + Az^2 \in \text{UST}$  if and only if  $|A| \leq \sqrt{3}/4$ . We prove this result for a subclass of UST in the next section.

3. THE SUBCLASS  $UST_*$ .

A rather natural way to construct a subclass of  $UST$  is to replace the derivative in (1.1) by a difference quotient. This generates a family of functions that shares most of the known sharp properties of the class  $UST$ . To be precise, we define  $UST_*$  as the class of normalized analytic functions  $f$  in  $D$  such that

$$\operatorname{Re} \left\{ \frac{f^*z/(1-\alpha z)(1-\beta z)}{f^*z/(1-\alpha z)(1-\gamma z)} \right\} \geq 0, \quad (3.1)$$

for all  $z \in D$ , and  $\alpha, \beta, \gamma$  in  $\bar{D}$ . Since  $\gamma = \alpha$  reduces (3.1) to (2.2), we have  $UST_* \subset UST$ .

The function  $z/(1-\alpha\rho z)(1-\gamma\rho z)$  for  $z \in D$ ,  $\rho > 0$  and  $\alpha, \gamma \in \bar{D}$  is starlike of order  $\alpha = (1-\rho)/(1+\rho)$ . We conclude from the generalized version of Theorem B that the functions  $F$  of Theorem 3 are in fact in the class  $UST_*$ .

**THEOREM 6.** If  $f$  is in the class  $R_\alpha$ , then  $F(z) = f(\rho z)/\rho$  is in  $UST_*$  whenever

$$\rho = (1-\alpha)/(1+\alpha) \text{ and } (\sqrt{2}-1)/(\sqrt{2}+1) \leq \alpha < 1.$$

**COROLLARY 1.** If  $f \in K$ , then  $\sqrt{2}f(z/\sqrt{2}) \in UST_*$ . This result is sharp.

To prove the sharpness in the Corollary, we observe from (3.1) that  $z/(1-\rho z) \in UST_*$  if and only if  $(1-\gamma\rho z)/(1-\beta\rho z)$  has positive real part for  $z \in D$  and some  $\rho > 0$  whenever  $\gamma, \beta \in \bar{D}$ . By the Lemma this requires  $0 < \rho \leq 1/\sqrt{2}$ .

There is another way to characterize  $UST_*$  (and  $UST$ ) that can be useful.

**THEOREM 7.** Let  $f$  be a normalized analytic function in  $D$ . Then  $f \in UST_*$  if and only if for  $z \in D$ ,  $z \neq 0$ ,

$$f^* \frac{z}{(1-\alpha z)} \frac{1 - \frac{1}{2}[(1+x)\gamma + (1-x)\beta]z}{(1-\beta z)(1-\gamma z)} \neq 0 \quad (3.2)$$

for all  $|x| = 1$  and  $\gamma, \beta, \alpha$  in  $\bar{D}$ .

This result follows directly from (3.1). The expression in braces in (3.1) cannot equal  $(x+1)/(x-1)$ ,  $|x| = 1$ , when its real part is positive. This yields (3.2) upon algebraic simplification.

**COROLLARY 2.**  $z + Az^2 \in UST_*$  if and only if  $|A| \leq \sqrt{3}/4$ .

**PROOF.** If  $a_2$  is the coefficient of  $z^2$  in the power series expansion of the second function in the Hadamard product (3.2), then with  $f(z) = z + Az^2$  this product is not zero in  $D$  if  $|Aa_2| \leq 1$ . Now, with  $\alpha = 1$ , the second coefficient is

$$a_2 = 1 + \frac{1}{2}[(1-x)\gamma + (1+x)\beta].$$

Now

$$|a_2| \leq |1 + \frac{1}{2}(\gamma + \beta)| + \frac{1}{2}|\gamma - \beta|$$

and we seek a maximum of the right hand side when  $|\gamma| \leq 1$ ,  $|\beta| \leq 1$ . A computation shows a maximum occurs when  $\gamma = 1$ ,  $\beta = e^{it}$ ,  $t = 2 \arcsin 1/\sqrt{3}$ . This gives us  $|a_2| \leq 4/\sqrt{3}$  and proves that  $|A| \leq \sqrt{3}/4$ . Since there is a choice of  $x$ ,  $|x| = 1$ , such that  $|a_2| = |1 + \frac{1}{2}(\gamma + \beta)| + \frac{1}{2}|\gamma - \beta|$ , we conclude that the result is sharp.

4. ARC LENGTH FOR THE CLASS  $UST$ .

In the final section we state a result on the length of images of circles under  $UST$  mappings which extends a well-known result of Keogh for bounded starlike functions [3]. Let  $C(\zeta, r)$  be a

circle centered at  $\zeta$  and radius  $r$  which is strictly inside the unit disk and let  $\Gamma$  be the image of this circle under the function  $f$  which is in the class UST. Furthermore, let

$$M_r = \max_{|z-\zeta|=r} |f(z) - f(\zeta)|,$$

and let  $L_r$  be the arc length of  $\Gamma$ .

**THEOREM 8.** Given the foregoing definitions we have

$$L_r \leq \frac{M_r}{r} \frac{|\zeta|}{|f(\zeta)|} (2|\zeta| + r) \left\{ 2\pi + 4\log \frac{1 + |\zeta| + r}{1 - |\zeta| - r} \right\} \quad (3.3)$$

where  $\zeta/f(\zeta) = 1$  if  $\zeta = 0$ . In particular,  $L_r = O\left(\log \frac{1}{1 - |\zeta| - r}\right)$ . This result reduces to Keogh's when  $\zeta = 0$ .

The technique of the proof of (3.3) is similar to Keogh's except that we use the following result of R.M. Gabriel [2].

If  $u(z)$  is subharmonic, positive, and continuous inside and on a circle  $\Gamma$ , and  $C$  is a circle inside  $\Gamma$ , then

$$\int_C u(z) |dz| < (1 + \rho/R) \int_{\Gamma} u(z) |dz| ,$$

where  $\rho$  is the distance between centers of  $C$  and  $\Gamma$  and  $R$  is the radius of  $\Gamma$ .

**ACKNOWLEDGEMENT.** The second author thanks Ted Suffridge for his insightful suggestions and John Anderson for his remarks on Theorem 8.

#### REFERENCES

- GOODMAN, A.W., On uniformly starlike functions, to appear in the *Journal of Math Analysis*.
- GABRIEL, R.M., An inequality concerning the integrals of positive subharmonic functions along certain circles, *J. London Math. Soc.* **V**, 1930, 129-131.
- KEOGH, F.R., Some theorems on conformal mapping of bounded star-shaped domains, *Proc. London Math. Soc. (3)* **9** (1959), 481-491.
- RUSCHEWEYH, ST. and SHEIL-SMALL, T., Hadamard products of schlicht functions and the Polya-Schoenberg conjecture, *Comment. Math. Helv.* **48** (1973), 119-135.
- RUSCHEWEYH, ST., "Convolutions in Geometric Function Theory", Les Presses de l'Universite de Montreal, 1982.
- SILVERMAN, H. and SILVIA, E.M., Prestarlike functions with negative coefficients, *Internat. J. Math. & Math. Sci.* **3** (1979), 427-439.

## Special Issue on Intelligent Computational Methods for Financial Engineering

### Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Lean Yu**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [yulean@amss.ac.cn](mailto:yulean@amss.ac.cn)

**Shouyang Wang**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; [sywang@amss.ac.cn](mailto:sywang@amss.ac.cn)

**K. K. Lai**, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [mskklai@cityu.edu.hk](mailto:mskklai@cityu.edu.hk)