

**A NOTE ON THE FREE CONVECTION BOUNDARY LAYER
ON A VERTICAL SURFACE WITH PRESCRIBED HEAT FLUX
AT SMALL PRANDTL NUMBER**

J. H. MERKIN

Department of Applied Mathematics
University of Leeds
Leeds LS2 9JT, U.K.

and

V. KUMARAN

Department of Mathematics
Anna University
Madras - 600025
India

(Received May 7, 1991)

ABSTRACT. It is shown that for a particular case of the surface heat flux the equations for small Prandtl number have simple analytical solutions. These are presented and compared with numerical solutions of the general equations.

KEY WORDS AND PHRASES. Free convection, boundary layers and heat flux.

1991 AMS SUBJECT CLASSIFICATION CODE. 76R10, 76N20.

1. INTRODUCTION.

In a recent paper [1], the solution for the free convection boundary-layer flow on a vertical plate with a prescribed surface heat flux valid for small Prandtl numbers was derived. The surface heat flux was taken to be proportional to x^λ , where x is the distance from the leading edge and λ is a constant, with the governing equations then being reducible to similarity form. Results in [1] were given for the case of uniform wall heat flux, i.e., $\lambda = 0$. A further consideration of this problem reveals that, for the case when $\lambda = 1$, simple analytical solutions are possible. It is the purpose of this note to present these solutions, and, as analytical solutions in free convection boundary-layer theory are somewhat of a rarity, this analysis is worth describing.

2. ANALYSIS.

Following [1], the governing similarity equations are, for $\lambda = 1$,

$$f''' + \theta + ff'' - f'^2 = 0 \quad (2.1a)$$

$$\theta'' + \sigma(f\theta' - f'\theta) = 0 \quad (2.1b)$$

with,

$$f(0) = 0, \quad f'(0) = 0, \quad \theta'(0) = -1, \quad f' \rightarrow 0, \quad \theta \rightarrow 0 \text{ as } \eta \rightarrow \infty, \quad (2.1c)$$

where primes denote differentiation with respect to the independent variable η and σ is the Prandtl number. There is an inner region, in which

$$f = \sigma^{-1/10} F(\zeta), \quad \theta = \sigma^{-2/5} H(\zeta), \quad \zeta = \sigma^{-1/10} \eta. \quad (2.2)$$

A consideration of the equations in this region leads to,

$$H = a_0 + \sigma^{1/2}(a_1 - \zeta) + \dots \quad (2.3a)$$

The equation for F is, at leading order, given by a Falkner-Skan equation, and, as $\zeta \rightarrow \infty$,

$$F \sim a_0^{1/2}\zeta + b_0 + \sigma^{1/2} \left[-\frac{\zeta^2}{2a_0^{1/2}} + \frac{1}{2a_0} (a_0 a_1 - b_0) \zeta + b_1 \right] + \dots \quad (2.3b)$$

The constants a_0 and a_1 are determined from the matching with the outer region, and b_0 is determined from the solution of the equation for F in the inner region. In the outer region,

$$f = \sigma^{-3/5}\phi(Y), \quad \theta = \sigma^{-2/5}h(Y), \quad Y = \sigma^{2/5}\eta. \quad (2.4)$$

Using (2.4) in equations (2.1abc) gives the equations for the outer region as

$$h + \phi\phi'' - \phi'^2 + \sigma\phi''' = 0 \quad (2.5a)$$

$$h'' + \phi h' - \phi' h = 0 \quad (2.5b)$$

(where primes now denote differentiation with respect to Y). The boundary conditions to be satisfied by equations (2.5ab) are that,

$$\phi' \rightarrow 0, \quad h \rightarrow 0 \quad \text{as } Y \rightarrow \infty \quad (2.6a)$$

and, from matching within the inner region, that

$$h \sim a_0 - Y + \dots + \sigma^{1/2}(a_1 + \dots) + \dots, \quad (2.6b)$$

$$\begin{aligned} \phi \sim a_0^{1/2} Y - \frac{1}{2} a_0^{-1/2} Y^2 + \dots \\ + \sigma^{1/2}(b_0 + \frac{1}{2a_0} (a_0 a_1 - b_0) Y + \dots) + \dots \end{aligned} \quad (2.6c)$$

for Y small.

(2.6bc) suggests looking for a solution of equations (2.5ab) by expanding

$$\phi = \phi_0 + \sigma^{1/2}\phi_1 + \dots, \quad h = h_0 + \sigma^{1/2}h_1 + \dots \quad (2.7)$$

At leading order we obtain the equations

$$h_0 + \phi_0\phi_0'' - \phi_0'^2 = 0 \quad (2.8a)$$

$$h_0'' + \phi_0 h_0' - \phi_0' h_0 = 0 \quad (2.8b)$$

It is straightforward to show that the solution of equations (2.8ab), which satisfies boundary conditions (2.6abc) is

$$a_0 = 1, \quad \phi_0 = 1 - e^{-Y}, \quad h_0 = e^{-Y}. \quad (2.9)$$

The solution can be continued to higher order terms. We find that, at $0(\sigma^{1/2})$,

$$a_1 = -b_0, \quad \phi_1 = b_0 e^{-Y}, \quad h_1 = -b_0 e^{-Y} \quad (2.10)$$

Using the value for a_0 given by (2.9), the appropriate Falkner-Skan equation for the leading order term F_0 in the inner layer can be solved. This gives, [2],

$$F_0''(0) = 1.23259, \quad b_0 = -0.64790.$$

Then using the value for b_0 , the (linear) equation for F_1 , the term of $0(\sigma^{1/2})$ in the inner region,

can be solved, giving

$$F_1''(0) = -0.41392, \quad b_1 = -0.62264.$$

3. RESULTS.

The analysis presented above gives, from (2.2), (2.3ab), (2.4), (2.9) and (2.10)

$$\left[\frac{d^2 f}{d\eta^2} \right]_{\eta=0} = \sigma^{-1/10} (1.23259 - 0.41392 \sigma^{1/2} + 0(\sigma)) \quad (3.1a)$$

$$\theta(0) = \sigma^{-25} (1 + 0.64790 \sigma^{1/2} + 0(\sigma)) \quad (3.1b)$$


$$f(\infty) = \sigma^{-3/5} (1 + 0(\sigma)) \quad (3.1c)$$



for σ small.

To check on the validity of the series approximations (3.1abc), we compared these with values obtained from a numerical solution of equations (2.1abc). The results are shown in figures 1, where we give the numerically determined values of $\left[\frac{d^2 f}{d\eta^2} \right]_0 \sigma^{1/10}$, $\theta(0)\sigma^{2/5}$ and $f(\infty)\sigma^{3/5}$ (shown by the broken line) and these quantities as calculated from (3.1abc) (shown by the full line). In all three cases we can see that the numerically determined values and (3.1abc) are in good agreement, even at the relatively large value of $\sigma = 0.2$, and that the agreement between the two sets of results improves as σ is decreased. It is worth noting that the linear slope of the numerical results in figure 1c appears to suggest that the correction to (3.1abc) is of $0(\sigma)$ and that no extra powers of σ are required (at least up to this order) in the expansions in the inner and outer regions (as was required in the general case given in [1]).

Figure 1 Graphs of (a) $\sigma^{1/10} \left[\frac{d^2 f}{d\eta^2} \right]_0$, (b) $\sigma^{2/5} \theta(0)$, and

(c) $\sigma^{3/5} f(\infty)$ obtained from a numerical solution of equations (1) (broken line) and from series expansions (11) (full line).





REFERENCES

1. MERKIN, J.H., Free convection on a heated vertical plate: the solution for small Prandtl number. *J. Engng. Math.* **23** (1989) 273-282.
2. ROSENHEAD, L. (editor), Laminar boundary layers. Clarendon Press, (Oxford), 1963.

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk