

SPACES OF COMPACT OPERATORS WHICH ARE M -IDEALS IN $L(X, Y)$

CHONG-MAN CHO

Hanyang University
Seoul 133-791, Korea

(Received December 10, 1991)

ABSTRACT. Suppose X and Y are reflexive Banach spaces. If $K(X, Y)$, the space of all compact linear operators from X to Y is an M -ideal in $L(X, Y)$, the space of all bounded linear operators from X to Y , then the second dual space $K(X, Y)^{**}$ of $K(X, Y)$ is isometrically isomorphic to $L(X, Y)$.

KEY WORDS AND PHRASES. Compact operators, M -ideal, dual space, projective tensor product.

1992 AMS SUBJECT CLASSIFICATION CODES. Primary 46B28, Secondary 46A32, 47D15.

1. INTRODUCTION

It is well known that if X and Y are reflexive Banach spaces one of which satisfies the approximation property then the second dual space $K(X, Y)^{**}$ of $K(X, Y)$, the space of compact linear operators from X to Y , is isometrically isomorphic to $L(X, Y)$, the space of bounded linear operators from X to Y (Diestel [1, p.17]). Harmand and Lima [2] proved that if X is a reflexive Banach space and $K(X)$ is an M -ideal in $L(X)$ then $K(X)^{**}$ is isometrically isomorphic to $L(X)$.

The purpose of this paper is to generalize the result of Harmand and Lima to the case of $K(X, Y)$ and $L(X, Y)$ by modifying their proof. In Theorem 3.3 we will prove that if X and Y are reflexive Banach spaces and $K(X, Y)$ is an M -ideal in $L(X, Y)$ then $K(X, Y)^{**}$ is isometrically isomorphic to $L(X, Y)$.

2. NOTATIONS AND PRELIMINARIES.

Let X and Y be Banach spaces. $X \simeq Y$ means that X and Y are isometrically isomorphic. $L(X, Y)$ (resp. $K(X, Y)$) will denote the space of all bounded linear operators (resp. compact linear operators) from X to Y . If $X = Y$, then we simply write $L(X)$ (resp. $K(X)$). X^* will denote the dual space of X and we will write $\langle x, x^* \rangle$ for the action of $x^* \in X^*$ on $x \in X$ instead of $x^*(x)$. B_X will denote the closed unit ball of X .

A closed subspace J of a Banach space X is called an L -summand if there exists a projection P on X such that $PX = J$ and $\|x\| = \|Px\| + \|x - Px\|$ for every x in X . In this case we write $X = J \oplus_1 J'$ where $J' = (I - P)X$. A closed subspace J of a Banach space X is called an M -ideal in X if J° , the annihilator of J in X^* , is an L -summand in X^* .

Let $X \hat{\otimes} Y$ be the projective tensor product of Banach spaces X and Y . If $u \in X \hat{\otimes} Y$, then there exist sequences (x_i) in X and (y_i) in Y such that $u = \sum_{i=1}^{\infty} x_i \otimes y_i$, with $\sum_{i=1}^{\infty} \|x_i\| \|y_i\| < \infty$. Moreover, we have $\|u\| = \inf \sum_{i=1}^{\infty} \|x_i\| \|y_i\| < \infty$, the infimum being taken over all representations $u = \sum_{i=1}^{\infty} x_i \otimes y_i$, $x_i \in X$, $y_i \in Y$ and $\sum_{i=1}^{\infty} \|x_i\| \|y_i\| < \infty$. (Diestel and Uhl [3, p. 227]).

Let Z be another Banach space and $T \in L(X, Z)$. We define $Tu = \sum_{i=1}^{\infty} x_i \otimes Ty_i$ for $u = \sum_{i=1}^{\infty} x_i \otimes y_i \in X \hat{\otimes} Y$. Then $Tu \in X \hat{\otimes} Z$ and $\|Tu\| \leq \|T\| \|u\|$. If $u = \sum_{i=1}^{\infty} x_i^* \otimes x_i \in X^* \hat{\otimes} X$ with $\sum_{i=1}^{\infty} \|x_i^*\| \|x_i\| < \infty$, the map $u \mapsto \langle x_i, x_i^* \rangle$ defines a bounded linear functional on $X^* \hat{\otimes} X$ with norm no larger than 1.

THEOREM 2.1 (Diestel and Uhl [3], Shatten [4]). Let X and Y be Banach spaces. The Banach space $L(X, Y^*)$ is isometrically isomorphic to $(Y \hat{\otimes} X)^*$ and under this identification $T \in L(X, Y^*)$ act on $u \in Y \hat{\otimes} X$ by $\langle u, T \rangle = \text{tr}(Tu)$.

THEOREM 2.2 (Feder and Sapher [5]). Let X and Y be Banach spaces. If either X^{**} or Y^* has the Radon-Nikodym Property, then map $V: Y^* \hat{\otimes} X^{**} \rightarrow K(X, Y)^*$ defined by $\langle T, V(u) \rangle = \text{tr}(T^{**}u)$ for $T \in K(X, Y)$ and $u \in Y^* \hat{\otimes} X^{**}$ is a quotient map.

3. SPACES OF COMPACT OPERATORS

Harmand and Lima [2] proved that if $K(X)$ is an M -ideal in $L(X)$ then there exists a net (T_α) in $B_{K(X)}$ such that

- (i) $T_\alpha z \rightarrow z$ for all $z \in X$
- (ii) $T_\alpha^* z^* \rightarrow z^*$ for all $z^* \in X^*$
- (iii) $\|T_\alpha - I\| \rightarrow 1$.

In the case of $K(X, Y)$ and $L(X, Y)$, we have the following analogue which also plays a key role in the proof of our main result (Theorem 3.3).

THEOREM 3.1. If X and Y are Banach spaces and $K(X, Y)$ is an M -ideal in $L(X, Y)$, then for each T in $B_{L(X, Y)}$ there is a net (T_α) in $B_{K(X, Y)}$ such that

- (i) $T_\alpha z \rightarrow Tz$ for all $z \in X$
- (ii) $T_\alpha^* y^* \rightarrow T^* y^*$ for all $y^* \in Y^*$.

PROOF. Suppose $K(X, Y)$ is an M -ideal in $L(X, Y)$. Then we can write $L(X, Y)^* = K(X, Y)^\circ \oplus J$ for some subspace J of $L(X, Y)^*$.

The map $\phi \rightarrow \phi + K(X, Y)^\circ$ defines an isometry from J onto $L(X, Y)^*/K(X, Y)^\circ$ and the map $\phi + K(X, Y)^\circ \rightarrow \phi|_{K(X, Y)}$ defines an isometry from $L(X, Y)^*/K(X, Y)^\circ$ onto $K(X, Y)^*$ (Rudin [6, p.91]). Hence the map $\phi \rightarrow \phi|_{K(X, Y)}$ gives an isometry from J onto $K(X, Y)^*$.

Let Q be the projection on $L(X, Y)^*$ with the range J . Then $\phi \in L(X, Y)^*$ is in the range of Q if and only if the restriction of ϕ to $K(X, Y)$ has the same norm as ϕ . If $T \in L(X, Y) \subseteq L(X, Y)^{**}$ with $\|T\| \leq 1$, then for $\phi \in K(X, Y)^\circ$ we have $(Q^*T)\phi = TQ(\phi) = 0$ thus $Q^*T \in K(X, Y)^\circ = J^* = K(X, Y)^{**}$. Since $Q^*T \in K(X, Y)^{**}$ and $\|Q^*T\| \leq 1$, by the Goldstein's theorem there is a net (T_α) in $B_{K(X, Y)}$ such that

$T_\alpha \rightarrow Q^*T$ in the weak*-topology on $J^* = K(X, Y)^{**}$.

We claim that $T_\alpha z \rightarrow Tz$ for all $z \in X$ and $T_\alpha^* y^* \rightarrow T^* y^*$ for all $y^* \in Y^*$. For $z^{**} \in X^{**}$ and $y^* \in Y^*$, define $\phi_{z^{**}} \otimes y^* \in L(X, Y)^*$ by

$$\langle A, \phi_{z^{**}} \otimes y^* \rangle = \langle A^* y^*, z^{**} \rangle.$$

Then we can easily see that $\phi_{z^{**}} \otimes y^* \in J = K(X, Y)^*$ and hence

$$\langle T_\alpha^* y^*, z^{**} \rangle \rightarrow \langle T^* y^*, z^{**} \rangle.$$

By the weak*-compactness of $B_{X^{**}}$ we get that

$$T_\alpha^* \rightarrow T^* y^* \text{ for all } y^* \in Y^*.$$

Similarly, for $y^* \in Y^*$ and $z \in X$ the functional $\psi_{y^*} \otimes z$ on $L(X, Y)$ defined by $\langle A, \psi_{y^*} \otimes z \rangle = \langle Az, y^* \rangle$ for $A \in L(X, Y)$ is in the range of Q and hence $T_\alpha z \rightarrow Tz$ for all $z \in X$.

The following proposition is essentially due to Harmand and Lima [2] who treated a special case $X = Y$.

PROPOSITION 3.2. Let X and Y be Banach spaces and V the map defined in Theorem 2.2. If $K(X, Y)$ is an M -ideal in $L(X, Y)$, then $T^{**} \in (\ker V)^\circ$ for every $T \in L(X, Y)$.

PROOF. Recall that by Theorem 2.1 we have $(Y^* \hat{\otimes} X^{**})^* \simeq L(X^{**}, Y^{**})$ and under this identification $S \in L(X^{**}, Y^{**})$ acts on $u \in Y^* \hat{\otimes} X^{**}$ by $\langle u, S \rangle = \text{tr}(Su)$.

Let $T \in L(X, Y)$, $\|T\| \leq 1$. By Theorem 3.1 there is a net (T_α) in $B_{K(X, Y)}$ such that $T_\alpha^* y^* \rightarrow T^* y^*$ for all $y^* \in Y^*$. Let $u = \sum_{i=1}^{\infty} y_i^* \otimes z_i^{**} \in \ker V$ with $\sum_{i=1}^{\infty} \|y_i^*\| \|z_i^{**}\| < \infty$. We may assume that $\|z_i^{**}\| \leq 1$ for all i and

$\|y_i^*\| \rightarrow 0$. Then we get

$$\begin{aligned}
 0 &= \langle T_\alpha, V(u) \rangle \\
 &= \text{tr}(T_\alpha^{**} u) \\
 &= \sum_{i=1}^{\infty} \langle y_i^*, T_\alpha^{**} x_i^{**} \rangle \\
 &= \sum_{i=1}^{\infty} \langle T_\alpha^* y_i^*, x_i^{**} \rangle \\
 &\rightarrow \sum_{i=1}^{\infty} \langle T^* y_i^*, x_i^{**} \rangle \\
 &= \text{tr}(T^{**} u) \\
 &= \langle u, T^{**} \rangle.
 \end{aligned}$$

Thus $T^{**} \in (\ker V)^\circ$.

THEOREM 3.3. If X and Y are reflexive Banach spaces and $K(X, Y)$ is an M -ideal in $L(X, Y)$ then $K(X, Y)^{**}$ is isometrically isomorphic to $L(X, Y)$.

PROOF. Since X and Y are reflexive, X and Y^* have the Radon-Nikodym property and hence by Theorem 2.2 the map $V: Y^* \hat{\otimes} X^{**} \rightarrow K(X, Y)^*$ defined by

$$\langle T, V(U) \rangle = \text{tr}(T^{**} u) \text{ for } u \in Y^* \hat{\otimes} X^{**}, T \in K(X, Y)$$

is a quotient map. Thus $V^*: K(X, Y)^{**} \rightarrow (Y^* \hat{\otimes} X^{**})^*$ is an isometry with the range $(\ker V)^\circ$ and hence we have

$$\begin{aligned}
 K(X, Y)^{**} &\simeq (\ker V)^* \\
 &\subseteq (Y^* \hat{\otimes} X^{**})^* \\
 &\simeq L(X^{**}, Y^*) \\
 &= L(X, Y).
 \end{aligned}$$

Since X and Y are reflexive, $T = T^{**}$ for all $T \in L(X, Y)$ and by Proposition 3.2 $(Y^* \hat{\otimes} X^{**})^* \subseteq (\ker V)^\circ$. Thus $K(X, Y)^{**} \simeq L(X, Y)$.

Recall that for $1 \leq p \leq \infty$ the l_p -sum $(\sum X_n)_p$ of a sequence of (X_n) of Banach spaces is the Banach space of all sequences (x_n) with $x_n \in X_n$ and with the norm $\| (x_n) \| = (\sum \|x_n\|^p)^{1/p} < \infty$.

COROLLARY 3.4. Suppose X and Y are closed subspaces of $(\sum X_n)_p$ and $(\sum Y_n)_q$ ($1 < p \leq q \leq \infty$, $\dim X_n < \infty$, $\dim Y_n < \infty$), respectively. If $K(X, Y)$ is dense in $L(X, Y)$ in the strong operator topology, then $K(X, Y)^{**} \simeq L(X, Y)$.

PROOF. X and Y are reflexive and $K(X, Y)$ is an M -ideal in $L(X, Y)$ (Cho [7]).

REMARK. If X and Y are as in Corollary 3.4 and either X or Y satisfies the compact approximation property, then $K(X, Y)$ is dense in $L(X, Y)$ in the strong operator topology [7] and hence $K(X, Y)^{**} \simeq L(X, Y)$.

ACKNOWLEDGEMENT. This work was supported by KOSEF Grant No. 891-0102-016-2.

REFERENCES

1. DIESTEL, J. Geometry of Banach Spaces-Selected Topics, Lecture Note in Math., Vol. 485, Springer-Verlag, Berlin and New York, 1975.
2. HARMAND, P. and LIMA, A., Banach Spaces which are M -ideals in their Duals, Trans. Amer. Math. Soc. 238 (1983), 253-264.
3. DIESTEL, J. and UHL, J., Vector Measures, American Mathematical Survey No. 15(1977).
4. SHATTEN, R., A Theory of Cross-space, Ann. of Math. Studies 26, 1950.
5. FEDER, M. and SAPHAR, P., Spaces of Compact Operators and their Dual Spaces, Israel J. Math. 21 (1975), 38-49.
6. RUDE, W., Functional Analysis, McGraw-Hill Book Company, 1973.
7. CHO, C. M., A Note on M -ideals of Compact Operators, Canadian Math Bull. 32(1989), 434-440.

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk