

**ON POINT DISSIPATIVE N-DIMENSIONAL SYSTEMS OF DIFFERENTIAL  
EQUATIONS WITH QUADRATIC NONLINEARITY**

ANIL K. BOSE  
ALAN S. COVER  
JAMES A. RENEKE

Department of Mathematical Sciences  
Clemson University  
Clemson, SC 29634-1907

(Received October 17, 1991)

**ABSTRACT.** Known sufficient conditions for quadratic dynamical system  $\mathbf{x}' = \mathbf{A}\mathbf{x} + \mathbf{f}(\mathbf{x})$  to be point dissipative given in terms of  $\mathbf{A}$  and  $\mathbf{f}$  for dimensions 2 and 3 are extended to allow for more general forms for the nonlinear term  $\mathbf{f}(\mathbf{x})$ . Furthermore, the conditions extend to  $n$  dimensions when  $\mathbf{f}$  is quadratic with zero set an  $(n - 1)$ -dimensional hyperplane.

**0. INTRODUCTION**

We are concerned with a class of vector equations of the form  $\mathbf{x}' = \mathbf{A}\mathbf{x} + \mathbf{f}(\mathbf{x})$  where the nonlinear term  $\mathbf{f}(\mathbf{x})$  is quadratic of the form

$$\mathbf{f}(\mathbf{x}) = \begin{bmatrix} \mathbf{x}^T \mathbf{C}_1 \mathbf{x} \\ \vdots \\ \mathbf{x}^T \mathbf{C}_n \mathbf{x} \end{bmatrix}$$

The  $n \times n$  matrices  $\{\mathbf{C}_i\}$  are assumed symmetric with the orthogonality property  $\mathbf{x}^T \mathbf{f}(\mathbf{x}) = 0$  for all  $\mathbf{x}$ . If  $\mathbf{x}^T \mathbf{f}(\mathbf{x}) = 0$  for all  $\mathbf{x}$  we say that  $\mathbf{f}$  is a conservative function. Note that if  $\mathbf{x}' = \mathbf{f}(\mathbf{x})$  with  $\mathbf{f}$  conservative then  $\|\mathbf{x}\|^2$  is constant. The problem is to determine conditions on  $\mathbf{A}$  and  $\mathbf{f}$  sufficient to have the system point dissipative, i.e., which guarantee the existence of a bounded region  $R$  with the property that every trajectory of the system eventually enters and remains within  $R$  [2].

Consider the Lyapunov function  $V(\mathbf{x}) = \frac{1}{2}(\mathbf{x} - \alpha)^T(\mathbf{x} - \alpha)$  for the system. For large  $\|\mathbf{x}\|$  the quadratic terms  $\mathbf{x}^T \mathbf{A} \mathbf{x} - \alpha^T \mathbf{f}(\mathbf{x})$  dominate  $\frac{d(V(\mathbf{x}(t)))}{dt}$ . Therefore our quest is to find conditions on

$A$  and  $f$  for which there is an admissible  $\alpha$  which makes these terms a negative definite function. If there exists an admissible  $\alpha$  then a classic result [ 4 ] implies the system is point dissipative.

A necessary condition for the existence of an admissible  $\alpha$  is that  $x^T A x < 0$  for each nontrivial  $x$  in the zeros of  $f$ . We have shown for  $n = 2$  and  $3$  [ 1 ] that this condition is also sufficient. Settling the obvious conjecture for all  $n$  bogs down in a proliferation of cases. However, we have shown in this paper that the necessary condition is sufficient for the simplest  $n$ -dimensional case, namely, when the zeros of  $f$  form an  $(n - 1)$ -dimensional hyperplane.

Finally, we extend the notion of an admissible  $\alpha$  to provide sufficient conditions for systems  $x' = Ax + f(x) + g(x)$  to be point dissipative when  $f$  is quadratic but not conservative and  $g$  is not quadratic. If there is a positive definite matrix  $S$  such that  $Sf$  is conservative then a vector  $\alpha$  is admissible for the dynamical system  $x' = Ax + f(x)$  if  $x^T S A x - \alpha^T H f(x)$  is a negative definite function where  $H^T H = S$ . If  $\alpha$  is an admissible vector for the system  $x' = Ax + f(x)$  then the system  $x' = Ax + f(x) + g(x)$  is point dissipative if there is an ordered triple  $(\varepsilon, C, M)$  such that  $-\alpha^T H g(x) \leq C \|x\|^2 - \varepsilon$  for all  $x$  with  $\|x\| \geq M$ .

Our interest in systems of the form  $x' = Ax + f(x)$  was stimulated by all of the work in the literature based on systems of the same form originally studied by Lorenz [ 5 ]. We hope to understand the richness of the class of chaotic systems, especially of dimension  $n > 3$ , by classifying a sufficiently rich class of point dissipative systems in terms of their compact attractors. This paper represents a step forward in that program by enlarging the class of systems that can be first classified as point dissipative in terms of their coefficients.

## 1. PRELIMINARIES

The proof of the principal result Theorem 3 uses some properties of skew symmetric matrices. These are matrices such that  $A^T = -A$ . Here are properties that are needed.

1.1. We note that  $k^T$  is in the kernel of a skew symmetric  $B$  relative to left multiplication if and only if  $k$  is in the kernel of  $B$  relative to right multiplication. This follows since

$$(k^T B)^T = -Bk \text{ when } B \text{ is skew symmetric.}$$

1.2. The rank of a skew symmetric matrix is even [ 3, page 217].

1.3. Again let  $B$  be an  $n$ -dimensional ( $n \geq 2$ ) skew symmetric matrix and  $C$  be the  $(n - 1)$ -dimensional principal submatrix obtained from  $B$  by removing the first row and the first column. If the  $\ker(B)$  is nondegenerate and is contained in  $\{x \mid x_1 = 0\}$  then  $C$  is singular. This follows since  $k$  in the  $\ker(B)$  implies  $k_1 = 0$  that is  $k = (0, k_2, \dots, k_n)^T$ . Now if we project  $k$  by  $P$  defined by  $Px = (x_2, \dots, x_n)^T$  then  $P$  is a projection from  $n$  to  $n - 1$  dimensional vector spaces. We note that  $k$  in the kernel of  $B$  implies that  $Pk$  is in  $\ker(C)$ . Since  $k^T = (0, k_2, \dots, k_n)$  we have

$$\begin{aligned} k^T B &= (k^T B_1, k^T B_2, \dots, k^T B_n) \\ &= (0, Pk^T C_1, \dots, Pk^T C_{n-1}) = (0, 0, \dots, 0) \end{aligned}$$

where  $B_j$  and  $C_j$  refer to the  $j^{\text{th}}$  column of  $B$  and  $C$ , respectively.

1.4. If the  $\ker(B)$  is contained in  $\{x \mid x_1 = 0\}$  then the  $\dim(\ker(B)) < \dim(\ker(C))$ . To see this let  $n$  be the dimension of  $B$ . We see from the proof of 3) that the  $\dim(\ker(C)) \geq \dim(\ker(B))$  and  $\dim(\ker(C)) \geq 1$ . Suppose that  $\dim(\ker(C)) = \dim(\ker(B))$ . Then  $\text{rank}(C) + \dim(\ker(C)) = n - 1$  and  $\text{rank}(B) + \dim(\ker(B)) = n$ . and so  $\text{rank}(C) = \text{rank}(B) - 1$ . But this impossible since the ranks of both the skew symmetric matrices  $B$  and  $C$  must be even.

Without loss of generality we can assume that the hyperplane of zeros of  $f$  is  $\{x \mid x_1 = 0\}$ . For consider the dynamical system  $x' = Ax + f(x)$  where the zeros of  $f$  are an  $(n - 1)$ -dimensional hyperplane, call it  $H$ . Let  $R$  be a rotation so that  $H$  goes onto the hyperplane  $Y_1 = \{y \mid y_1 = 0\}$  under  $R^{-1}$ . Let  $x = Ry$ . Then the original dynamical system is represented by  $y' = R^{-1}ARy + R^{-1}f(Ry)$ . Note that the zeros of  $R^{-1}f(Ry)$  are precisely the zeros of  $f(x)$  under the rotation  $R^{-1}$ . This is true for any rotation  $R$ . We note that  $Z(R^{-1}f(Ry)) \supseteq R(Z(f(x)))$ . This applies to both  $R$  and  $R^{-1}$ .

Let  $x_0 \neq 0$  be in  $H$ . The hypothesis of our theorem requires that  $x_0^T Ax_0 < 0$ . Then  $y_0 = Rx_0$  and  $y_0^T (R^{-1}AR) y_0 = x_0^T Ax_0 < 0$ . Since  $R$  has an inverse the hypotheses for the dynamical system holds whether represented in terms of  $x$  or  $y$ .

Now assuming that  $Z(f) = \{x \mid x_1 = 0\}$  we notice that  $f$  has a convenient representation. Each coordinate function of  $f$ ,  $f_k$ ,  $k = 1, 2, \dots, n$ , must have  $\{x \mid x_1 = 0\}$  contained in its zero set. If we represent  $f_k$ ,  $k = 1, 2, \dots, n$  by

$$f_k(x) = \sum_{i=1}^n \sum_{j=1}^n b_{k \ i \ j} x_i x_j$$

then

$$f_k(0, x_2, \dots, x_n) = \sum_{i=1}^n \sum_{j=1}^n b_{k \ i \ j} x_i x_j = 0 \quad \text{for all } x_2, x_3, \dots, x_n$$

Therefore,  $b_{k \ i \ j} = 0$  if neither  $i$  nor  $j$  is 1. Or

$$f_k(x) = \sum_{j=1}^n b_{k \ 1 \ j} x_1 x_j = x_1 \sum_{j=1}^n b_{k \ 1 \ j} x_j$$

and

$$f(x) = \begin{bmatrix} x_1 \sum_{j=1}^n b_{1 \ 1 \ j} x_j \\ x_1 \sum_{j=1}^n b_{2 \ 1 \ j} x_j \\ \vdots \\ \vdots \\ x_1 \sum_{j=1}^n b_{n \ 1 \ j} x_j \end{bmatrix} = x_1 \begin{bmatrix} b_{111} & b_{112} & \dots & b_{11n} \\ b_{211} & b_{212} & \dots & b_{21n} \\ \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ b_{n11} & b_{n12} & \dots & b_{n1n} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ \vdots \\ x_n \end{bmatrix}$$

Let  $B$  denote the matrix  $(b_{k \ 1 \ j})_{1 \ 1}^{n \ n}$ . Since  $x^T f(x) = x_1 x^T B x = 0$  for all  $x$

$x^T f(x) = \sum_{j=1}^n b_{j1j} x_1 x_j^2 + \sum_{k=1}^n \sum_{j=1}^n (b_{k1j} + b_{j1k}) x_1 x_j x_k = 0$  which is the zero polynomial having zero coefficients.  $B$  is skew symmetric and we drop the second subscript which is always one and we have the representation

$$f(x) = x_1 \begin{bmatrix} 0 & b_{12} & b_{13} & \dots & b_{1n} \\ -b_{12} & 0 & b_{23} & \dots & b_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -b_{1n} & -b_{2n} & -b_{3n} & \dots & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

We use the Lyapunov function  $V(x) = \frac{1}{2}(x - \alpha)^T (x - \alpha)$  then  $\frac{d(V(x)(t)))}{dt} = x^T A x - \alpha^T f(x) +$  linear terms. For large  $\|x\|$  the quadratic terms  $x^T A x - \alpha^T f(x)$  dominate  $\frac{d(V(x(t)))}{dt}$  therefore our quest is find an  $\alpha$  which turns the quadratic into a negative definite function.

## 2. EXTENSION OF PREVIOUS RESULTS

A sufficient condition for a quadratic dynamical system to be point dissipative has been given [ 1 ] when the dimension is two or three. This condition uses a relation between the quadratic and linear parts of the system when  $f$  is conservative. The following lemma allows us to extend the condition to the case where there exists a positive definite matrix  $S$  such that  $Sf$  is conservative.

Lemma 1. Let

$$x' = Ax + f(x) \quad (2.1)$$

be a quadratic dynamical system for which there exists a positive definite matrix  $S$  such that  $Sf$  is conservative. Then there exists matrix  $H$  for which the change of variables  $y = Hx$  transforms the dynamical system (2.1) into  $y' = By + g(y)$  which has a conservative quadratic term. Furthermore,  $x^T S A x = y^T B y$ .

Proof. We can factor  $S$  by  $S = H^T H$  [3]. Let  $y = Hx$  or  $x = H^{-1}y$ . The system transforms into  $y' = (H A H^{-1})y + H f(H^{-1}y)$  or

$$y' = By + g(y) \quad (2.2)$$

The system (2.2) has a conservative quadratic term since  $y^T g(y) = x^T H^T H f(x) = x^T S f(x) = 0$ . Note that  $x^T S A x = y^T (H^{-1})^T H^T H A H^{-1} y = y^T H A H^{-1} y = y^T B y$ .

When  $Sf$  is conservative then we say that a vector  $\alpha$  is admissible for the dynamical system  $x' = Ax + f(x)$  if  $-x^T S A x + \alpha^T H f(x)$  is a positive definite function where  $S = H^T H$ .

When  $f$  is conservative then the condition for an admissible  $\alpha$  reduces to  $-x^T A x + \alpha^T f(x)$  is positive definite. The proofs of the theorems when the quadratic part of the system is conservative entail demonstrating the existence of an admissible  $\alpha$  for the system.

These results [ 1 ] can be restated as follows:

**Theorem 1.** A quadratic dynamical system  $x' = Ax + f(x)$  is point dissipative when there exists a positive definite matrix  $S$  such that  $Sf$  is conservative and there exists an admissible  $\alpha$ . If the system has dimension 2 or 3,  $Sf$  is conservative and  $z^T S A z < 0$  for any  $z$  which is a nontrivial zero of  $f$  then the system is point dissipative.

**Proof.** We can transform the dynamical system by  $y = Hx$  where  $S = H^T H$  and by Lemma 1 the resulting dynamical system satisfies the hypothesis of the previous theorem. Hence, the resulting dynamical system as well as the original system are both point dissipative.

Another direction of generalizing the past results is to consider nonlinear dynamical systems which have nonquadratic nonlinear terms as well as quadratic terms. Relative to the nonlinear terms there again must exist a positive definite matrix  $S$  such that the nonlinear terms premultiplied by  $S$  are conservative.

**Theorem 2.** When there exists a positive definite matrix  $S$  such that  $Sg$  and the quadratic function  $Sf$  are conservative and

Condition (A) For some admissible  $\alpha$  for  $x' = Ax + f(x)$  there exists an ordered triple of numbers  $(\epsilon, C, M)$  such that  $-\alpha^T H g(x) \leq C \|x\|^{2-\epsilon}$  for all  $x$  with  $\|x\| \geq M$ .  
then

$$x' = Ax + f(x) + g(x)$$

is point dissipative.

Note that condition (A) can be replaced by either of the stronger conditions (B) or (C).

Condition (B) There is an admissible  $\alpha$  for  $x' = Ax + f(x)$  and  $\|g\| = o\|x\|^2$ .

Condition (C) There is an admissible  $\alpha$  for  $x' = Ax + f(x)$  and  $g$  is bounded.

### 3. THE PROOF OF THE THEOREM

**Theorem 3.** A quadratic dynamical system  $x' = Ax + f(x)$ ,  $A$  a matrix and  $f$  a quadratic function, is point dissipative when

- (1) there exists a positive definite matrix  $S$  such that  $Sf$  is conservative,
- (2) the zeros of  $f$  are an  $(n - 1)$ -dimensional hyperplane,

and

- (3)  $z^T S A z < 0$  for any  $z$  which is a nontrivial zero of  $f$ .

**Proof.** From Lemma 1 the system can be transformed into a system which has a conservative quadratic term. Moreover, the zero properties are preserved. So we assume that  $f$  is conservative. We can assume that  $Z(f) = \{ x \mid x_1 = 0 \}$  and  $f(x) = x_1 B x$  where  $B$  is

skew symmetric as shown in Section 1.3. Since  $Z(f) = \{ x \mid x_1 = 0 \}$ ,  $\mathbf{k}$  in the kernel of  $B$  implies  $(0, k_2, \dots, k_n)$  or that  $k_1 = 0$ . Let  $\mathbf{h}$  be a vector which is not in the kernel of  $B$  but  $(h_2, \dots, h_n)^T$  is in the  $\ker(C)$ , see Section 1.4. Here again  $C$  is the principal submatrix of  $B$  formed by removing the first row and column of  $B$ . Moreover, if the rank of  $C \neq 0$ , we can choose  $\mathbf{h}$  so that  $h_1 = 0$ .

Let  $G = \{ \alpha \mid \alpha = t \mathbf{h}, t \text{ a real number} \}$  and we will show  $G$  is contained in the set of admissible  $\alpha$ 's. Note that if  $\alpha$  is in  $G$  then  $\alpha^T f(x) = x_1 t \mathbf{h}^T B x = x_1 t (q, 0, 0, \dots, 0) x = t q x_1^2$  since  $P\mathbf{h}$  is in the  $\ker(C)$ . Also  $q \neq 0$  since  $\mathbf{h}$  is not in the  $\ker(B)$ . Notice that  $q \neq 0$  whether or not the  $\text{rank}(C) = 0$  furthermore, we can choose  $t$  so that  $\alpha^T f(x) \geq 0$  for all  $x$ .

We can restrict our attention to the sphere  $\|x\| = 1$  which is compact. There is a closed cone on a closed cone which contains the hyperplane  $\{ x \mid x_1 = 0 \}$  such that  $x^T A x < 0$  for all  $x \neq 0$  in this cone. Hence, there is an  $\epsilon > 0$  such that  $x^T A x < 0$  on  $\{ x \mid \|x\| = 1 \text{ and } -\epsilon \leq x_1 \leq \epsilon \}$ . Let  $M = \max(x^T A x)$  on  $\{ x \mid \|x\| = 1 \}$ . By picking  $t$  large enough in magnitude we can assure that  $t q \epsilon^2 > M$ . Hence,  $\alpha^T f(x) \geq t q \epsilon^2 > M$  on  $\{ x \mid \|x\| = 1 \text{ and } |x_1| \geq \epsilon \}$ . Thus for all of  $\{ x \mid \|x\| = 1 \}$ ,  $x^T A x - \alpha^T f(x) < 0$ . And so for all nonzero  $x$  we have that  $x^T A x - \alpha^T f(x) < 0$  which implies that the system is point dissipative.

This result can be generalized by adding to the differential equation any conservative function  $g(x)$  whose growth is restricted. The corollary states this condition.

Corollary Let  $g$  and the quadratic function be conservative. If  $Z(f)$  is an  $(n - 1)$ -hyperplane and  $x$  in  $Z(f)$  implies  $x^T A x < 0$  and

Condition (A) for some admissible  $\alpha$  there exists an ordered triple of numbers  $(\epsilon, C, M)$  such that  $-\alpha^T g(x) \leq \|x\|^{2-\epsilon}$  for all  $x$  with  $\|x\| \geq M$ .  
then

$$x' = Ax + f(x) + g(x)$$

is point dissipative.

Note that condition (A) can be replaced by either of the stronger conditions (B) or (C).

Condition (B) There is an admissible  $\alpha$  for  $x' = Ax + f(x)$  and  $\|g\| = o\|x\|^2$ .

Condition (C) There is an admissible  $\alpha$  for  $x' = Ax + f(x)$  and  $g$  is bounded.

#### 4. EXAMPLES

Consider the dynamical system

$$x' = \begin{bmatrix} 2 & -1 & -1 & -1 \\ -1 & -3 & -1 & -1 \\ -1 & -1 & -3 & -1 \\ -1 & -1 & -1 & -3 \end{bmatrix} x + \begin{bmatrix} x_1 x_2 + x_1 x_3 + x_1 x_4 \\ -x_1^2 + x_1 x_3 + x_1 x_4 \\ -x_1^2 - x_1 x_2 + x_1 x_4 \\ -x_1^2 - x_1 x_2 - x_1 x_3 \end{bmatrix}$$

In this example the conservative quadratic function can be written as

$$\mathbf{f}(\mathbf{x}) = x_1 \begin{bmatrix} 0 & 1 & 1 & 1 \\ -1 & 0 & 1 & 1 \\ -1 & -1 & 0 & 1 \\ -1 & -1 & -1 & 0 \end{bmatrix} \mathbf{x} = x_1 \mathbf{Bx}$$

$\mathbf{B}$  is a nonsingular matrix and  $\mathbf{C} = \mathbf{B}_{11}$  is singular. The kernel of  $\mathbf{C}$  is generated by  $(1, -1, 1)$ . Since  $\mathbf{B}$  is nonsingular the zero set of  $\mathbf{f}(\mathbf{x})$  is  $Z(\mathbf{f}) = \{ \mathbf{z} \mid \mathbf{z} = (0, z_2, z_3, z_4) = \{ \mathbf{x} \mid x_1 = 0 \} \}$ . The hypothesis of the theorem hold since

$$(0, z_2, z_3, z_4) \mathbf{A} \begin{bmatrix} 0 \\ z_2 \\ z_3 \\ z_4 \end{bmatrix} = (z_2, z_3, z_4) \mathbf{A}_{11} \begin{bmatrix} z_2 \\ z_3 \\ z_4 \end{bmatrix} \text{ and } \mathbf{A}_{11} = \begin{bmatrix} -3 & -1 & -1 \\ -1 & -3 & -1 \\ -1 & -1 & -3 \end{bmatrix}$$

is a negative definite matrix. We can use  $\alpha^T = (0, t, -t, t)$  and  $\alpha^T \mathbf{f}(\mathbf{x}) = x_1 (0, t, -t, t) \cdot (x_2 + x_3 + x_4, -x_1 + x_3 + x_4, x_1 + x_2 - x_4, -x_1 - x_2 - x_3)^T = tx_1^2$ . So the quadratic term of the Lyapunov function is

$$\mathbf{x}^T \mathbf{A} \mathbf{x} - \alpha^T \mathbf{f}(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x} - tx_1^2 \mathbf{x}^T (\mathbf{A} - \mathbf{Q}) \mathbf{x} = \mathbf{x}^T \begin{bmatrix} 2-t & -1 & -1 & -1 \\ -1 & -3 & -1 & -1 \\ -1 & -1 & -3 & -1 \\ -1 & -1 & -1 & -3 \end{bmatrix} \mathbf{x}$$

This quadratic function is negative definite when  $t > 2.6$ . By the theorem this nonlinear quadratic dynamical system is point dissipative.

Indeed in the example  $q$  turns out to be 1. Therefore,  $\alpha^T \mathbf{f}(\mathbf{x})$  can only change the  $a_{11}$  element of  $\mathbf{A}$ . This turns out to be just what we need and want. Because the zero set of  $\mathbf{f}$  is  $\{(0, z_2, z_3, z_4)\}$  we must have that  $-\mathbf{A}_{11}$  is positive definite. If  $t$  can be chosen so that  $\det(-\mathbf{A} + \mathbf{Q})$  is positive then that will be enough to insure that  $-\mathbf{A} + \mathbf{Q}$  is positive definite.

Let us choose  $t = 3$  and return to the derivative of the Lyapunov function. We can set it equal to zero and have the equation of an ellipsoid which contains the attractor of the system. The equation is

$$\mathbf{x}^T \begin{bmatrix} -1 & -1 & -1 & -1 \\ -1 & -3 & -1 & -1 \\ -1 & -1 & -3 & -1 \\ -1 & -1 & -1 & -3 \end{bmatrix} \mathbf{x} - (0, 3, -3, 3)^T \begin{bmatrix} 2 & -1 & -1 & -1 \\ -1 & -3 & -1 & -1 \\ -1 & -1 & -3 & -1 \\ -1 & -1 & -1 & -3 \end{bmatrix} \mathbf{x} = 0$$

If we use the rotation

$$\mathbf{x} = \mathbf{R} \mathbf{y} = \begin{bmatrix} 0.0000 & 0.0000 & 0.9370 & 0.3493 \\ 0.8165 & 0.0000 & -0.2017 & 0.5410 \\ -0.4082 & 0.7071 & -0.2017 & 0.5410 \\ -0.4082 & 0.7071 & -0.2017 & 0.5410 \end{bmatrix} \mathbf{x}$$

the above equation becomes

$$y^T \begin{bmatrix} -2 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 0 & 0 & -3 + \sqrt{7} & 0 \\ 0 & 0 & 0 & -3 - \sqrt{7} \end{bmatrix} y + (0.3330, 7.9233, 10.61307, -2.1149) y = 0$$

or

$$2(x_1 - 0.16651)^2 + 2(x_2 - 3.9617)^2 + 0.3542(x_3 - 5.3065)^2 + 5.6458(x_4 - 1.0574)^2 = 47.7333$$

The distance to the center is 6.7082 and the half-diameter of the ellipsoid is 11.6070 and our bound is the sum of these two numbers is 18.3162

Consider the dynamical system

$$\dot{x} = \begin{bmatrix} -4 & 4 \\ -4 & 1 \end{bmatrix} x + \begin{bmatrix} 6xy - 4y^2 \\ -6x^2 + 4xy \end{bmatrix} + \begin{bmatrix} 2ye^{-x-y} \\ -2xe^{-x-y} \end{bmatrix}$$

where the linear part  $Ax$  and the quadratic function

$$f(x) = \begin{bmatrix} 6xy - 4y^2 \\ -6x^2 + 4xy \end{bmatrix}$$

are related by  $z^T A z < 0$  when  $z$  is a nontrivial zero of  $f$ . The admissible  $\alpha$ 's are those for which  $x^T A x - \alpha^T f(x)$  is a negative definite function. This is equivalent to the matrix

$$\begin{bmatrix} 4 - 6\alpha_2 & 3\alpha_1 + 2\alpha_2 \\ 3\alpha_1 + 2\alpha_2 & -1 - 4\alpha_1 \end{bmatrix}$$

being positive definite. For the diagonal terms to be positive definite  $\alpha_1 < -1/4$ ,  $\alpha_2 < 2/3$  and for the determinant of the matrix to be positive

$$-9\alpha_1^2 + 12\alpha_1\alpha_2 - 4\alpha_2^2 - 16\alpha_1 + 6\alpha_2 - 4 > 0$$





Figure 1

The shaded region is the set of admissible  $\alpha^T$  relative to  $A$  and  $f$ . Now if for one of these  $\alpha$ 's condition (A) hold the the dynamical system is point dissipative. This is the case when we choose  $\alpha^T = (-1/2, 1/2)$

$$-\alpha^T g(x) = -\left(\frac{-1}{2}, \frac{1}{2}\right) \begin{bmatrix} 2x e^{-x-y} \\ -2y e^{-x-y} \end{bmatrix} = -(x+y) e^{-x-y} \leq \frac{1}{e}$$

and the value of the determinate is 0.75 and the diagonal elements are 1 and 1. Here is the graph [6] of some of the trajectories.



$$\frac{dx}{dt} = -4x + 4y + 6x^2 - 4xy + 2xe^{-x-y}$$

$$\frac{dy}{dt} = -4x + y - 6xy + 4y^2 - 2ye^{-x-y}$$

Figure 2

The attractor above the origin is magnified in the following Figure 3. It indicates that there is an attractor at  $(-0.5, 1)$  surrounded by a limit cycle which is an repeller. There is also an attractor at  $(0, 0)$  and a saddle point near  $(1, 2)$ .




Figure 3

#### BIBLIOGRAPHY

1. BOSE, A. K., A. S. Cover and J. A. Reneke, On Point-Dissipative Systems of Differential Equations with Quadratic Nonlinearity, *Int. J. Math. and Math. Sci.*, 14(1990), 99-110.
2. HALE, J. K., L. T. Magalhaes and W. M. Oliva, An Introduction to Infinite Dimensional Dynamical Systems. Geometric Theory, New York, Berlin, Heidelberg, Tokyo, Applied Mathematical Sciences, Springer-Verlag.
3. HORN, R. A. and C. R. Johnson, Matrix Analysis, Cambridge, Cambridge University Press, 1985.
4. LASALLE, Joseph and Solomon Lefschetz, Stability by Liapunov's Direct Method, with Applications, New York, Academic Press, 1961.
5. LORENZ, E. N., Deterministic non-periodic flow, *J. Atmos. Sci.*, 20(1963), 130-141.
6. Phase Portraits, Version 2.0, Department of Mathematics and Computer Science, Drexel University, 1988.

## Special Issue on Intelligent Computational Methods for Financial Engineering

### Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Lean Yu**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [yulean@amss.ac.cn](mailto:yulean@amss.ac.cn)

**Shouyang Wang**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; [sywang@amss.ac.cn](mailto:sywang@amss.ac.cn)

**K. K. Lai**, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [mskklai@cityu.edu.hk](mailto:mskklai@cityu.edu.hk)