

## RESEARCH NOTES

### RETRACTS IN EQUICONNECTED SPACES

JIN HO KWAK

Department of Mathematics  
Pohang Institute of Science and Technology  
Pohang, 790-600  
Korea

JAEUN LEE

Department of Mathematics  
Kyungpook University  
Taegu, 702-701  
Korea

(Received June 6, 1990)

**ABSTRACT.** The object of this paper is to show that the concepts “retract” and “strong deformation retract” coincide for a subset of an equiconnected space. Also, we have a similar local version in a locally equiconnected space.

**KEY WORDS AND PHRASES.** Retract, strong deformation retract, and equiconnected.

**1991 AMS SUBJECT CLASSIFICATION CODE.** 54C15.

#### 1. INTRODUCTION.

Let  $X$  be a topological space, let  $A$  be a subset of the product space  $X \times X$  and let  $\lambda$  be a map from  $A \times I$  (where  $I$  is the interval) to  $X$ . The map  $\lambda$  is said to have the connecting property on  $A$  if  $\lambda(x, y, 0) = x$ ,  $\lambda(x, y, 1) = y$  for all  $(x, y) \in A$  and  $\lambda(x, x, t) = x$  for all  $t \in I$ ,  $(x, x) \in A$ . A topological space  $X$  is equiconnected (EC) if there is a map  $\lambda$  from  $X \times X \times I$  to  $X$  which has the connecting property on  $X \times X$ . A space  $X$  is locally equiconnected (LEC) if there is a neighborhood  $\Delta(X)$  of the diagonal in  $X \times X$  and a map  $\lambda$  from  $\Delta(X) \times I$  to  $X$  which has the connecting property on  $\Delta(X)$ . A subset  $S$  of  $X$  is called a (strong) neighborhood deformation retract of  $X$  if there is a neighborhood  $N$  of  $S$  such that  $S$  is a (strong) deformation retract of  $N$  over  $X$ .

Now we state a main theorem.

**THEOREM 1.** The following three statements are equivalent for a subset  $S$  of a LEC space  $X$ .

- i)  $S$  is a neighborhood retract of  $X$ ,
- ii)  $S$  is a neighborhood deformation retract of  $X$ ,
- iii)  $S$  is a strong neighborhood deformation retract of  $X$ .

It is well known (c.f., XV 8.1 in [2]) that a subset of a metrizable space  $X$  is a strong deformation retract of  $X$  if it is both a deformation retract of  $X$  and a strong neighborhood deformation retract of  $X$ . We therefore have

**COROLLARY 1.** If  $S$  is a subset of a metrizable LEC space  $X$ , then the following are equivalent.

- i)  $S$  is a deformation retract of  $X$ ,
- ii)  $S$  is a strong deformation retract of  $X$ .

Furthermore, if  $X$  is contractible, the above statements are equivalent to

- iii)  $S$  is a retract of  $X$ .

PROOF. We need only to show the implication  $i) \Rightarrow iii)$ . By assuming  $i)$ , there is an open neighborhood  $N$  of  $S$  such that  $S$  is a retract of  $N$  with retraction  $r : N \rightarrow S$ . Since every open subset of an LEC space is LEC,  $N$  is LEC. Let  $\lambda : \Delta(N) \times I \rightarrow N$  be a local equiconnection, where  $\Delta(N)$  is an open neighborhood of the diagonal in  $N \times N$ . We define a map  $F : N \rightarrow N \times N$  by  $F(x) = (x, r(x))$  for all  $x \in N$ . Then,  $F$  is continuous and  $F^{-1}(\Delta(N))$  is a neighborhood of  $S$  in  $N$ , and the map  $G : F^{-1}(\Delta(N)) \times I \rightarrow X$  defined by  $G(x, t) = \lambda(F(x), t)$  for all  $(x, t) \in F^{-1}(\Delta(N)) \times I$  is the desired homotopy to complete the proof.

A similar argument gives

**THEOREM 2.** If  $S$  is a subset of an EC space  $X$ , then the following are equivalent:

- i)  $S$  is a retract of  $X$ ,
- ii)  $S$  is a deformation retract of  $X$ ,
- iii)  $S$  is a strong deformation retract of  $X$ .

#### REFERENCES

1. HIMMELBERG, C. J. Some theorems on equiconnected and locally equiconnected spaces, Trans. Amer. Math. Soc. 115 (1965), 43-53.
2. DUGUNDJI, J. Topology, Allyn and Bacon Inc., Boston, 1966.

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)