

**REMARKS ON ORTHOGONAL POLYNOMIALS WITH RESPECT TO
VARYING MEASURES AND RELATED PROBLEMS**

XIN LI

Department of Mathematics
University of Central Florida
Orlando, FL 32816

(Received March 26, 1992 and in revised form May 4, 1992)

ABSTRACT. We point out the relation between the orthogonal polynomials with respect to (w.r.t.) varying measures and the so-called orthogonal rationals on the unit circle in the complex plane. This observation enables us to combine different techniques in the study of these polynomials and rationals. As an example, we present a simple and short proof for a known result on the weak-star convergence of orthogonal polynomials w.r.t. varying measures. Some related problems are also considered.

KEY WORDS AND PHRASES. Orthogonal polynomial and rational, weak-star convergence.

1991 AMS SUBJECT CLASSIFICATION CODES. Primary 42A05, Secondary 33C45, 41A28.

1. INTRODUCTION.

Let σ be a finite positive measure on the unit circle $\Gamma := \{z \in C \mid |z| = 1\}$, and let $w_n(z) = z^n + \dots$ be a sequence of polynomials whose zeros $\{z_{n,k}\}_{k=1}^n$ all lie in the closed unit disk $|z| \leq 1$. Assume

$$\int_{\Gamma} \frac{d\sigma}{|w_n(z)|^2} < \infty,$$

then, for each n , we can consider the orthonormal polynomials $\varphi_{n,m}(z) = \alpha_{n,m} z^m + \dots$ ($\alpha_{n,m} > 0$) w.r.t. $d\sigma/|w_n(z)|^2$, i.e., polynomials satisfying

$$\frac{1}{2\pi} \int_{\Gamma} z^{-j} \varphi_{n,m}(z) \frac{d\sigma}{|w_n(z)|^2} = 0, \quad j = 0, 1, \dots, m-1,$$

and

$$\frac{1}{2\pi} \int_{\Gamma} |\varphi_{n,m}(z)|^2 \frac{d\sigma}{|w_n(z)|^2} = 1.$$

The sequences $\{\varphi_{n,m}(z)\}_{m=0}^{\infty}$ are called orthogonal polynomials w.r.t. varying measures $d\sigma/|w_n(z)|^2$ ($n = 1, 2, \dots$). They appear in the study of simultaneous Padé approximation and related problems. The following weak-star convergence result plays an important role in the study of convergence of $\{\varphi_{n,n}(z)\}_{n=1}^{\infty}$.

THEOREM 1. (Lopez, [8]) If the zeros of $w_n(z)$ satisfy the condition $\lim_{n \rightarrow \infty} \sum_{i=1}^n (1 - |z_{n,i}|) = \infty$, then

$$\left| \frac{w_n(z)}{\varphi_{n,n}(z)} \right|^2 |dz| \rightarrow d\sigma, \quad (n \rightarrow \infty),$$

in the weak-star topology.

One nice thing about this result is that it holds without any condition on the finite positive measure $d\sigma$. We will present a simple and short proof of Theorem 1. But before we get to the proof, we remark that the so-called orthogonal rationals (with prescribed poles) are also of current interest, especially in problems related to electrical engineering. (See, e.g., [2, 3, 4, 5]) These applications were more concerned with the algebraic properties of orthogonal rationals. It turns out that orthogonal rationals are just special case of orthogonal polynomials w.r.t. varying measures. It is the main purpose of this note to point out this relation. To see this relation, let us recall the definition of orthogonal rationals with respect to $d\sigma$.

Let $\{z_i\}_{i=1}^\infty$ be points in $|z| \leq 1$, and denote $Q_n(z) := \prod_{i=1}^n (z - z_i)$.

DEFINITION 2. (cf. [2]) For $n = 1, 2, \dots$, the rationals $\frac{P_n(z)}{Q_n(z)}$ with $P_n(z) = \alpha_n z^n + \dots (\alpha_n > 0)$ satisfying

$$\frac{1}{2\pi} \int_{\Gamma} \frac{P_n(z)}{Q_n(z)} \overline{r(z)} d\sigma = 0$$

for $r(z) \in \{\frac{q(z)}{Q_n(z)} : q \in \mathfrak{P}_{n-1}\}$, and

$$\frac{1}{2\pi} \int_{\Gamma} \left| \frac{P_n(z)}{Q_n(z)} \right|^2 d\sigma = 1,$$

are called orthogonal rationals w.r.t. $d\sigma$.

Now, from Definition 2, it can be seen that with $Q_n(z)$ treated as $w_n(z)$, the corresponding $\varphi_{n,n}(z)$ is $P_n(z)$. So orthogonal rationals are indeed special case of orthogonal polynomials w.r.t. varying measures.

We go back to consider orthogonal polynomials w.r.t. varying measures. The following result, which is essentially proved in the orthogonal rational setting (cf. e.g., [2]), will help us to simplify the proof of Theorem 1 in [8].

LEMMA 3. If r and s belong to $\mathfrak{R}(w_n) := \{p(z)/w_n(z) : p \in \mathfrak{P}_n\}$, then

$$\frac{1}{2\pi} \int_{\Gamma} r(z) \overline{s(z)} \left| \frac{w_n(z)}{\varphi_{n,n}(z)} \right|^2 |dz| = \frac{1}{2\pi} \int_{\Gamma} r(z) \overline{s(z)} d\sigma.$$

PROOF. By [6, formula (1.20)], we have

$$\frac{1}{2\pi} \int_{\Gamma} \frac{p(z) \overline{q(z)}}{|\varphi_{n,n}(z)|^2} |dz| = \frac{1}{2\pi} \int_{\Gamma} p(z) \overline{q(z)} \frac{d\sigma}{|w_n(z)|^2}$$

for all $p, q \in \mathfrak{P}_n$. So,

$$\frac{1}{2\pi} \int_{\Gamma} \frac{p(z)}{w_n(z)} \overline{\left(\frac{q(z)}{w_n(z)} \right)} \left| \frac{w_n(z)}{\varphi_{n,n}(z)} \right|^2 |dz| = \frac{1}{2\pi} \int_{\Gamma} \frac{p(z)}{w_n(z)} \overline{\left(\frac{q(z)}{w_n(z)} \right)} d\sigma,$$

which implies the lemma.

REMARKS. (i) Taking $r = s = 1$ in Lemma 3 gives us

$$\int_{\Gamma} \left| \frac{w_n(z)}{\varphi_{n,n}(z)} \right|^2 |dz| = \int_{\Gamma} d\sigma. \quad (1)$$

(ii) Lemma 3 is equivalent to the following:

$$\frac{1}{2\pi} \int_{\Gamma} \frac{z^{-n} p_{2n}(z)}{|w_n(z)|^2} \left| \frac{w_n(z)}{\varphi_{n,n}(z)} \right|^2 |dz| = \frac{1}{2\pi} \int_{\Gamma} \frac{z^{-n} p_{2n}(z)}{|w_n(z)|^2} d\sigma, \quad (2)$$

for all $p_{2n} \in \mathcal{P}_{2n}$.

We also need the following result about weighted approximation with varying weights in L_{∞} -norm.

LEMMA 4. The space

$$T_n := \left\{ \frac{z^{-n} p_{2n}(z)}{|w_n(z)|^2} : p_{2n} \in \mathcal{P}_{2n} \right\}$$

is dense in $C(\Gamma)$ in the L_{∞} -norm if and only if

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n (1 - |z_{n,i}|) = \infty.$$

PROOF. This is a simple generalization of some known results (see, e.g., [1, p. 244] and [2, Theorem 8.1.2]). We only mention the following fact (cf. [1, p. 243])

$$\inf_{p \in \mathcal{P}_{2n+m-1}} \left\| \frac{z^{2n+m} + p(z)}{\prod_{z_{n,i} \neq 0} (z - z_{n,i})(z - 1/z_{n,i})} \right\| = \prod_{z_{n,i} \neq 0} |z_{n,i}|.$$

PROOF OF THEOREM 1. Now the proof of Theorem 1 follows from (1), (2), and Lemma 4 by using a standard argument (cf. e.g., [9, p. 248]).

Recently, in the study of frequency analysis (see, e.g., [7]), the following orthogonal polynomials w.r.t. varying measures appear naturally, i.e., polynomials $\varphi_{n,m}(z) = \beta_{n,m} z^m + \dots + \beta_{n,m} > 0$ satisfying

$$\frac{1}{2\pi} \int_{\Gamma} z^{-j} \psi_{n,m}(z) |w_n(z)|^2 d\sigma = 0, \quad j = 0, 1, \dots, m-1$$

and

$$\frac{1}{2\pi} \int_{\Gamma} |\psi_{n,m}(z) w_n(z)|^2 d\sigma = 1.$$

For convenience, let us assume that the zeros of $w_n(z)$ all lie outside the unit circle Γ , i.e., $|z_{n,i}| > 1$. The asymptotics of $\psi_{n,m}(z) w_n(z)$ are wanted in application ([7]). Here we will not discuss this problem but instead point out the relation of these polynomials $\{\psi_{n,m}\}$ with the following ones $\{\phi_{n,m}\}$: Assume $A_{n,i} > 0$ ($1 \leq i \leq n, n \geq 1$) are given, then define polynomials $\phi_{n,m}(z) = \gamma_{n,m} z^m + \dots + \gamma_{n,m} > 0$ satisfying

$$\frac{1}{2\pi} \int_{\Gamma} z^{-j} \phi_{n,m}(z) d\sigma + \sum_{i=1}^n A_{n,i} z_{n,i}^{-j} \phi_{n,i}(z_{n,i}) = 0$$

for $j = 0, 1, \dots, m+n-1$, and

$$\frac{1}{2\pi} \int_{\Gamma} |\phi(z)|^2 d\sigma + \sum_{i=1}^n A_{n,i} |\phi_{n,i}(z_{n,i})|^2 = 1.$$

The polynomials $\{\phi_{n,m}\}_{m=0}^{\infty}$ are the orthonormal polynomials w.r.t. measure $d\sigma/(2\pi) + \sum_{i=1}^n A_{n,i} \delta_{z_{n,i}}$ having finitely many mass points off the unit circle Γ (where δ_z denotes the point unit measure with support at z). We have

THEOREM 5. For each n fixed, there hold

$$\lim_{\substack{A_{n,i} \rightarrow \infty \\ 1 \leq i \leq n}} \phi_{n,m}(z) = \psi_{n,m}(z)w_n(z),$$

locally uniformly in C , where $m = 0, 1, 2, \dots$.

PROOF. We use a normal family argument. For n and m chosen, since $\int_{\Gamma} |\phi_{n,m}(z)|^2 d\sigma \leq 2\pi$, then the set $\Phi = \{\phi_{n,m} : A_{n,i} > 0, 1 \leq i \leq n\}$ as a subset of finite dimensional normed space is a uniformly bounded set. Thus Φ as a set of analytic functions (polynomials) is a normal family over C . Let $f(z) \in \mathcal{P}_{n+m}$ be a limit of a subsequence $\Phi' \subset \Phi$, then since $\sum_{i=1}^n A_{n,i} |\phi_{n,m}(z_{n,i})|^2 \leq 1$, we have

$$f(z_{n,i}) = \lim_{\substack{\phi_{n,m} \in \Phi' \\ A_{n,i} \rightarrow \infty}} \phi_{n,m}(z_{n,i}) = 0$$

for $i = 1, 2, \dots, n$. So $f(z) = w_n(z)g(z)$ with some $g(z) \in \mathcal{P}_m$. Now we just need to use the extremal property of (monic) orthogonal polynomials to assert that $g(z) = \psi_{n,m}(z)$. This completes the proof of Theorem 5.

REFERENCES

1. ACHIESER, N.I., Theory of Approximation, Frederick Ungar Publ. Co., New York, 1956.
2. BULTHEEL, A., GONZALEZ-VERA, P., HENDRIKSEN, E., & NJASTAD, O., A Szegő theory for rational functions, manuscript, 1991.
3. DEWILDE, P., & DYM, H., Lossless inverse scattering, digital filters, and estimation theory, IEEE Trans. Inform. Theory, Vol. IT-30 (1984), 644-663.
4. DEWILDE, P., & DYM, H., Schur recursions, error formulas, and convergence of rational estimators for stationary stochastic sequences, IEEE Trans. Inform. Theory, Vol. IT-27 (1981), 446-461.
5. DJRBASHIAN, M.M., A survey on the theory of orthogonal systems and some open problems, Orthogonal Polynomials: Theory and Applications, (P. Nevai ed.), Kluwer Acad. Publ., Boston, 1990, 135-146.
6. GERONIMUS, Y.A., Orthogonal Polynomials. Consultants Bureau, New York, 1961.
7. JONES, W.B., NJASTAD, O., & SAFF, E.B., Szegő polynomials associated with Wiener-Levinson filters, J. Comp. Appl. Math. 32 (1990), 387-406.
8. LOPEZ LAGOMASINO, G., Asymptotics of polynomials orthogonal with respect to varying measures, Constr. Approx. 5 (1989), 199-219.
9. MÁTÉ, A., NEVAI, P. & TOTIK, V., Strong and weak convergence of orthogonal polynomials, Amer. J. Math. 109 (1987), 239-282.

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru