

ON TOTALLY UMBILICAL CR-SUBMANIFOLDS OF A KAEHLER MANIFOLD

M.A. BASHIR

Mathematics Department
College of Science
King Saud University
P.O. Box 2455
Riyadh 11451
Saudi Arabia

(Received June 4, 1991 and in revised form April 21, 1992)

ABSTRACT. Let M be a compact 3-dimensional totally umbilical CR -submanifold of a Kaehler manifold of positive holomorphic sectional curvature. We prove that if the length of the mean curvature vector of M does not vanish, then M is either diffeomorphic to S^3 or RP^3 or a lens space $L^3_{p,q}$.

KEY WORDS AND PHRASES. CR -submanifolds. Totally umbilical submanifolds. Kaehler manifold.

1991 AMS SUBJECT CLASSIFICATION CODES. Primary 53C40. Secondary 53C55.

1. INTRODUCTION.

Totally umbilical CR -submanifolds of a Kaehler manifold have been considered by Bejancu [2], Blair, and Chen [3]. Recently Deshmukh and Husain [5] have also studied these submanifolds. In fact, they have proved a classification theorem when the dimension of the submanifold M is ≥ 5 . In this paper we consider 3-dimensional totally umbilical CR -submanifolds of a Kaehler manifold. For this case we have obtained the following theorem:

THEOREM 1.1. Let M be a compact 3-dimensional totally umbilical CR -submanifold of a Kaehler manifold \bar{M} , of positive holomorphic sectional curvature. If the length of the mean curvature vector of M does not vanish then M is diffeomorphic either to S^3 , RP^3 or the lens space $L^3_{p,q}$.

2. PRELIMINARIES.

Let \bar{M} be an m -dimensional Kaehler manifold with almost complex structure J . A $(2p+q)$ -dimensional submanifold M of \bar{M} is called a CR -submanifold if there exists a pair of orthogonal complementary distributions D and \bar{D} such that $JD = D$ and $J\bar{D} \subset \nu$, where ν is the normal bundle of M and $\dim \bar{D} = q$ [1]. Thus the normal bundle ν splits as $\nu = J\bar{D} \oplus \mu$, where μ is invariant subbundle of ν under J . A CR -submanifold is said to be proper if neither $D = \{0\}$ nor $\bar{D} = \{0\}$.

We denote by $\bar{\nabla}$, ∇ , $\overset{\perp}{\nabla}$ the Riemannian connection on \bar{M} , M and the normal bundle respectively. They are related by

$$\bar{\nabla}_X Y = \nabla_X Y + h(X, Y) \quad (2.1)$$

$$\bar{\nabla}_X N = -A_N X + \overset{\perp}{\nabla}_X N, \quad N \in \nu \quad (2.2)$$

where $h(X, Y)$ and $A_N X$ are the second fundamental forms which are related by

$$g(h(X, Y), N) = g(A_N X, Y) \quad (2.3)$$

Now a CR -submanifold is said to be totally umbilical if

$$h(X, Y) = g(X, Y)H$$

where $H = \frac{1}{n}(\text{trace } h)$ is the main curvature vector. If M is totally umbilical CR -submanifold, then equations (2.1) and (2.2) become

$$\bar{\nabla}_X Y = \nabla_X Y + g(X, Y)H \quad (2.4)$$

$$\bar{\nabla}_X N = -g(H, N)X + \frac{1}{n} \nabla_X N \quad (2.5)$$

For $X, Y, Z, W \in X(M)$, the equation of Gauss is given by

$$R(X, Y; Z, W) = \bar{R}(X, Y; Z, W) + g(h(X, W), h(Y, Z)) - g(h(X, Z), h(Y, W)) \quad (2.6)$$

3. 3-DIMENSIONAL CR -SUBMANIFOLD OF A KAEHLER MANIFOLD.

(A) Let M be a compact totally umbilical 3-dimensional CR -submanifold of a Kaehler manifold \bar{M} . If $\dim D = 0$, then M will be totally real. Therefore, we assume that $\dim D \neq 0$. Since M is 3-dimensional it follows that $\dim D = 2$. We can then choose a frame field $\{X, JX, Z\}$ on M , where $X \in D$ and $Z \in D^\perp$. We first have the following:

LEMMA 1. Let $\{X, JX, Z\}$ be a frame field on M , $X \in D$, $Z \in D^\perp$. Then $\nabla_Z Z = 0$ and $H \in JD^\perp$.

PROOF. Using (2.4) and (2.5) in the equation $\bar{\nabla}_Z JZ = J\bar{\nabla}_Z Z$, we obtain

$$-g(H, JZ)JZ + J\bar{\nabla}_Z JZ = -\nabla_Z Z - h(Z, Z) \quad (3.1)$$

Taking inner produce in (3.1) with $W \in D$ we have

$$g(\nabla_Z Z, W) = 0 \quad W \in D \quad (3.2)$$

From (3.2) we have $\nabla_Z Z \in D^\perp$. Since $g(Z, Z) = 1$, we also have $\nabla_Z Z \in D$. Therefore $\nabla_Z Z = 0$.

Now for $X, Y \neq 0$ in D we use (2.4) and the equation $J\bar{\nabla}_X Y = \bar{\nabla}_X JY$ to get

$$J\nabla_X Y + g(X, Y)JH = \nabla_X JY + g(X, JY)H \quad (3.3)$$

Taking inner produce in (3.1) with $N \in \mu$ we have

$$g(X, Y)g(JH, N) = g(X, JY)g(H, N) \quad (3.4)$$

In particular if we let $Y = JX$ in (3.4) we get

$$\|X\|g(H, N) = 0, \quad N \in \mu. \quad \text{Therefore } H \in JD^\perp. \quad (3.5)$$

Consider the frame field $\{X, JX, Z\}$ on M . Since M is totally umbilical the equation $h(Y, W) = g(Y, W)H$ for $Y, W \in X(M)$ implies that

$$h(X, JX) = h(X, Z) = h(JX, Z) = 0$$

$$h(X, X) = h(JX, JX) = h(Z, Z) = H \equiv \alpha JZ \quad (3.6)$$

for some smooth function α on M , since $H \in JD^\perp$.

Using (2.3) with $N = JZ$ we get

$$AX = \alpha X, \quad AJX = \alpha JX, \quad AZ = \alpha Z \quad (3.7)$$

So the frame field $\{X, JX, Z\}$ diagonalizes A . Now using the equation $(\bar{\nabla}_X J)(X) = 0$ and $(\bar{\nabla}_{JX} J)(X) = 0$ with the help of (3.6) we get

$$g(\bar{\nabla}_X X, Z) = 0, \quad g(\bar{\nabla}_{JX} X, Z) = 0 \quad (3.8)$$

Also using the equation $\nabla_Z Z = 0$ from Lemma 1 we have

$$g(\bar{\nabla}_Z X, Z) = 0, \quad g(\bar{\nabla}_Z JX, Z) = 0 \quad (3.9)$$

Then using the equation $(\bar{\nabla}_X J)(Z) = 0$ and (3.7) we obtain

$$g(\bar{\nabla}_X Z, X) = 0, \quad g(\bar{\nabla}_X Z, JX) = \alpha \quad (3.10)$$

and using the equation $(\bar{\nabla}_{JX} J)(Z) = 0$ we have

$$g(\bar{\nabla}_{JX} Z, X) = -\alpha, \quad g(\bar{\nabla}_{JX} Z, JX) = 0 \quad (3.11)$$

Using equations (3.8), (3.9), (3.10), and (3.11) one can write the following equations for the frame field $\{X, JX, Z\}$:

$$\begin{aligned} \bar{\nabla}_X Z &= \alpha JX, & \bar{\nabla}_{JX} Z &= -\alpha X, & \bar{\nabla}_Z Z &= 0 \\ \bar{\nabla}_X X &= \alpha JX, & \bar{\nabla}_{JX} X &= -bJX + \alpha Z, & \bar{\nabla}_Z X &= cJX \\ \bar{\nabla}_X JX &= -aX - \alpha Z, & \bar{\nabla}_{JX} JX &= bX, & \bar{\nabla}_Z JX &= -cX \end{aligned} \quad (3.12)$$

for some smooth functions a, b and c .

Now we are ready to prove the following:

LEMMA 2. For the frame field $\{X, JX, Z\}$ we have

- (i) $R(X, Z; Z, X) = \|H\|^2$
- (ii) $R(X, JX; JX, X) = \bar{R}(X, JX; JX, X) + \|H\|^2$
- (iii) $R(Z, JX, JX, Z) = \|H\|^2$

PROOF. Using equations (3.12) in the equation

$R(X, Z; Z, X) = g(\bar{\nabla}_X \bar{\nabla}_Z Z - \bar{\nabla}_Z \bar{\nabla}_X Z, -\bar{\nabla}_{[X, Z]} Z, X)$, we obtain (i) and (iii). (ii) follows from the Gauss equation (2.6) and the equation $h(X, Y) = g(X, Y)H$.

PROOF OF THE THEOREM. Since $\bar{R}(X, JX; JX, X) > 0$ and $\|H\| \neq 0$ it follows from (i), (ii), and (iii) of Lemma 2 that all plane sections of M have strictly positive sectional curvature. Therefore, the Ricci-curvature of M is strictly positive. Hence by Hamilton's theorem (cf. [4]) it follows that M is diffeomorphic to either S^3, RP^3 or the lens space $L^3_{p, q}$.

ACKNOWLEDGEMENT. The author would like to thank Dr. Sharief Deshmukh for useful discussion while preparing this paper.

REFERENCES

1. BEJANCU, A. *CR-submanifolds of a Kaehler manifold*, *Proc. Amer. Math. Soc.*, **69** (1978), 135-142.
2. BEJANCU, A. *Umbilical CR-submanifolds of a Kaehler manifold*, *Rend. Mat.* **13** (1980), 431-466

3. BLAIR, D.E. and CHEN, B.Y., On *CR*-submanifolds of Hermitian manifolds, *Israel J. Math.* **34** (1980), 353-363
4. HAMILTON, R.S., Three manifolds with positive Ricci Curvature, *J. Differential Geom.* **17** (1982), 255-306
5. DESHMUKH, S. and HUSAIN, S., Totally umbilical *CR*-submanifolds of a Kaehler manifold, *Kodai Math. J.* **9** (1986), 425-429.

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru