

A TYCHONOFF NON-NORMAL SPACE

V. TZANNES

Department of Mathematics
University of Patras
Patras 26110, Greece

(Received July 13, 1992 and in revised form February 3, 1993)

ABSTRACT. A Tychonoff non-normal space is constructed which can be used for the construction of a regular space on which every weakly continuous (hence every θ -continuous or η -continuous) map into a given space is constant.

KEY WORDS AND PHRASES. Tychonoff, non-normal, weakly, θ -, η -continuous maps.

1980 AMS SUBJECT CLASSIFICATION CODE. 54D15, 54C30.

1. INTRODUCTION.

We construct for every Hausdorff space R a Tychonoff non-normal space S such that if f is a weakly continuous map of S into R then there exist two closed subsets $K', L', K' \cap L' = \emptyset$ such that $f(K') = f(L') = \{r\}$, $r \in R$. Therefore, applying the method of Jones [1], we can first construct a regular space containing two points $-\infty$, $+\infty$ such that $f(-\infty) = f(+\infty)$, for every weakly continuous map f of this space into R and then, applying the method of Iliadis and Tzannes [2], a regular space on which every weakly continuous (hence every θ -continuous or η -continuous (Dickman, Porter and Rubin [3])) map into R is constant. The construction of S is a modification of the space $T_1(R)$ in Iliadis and Tzannes [2]. For regular spaces on which every continuous map into a given space is constant see also Armentrout [4], Brandenburg and Mysiak [5], van Douwen [6], Herrlich [7], Hewitt [8], Tzannes [9] and Jounglove [10]. A map $f : X \rightarrow Y$, where X, Y are topological spaces is called 1) weakly continuous if for every $x \in X$ and U open neighbourhood of $f(x)$ there exists an open neighbourhood V of x , such that $f(V) \subseteq \text{Cl}U$, 2) θ -continuous if for every $x \in X$ and open neighbourhood U of $f(x)$, there is an open neighbourhood V of x such that $f(\text{Cl}V) \subseteq \text{Cl}U$ 3) η -continuous if for every regular-open sets U, V of Y ,

$$(i) \quad f^{-1}(U) \subseteq \text{IntCl}f^{-1}(U)$$

$$(ii) \quad \text{IntCl}f^{-1}(U \cap V) \subseteq \text{IntCl}f^{-1}(U) \cap \text{IntCl}f^{-1}(V).$$

Every η -continuous is θ -continuous (Dickman, Porter and Rubin [3, Proposition 3.3. (c)]) and every θ -continuous is obviously weakly continuous.

We denote 1) by $|X|$ the cardinality, of X , 2) by $\psi(X) = \sup\{\psi(X, x) : x \in X\}$ the pseudocharacter of X , where $\psi(X, x)$ is the pseudocharacter of X at x , that is the minimal cardinality of pseudobases of x . (The set U_α consisting of open neighbourhoods of x , is called a pseudobasis if $\bigcap U_\alpha = \{x\}$), 3) by $\psi^+(X)$ the smallest cardinal number greater than $\psi(X)$.

2. THE SPACE S.

Let R be a Hausdorff space and K, L two uncountable sets such that $|K| = |L| = \aleph > |R|$.

For every $k_i \in K$ (resp. $l_i \in L$) we consider an uncountable set K_i (resp. L_i) and a set M such that $|K_i| = |L_i| = |M| \geq \psi^+(R)$. On the set $S = M \cup K_i \cup L_i$ we define the following topology: Every point belonging to K_i, L_i is isolated. For every $k_i \in K$ (resp. $l_i \in L$) a basis of open neighbourhoods are the sets $O(k_i) = \{k_i\} \cup C_i$ (resp. $O(l_i) = \{l_i\} \cup D_i$), where C_i, D_i consist of all but finite number of elements of K_i, L_i , respectively. For every point $m \in M$ a basis of open neighbourhoods are the sets $O(m) = \{m\} \cup P \cup Q$, where P, Q contain all but finite number of elements of the sets $\{h_i(m) : i \in I\}, \{g_i(m) : i \in I\}$, respectively, where I is an index set, $|I| = \aleph_0$ and h_i, g_i are one-to-one maps of M onto K_i, L_i , respectively.

One can show that the space S is Tychonoff and non-normal.

Let f be a weakly continuous map of S into R . Since $|K| > |R|$, it follows that for some $r_1 \in R$ there exists $K' \subseteq K$ such that $|K'| = |K|$ and $f(K') = \{r_1\}$. Let $\{k_n : n = 1, 2, \dots\}$ be a countable subset of K' . Since for every open neighbourhood U of r_1 the set $f^{-1}(ClU)$ contains an open neighbourhood of $k_n, n = 1, 2, \dots$, it follows that $|K_n \setminus f^{-1}(r_1)| \leq \psi(R, r_1)$. Consequently, if h_n is the one-to-one map of M onto K_n then $|h_n^{-1}(K_n \setminus f^{-1}(r_1))| \leq \psi(R, r_1)$ and hence $|\bigcup_{n=1}^{\infty} h_n^{-1}(K_n \setminus f^{-1}(r_1))| \leq \psi(R, r_1)$. Repeating all the above for the set L we have that for some $r_2 \in R$ there exist $L' \subseteq L, |L'| = |L|, f(L') = \{r_2\}$ and a countable subset $\{l_n : n = 1, 2, \dots\} \subseteq L'$ such that if V is an open neighbourhood of r_2 then $|L_n \setminus f^{-1}(r_2)| \leq \psi(R, r_2)$ and hence $|\bigcup_{n=1}^{\infty} g_n^{-1}(L_n \setminus f^{-1}(r_2))| \leq \psi(R, r_2)$. Therefore if $M' = \bigcup_{n=1}^{\infty} (h_n^{-1}(K_n \setminus f^{-1}(r_1)) \cup g_n^{-1}(L_n \setminus f^{-1}(r_2)))$ then $M \setminus M' \neq \emptyset$. Let $m \in M \setminus M'$ and ClW be a closed neighbourhood of $f(m)$ such that $r_1, r_2 \notin ClW$. There exists an open neighbourhood $O(m)$ of m such that $f(O(m)) \subseteq ClW$, while for every $n = 1, 2, \dots, h_n(m) \in f^{-1}(r_1), g_n(m) \in f^{-1}(r_2)$ which imply that $f(m) = r_1 = r_2$.

REFERENCES

1. JONES, F.B. Hereditarily separable, non-completely regular spaces, *Proceedings of the Blacksburg Virginia Topological Conference*, Springer-Verlag (375), 149-151.
2. ILIADIS, S. and TZANNES, V. Spaces on which every continuous map into a given space is constant, *Can. J. Math.* 38 (1986), 1281-1296.
3. DICKMAN, R.F. Jr., PORTER, J.R. and RUBIN, L.R. Completely regular absolutes and projective objects, *Pacific J. Math.*, 94 (2) (1981), 277-295.
4. ARMENTROUT, S. A Moore space on which every real-valued continuous function is constant, *Proc. Amer. Math. Soc.* 12 (1961), 106-109.
5. BRANDENBURG, H. and MYSIOR, A. For every Hausdorff space Y there exists a non-trivial Moore space on which all continuous functions into Y are constant, *Pacific J. Math.* 1 (1984), 1-8.
6. VAN DOUWEN, E.K. A regular space on which every continuous real-valued function is constant, *Nieuw Archief voor Wiskunde* 20 (1972), 143-145.
7. HERRLICH, H. Wann sind alle stetigen Abbildungen in Y Konstant? *Math. Zeitschr.* 90 (1965), 152-154.
8. HEWITT, E. On two problems of Urysohn, *Annals of Mathematics* 47 (1946), 503-509.
9. TZANNES, V. A Moore strongly rigid space, *Can. Math. Bull.* (34) (4) (1991), 547-552.
10. YOUNGLOVE, J.N. A locally connected, complete Moore space on which every real-valued continuous function is constant, *Proc. Amer. Math. Soc.* 20 (1969) 527-530

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru