

## COMPACT DIAGONAL LINEAR OPERATORS ON BANACH SPACES WITH UNCONDITIONAL BASES

YUN SUNG CHOI and SUNG GUEN KIM

Department of Mathematics  
Pohang Institute of Science and Technology  
Pohang, Korea

(Received April 13, 1992 and in revised form July 3, 1992)

**ABSTRACT.** Let  $E$  and  $F$  be Banach spaces with equivalent normalized unconditional bases. In this note we show that a bounded diagonal linear operator  $T : E \rightarrow F$  is compact if and only if its entries tend to 0, using the concept of weak uniform continuity.

**KEY WORDS and PHRASES.** Weakly uniformly continuous, diagonal operators, compact operators, unconditional bases.

**1992 AMS SUBJECT CLASSIFICATION CODES.** 47B05, 47B37.

### 1. Introduction.

Let  $E$  and  $F$  be two complex Banach spaces and  $E^*$  be the dual space of  $E$ . A function  $f : E \rightarrow F$  is said to be weakly uniformly continuous on bounded subsets of  $E$  if for each bounded set  $B \subset E$  and  $\epsilon > 0$ , there are  $\phi_1, \dots, \phi_k \in E^*$  and  $\delta > 0$  such that if  $x, y \in B$ ,  $|\phi_i(x - y)| \leq \delta$  ( $i = 1, \dots, k$ ), then  $\|f(x) - f(y)\| \leq \epsilon$ . R.M. Aron and J.B. Prolla [1] showed that a bounded linear operator  $T : E \rightarrow F$  is compact if and only if  $T$  is weakly uniformly continuous on bounded subsets of  $E$ . Applying this result we generalize the following well-known Hilbert space fact to a Banach space with an unconditional basis: A diagonal bounded linear operator is compact if and only if its entries tend to 0. See, for example, [2, Proposition 4.6].

We recall some relevant definitions and results about a Banach space with an unconditional basis. Let  $E$  be a complex Banach space with an unconditional basis  $(e_n)$ . For every choice of signs  $\theta = (\theta_n)$ , we have a bounded linear operator  $M_\theta$  on  $E$  defined by

$$M_\theta(\sum a_n e_n) = \sum a_n \theta_n e_n. \quad (1.1)$$

The uniform bounded principle implies that the number  $K = \sup \|M_\theta\|$  is finite, which is called the unconditional constant of  $(e_n)$ . Then for every choice of a complex sequence  $(a_n)$  such that  $\sum a_n e_n$  converges and every choice of a bounded complex sequence  $(\alpha_n)$ , we have

$$\|\sum \alpha_n a_n e_n\| \leq 2K(\sup |\alpha_n|) \|\sum a_n e_n\|. \quad (1.2)$$

For details see [3].

## 2. Main Results.

THEOREM 1. Let  $(\epsilon_n)$  and  $(f_n)$  be equivalent normalized unconditional bases of  $E$  and  $F$ , respectively. Given a bounded sequence  $(\alpha_n)$ , let  $T : E \rightarrow F$  be the bounded linear operator with  $T(\epsilon_n) = \alpha_n f_n$  for each  $n$ . Then  $T$  is compact if and only if  $\alpha_n \rightarrow 0$ .

Proof. Suppose  $T$  is compact. Let  $(P_n)$  be the sequence of the natural projections associated with  $(f_n)$ . Then  $(P_n \circ T)$  converges uniformly to  $T$  on the closed unit ball  $B_E$ , from which it follows that  $\alpha_n \rightarrow 0$ .

Conversely suppose that  $\alpha_n \rightarrow 0$ . We will show that  $T$  is weakly uniformly continuous on bounded subsets of  $E$ . Let  $B_r$  be the closed ball of  $E$  with the radius  $r$  and the center 0 and  $C$  be the positive number with  $|\alpha_n| \leq C$  for all  $n$ . Given  $\epsilon > 0$ ,  $x = \sum a_n \epsilon_n$  and  $y = \sum b_n \epsilon_n$  in  $B_r$ ,

$$\|T(x) - T(y)\| = \left\| \sum \alpha_n (a_n - b_n) f_n \right\| \quad (2.1)$$

$$\leq C \sum_{n=1}^{N-1} |a_n - b_n| + \left\| \sum_{n=N}^{\infty} \alpha_n (a_n - b_n) f_n \right\| \quad (2.2)$$

$$\leq C \sum_{n=1}^{N-1} |a_n - b_n| + 2K \left( \sup_{n \geq N} |\alpha_n| \right) \left\| \sum_{n=N}^{\infty} (a_n - b_n) f_n \right\|, \quad (2.3)$$

where  $K$  is the unconditional constant of  $(f_n)$ . Since  $(\epsilon_n)$  and  $(f_n)$  are equivalent, it is easy to see that

$$\left\| \sum_{n=N}^{\infty} (a_n - b_n) f_n \right\| \leq 2(1 + K)r\|T\|. \quad (2.4)$$

Let  $(f_n^*)$  be the sequence of coefficient functionals associated with  $(f_n)$ . Since  $\alpha_n \rightarrow 0$ , choosing sufficiently large  $N$ , we conclude that

$$\|T(x) - T(y)\| \leq \epsilon \quad (2.5)$$

if  $|f_1^*(x - y)|, \dots, |f_{N-1}^*(x - y)|$  are sufficiently small. Hence  $T$  is weakly uniformly continuous on bounded subset of  $E$ .

From the above proof it is easy to see that the Banach space  $c_0$  of null complex sequences is isomorphic with the Banach space of compact diagonal linear operators  $T : E \rightarrow F$ , where  $E$  and  $F$  are Banach spaces with equivalent unconditional bases. We would like to remark that if  $(\epsilon_n)$  and  $(f_n)$  are not equivalent, then given a bounded complex sequence  $(\alpha_n)$ , the map  $T(\epsilon_n) = \alpha_n f_n$  is not necessarily extended to a bounded linear operator from  $E$  into  $F$ . For example take  $E = \ell_2$ ,  $F = \ell_1$  and  $\alpha_n = 1$  for all  $n$  with respect to the canonical bases of them.

ACKNOWLEDGEMENT. Both authors would like to thank the referees for helpful suggestions and remarks.

#### REFERENCE

1. ARON, R.M. and PROLLA, J.B. Polynomial approximation of differentiable functions on Banach spaces, J. Reine Angew. Math. 313 (1980), 195-216.
2. CONWAY, J.B. A Course in Functional Analysis, Springer-Verlag, New York-Berlin-Heidelberg-Tokyo, 1985.
3. LINDENSTRAUSS, J. and TZAFIRI, L. Classical Banach Spaces I, Springer-Verlag, Berlin-Heidelberg-New York, 1977.

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)