

VECTOR FIELDS GENERATING MORE THAN ONE FLOW

SLAWOMIR DOROSIEWICZ* and KAZIMIERZ NAPIÓRKOWSKI**

* Institute of Econometry, Central School of Planning and Statistics, Al.
Niepodległości 162, 02-519 Warsaw, Poland.

** Department of Mathematical Methods in Physics, Warsaw University, Hoża 74,
00-682 Warsaw, Poland.

(Received February 18, 1993 and in revised form July 19, 1993)

ABSTRACT. Vector fields generating more than one flow on a manifold are constructed. For one-dimensional case a complete description of the set of flows is given. For dimensions larger than one a method of constructing vector fields with dense or open branched sets sets is given. Density of vector fields with nonempty sets of branched points is studied.

KEY WORDS AND PHRASES. Dynamical systems, uniqueness, Cauchy problem.

1991 AMS SUBJECT CLASSIFICATION CODE: 34A10.

1. INTRODUCTION.

There are many elementary examples of vector fields for which the corresponding differential equation has non-unique solution of the Cauchy problem. The examples, as a rule, do not define flows on the underlying manifold. The reason is that considered vector fields have singularity at isolated points. In the present paper we show, mainly on examples, how to construct vector fields which generate more than one flow.

The basic sections in this paper are Section 3, 4 and 5. Section 3 describes the one-dimensional case. It occurs that the set of singularities (i.e. zeros) of the vector field has to be of special type, namely, it must contain the support of a nonatomic measure satisfying some additional conditions. The set of all flows generated by a given vector field can be completely described by means of class of measures supported on the set of zeros of the vector field.

Two and more dimensional case is considered in section 4. Situation is much more complicated. The points in which the non-uniqueness of the flow takes place (called branched points) are no longer zeros of the vector field. Another phenomenon is that the set of branched points can be dense or open. We are not able to give a complete description as in Section 3, but some methods of constructing examples based on combining one dimensional examples with a constant vector field are given. Approximation methods developed in this section are use in Section 5 for a study of density properties of the set of vector fields generating more than one flow.

2. GENERAL PROPERTIES OF FLOWS AND QUASI-FLOWS.

DEFINITION 2.1

Let X be a Hausdorff space.

(i) A continuous map $\varphi: \mathbb{R} \times X \rightarrow X$, such that $\varphi(0, x) = x$ for $x \in X$, satisfying the group property: $\varphi(t+s, x) = \varphi(t, \varphi(s, x))$ and for all $t, s \in \mathbb{R}$ and $x \in X$ will be called a *flow* on X .

(11) If a map $\varphi: \mathbb{R} \times X \rightarrow X$ satisfies the group property and for all $x \in X$ the map $\varphi(\cdot, x): \mathbb{R} \rightarrow X$ is continuous, then φ will be called a *quasi-flow* on X .

DEFINITION 2.2

Let φ be a quasi-flow on a Hausdorff space X .

- (1) The set $\{\varphi(t, x) : t \in \mathbb{R}\}$ is called φ -orbit of x and is denoted by $\mathcal{O}_\varphi(x)$.
- (11) A point $x \in X$ is called φ -fixed point if φ -orbit of x is trivial i.e. $\mathcal{O}_\varphi(x) = \{x\}$. A set of φ -fixed points will be denoted by $S(\varphi)$.

EXAMPLE 2.3

For any given integer n , let φ be the map $\mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by $\varphi(t, x_1, \dots, x_n) = (x_1 + t, x_2, \dots, x_n)$. It is easy to see that φ is a flow on \mathbb{R}^n ; we denote it by i_n .

DEFINITION 2.4

Let U be an open subset of a Hausdorff space X , and let φ, ψ be flows on X . We say that φ and ψ are *equal on U* if the following condition is satisfied: for any $(t, x) \in \mathbb{R} \times U$ if $\varphi(t, x) \in U$ and $\psi(t, x) \in U$ then $\varphi(t, x) = \psi(t, x)$.

DEFINITION 2.5

Let X and Y be Hausdorff spaces, $U \subset X$ be a nonempty subset. Suppose that φ and ψ are the flows on X , Y respectively. We say that φ and ψ are *conjugate on U* if there exists a homeomorphism $f: X \rightarrow Y$, such that:

$$\psi(t, f(x)) = f(\varphi(t, x))$$

for all $x \in U$ and $t \in \mathbb{R}$ such that $\varphi(t, x) \in U$. If $U = M$, then we say that the flows φ, ψ are *conjugate*.

Elementary properties of flows on one dimensional connected manifolds are described in the following two propositions.

PROPOSITION 2.6

Let φ be a quasi-flow on \mathbb{R} . Then φ is a flow on \mathbb{R} , and for any point $x \in \mathbb{R}$ exactly one of the following conditions is satisfied:

- (a) x is a φ -fixed point,
- (b) the orbit $\mathcal{O}_\varphi(x)$ is an open interval containing the point x . Moreover, if a set \mathcal{O} is a nontrivial φ -orbit then the restriction $\varphi|_{\mathbb{R} \times \mathcal{O}}$ of φ to the set $\mathbb{R} \times \mathcal{O}$ is conjugate with the unit flow i_1 , i.e. there exists a homeomorphism $\sigma: \mathbb{R} \rightarrow \mathcal{O}$, which satisfies the equality: $\varphi(t, x) = \sigma(t + \sigma^{-1}(x))$ for all $x \in \mathcal{O}$ and $t \in \mathbb{R}$.

A similar result can be formulated for the flows on the one dimensional sphere:

PROPOSITION 2.7

Let φ be a quasi-flow on S^1 , then φ is a flow, and for any point $x \in S^1$ exactly one of the following conditions is satisfied:

- (a) x is φ -fixed point,
- (b) the orbit containing x is equal S^1 ,
- (c) the orbit $\mathcal{O}_\varphi(x)$ is diffeomorphic to \mathbb{R} , i.e. $\mathcal{O}_\varphi(x)$ is an open arc.

The proof is elementary and will be omitted.

The following definition introduces a notion very useful in further considerations: the time measure of a flow on \mathbb{R} . This notion is a special case of the

time measure of a quasi-flow on a Hausdorff space (see [2] p. 151).

DEFINITION 2.8

Let β denote the family of Borel sets on \mathbb{R} and $\bar{\mathbb{R}}$ denote $\mathbb{R} \cup \{\infty\}$, let φ be a flow on \mathbb{R} . The time measure of the flow φ is a Borel measure $\mu: \beta \rightarrow \bar{\mathbb{R}}$ defined as follows:

- (a) if x is a φ -fixed point then $\mu(\{x\}) = +\infty$,
- (b) if $x \in S(\varphi)$ then for any Borel subset $A \subset \varphi(x)$

$$\mu(A) = \mu_L(\varphi^{-1}(A, x)) = \mu_L(\{\tau \in \mathbb{R}^1 : \varphi(\tau, x) \in A\})$$

where μ_L is the Lebesgue measure on the real line \mathbb{R} ,

- (c) for any Borel set A :

$$\mu(A) = \sum_{\emptyset \in \Gamma} \mu(A \cap \emptyset) + \mu(A \cap S(\varphi)) ,$$

where Γ is the family of all nontrivial φ -orbits.

PROPOSITION 2.9

A necessary and sufficient condition for a Borel measure μ on \mathbb{R} to be the time measure of a flow on \mathbb{R} is that for each point $x \in \mathbb{R}$ exactly one of the following conditions is satisfied:

- (i) $\mu(\{x\}) = +\infty$,
- (ii) there exist γ, δ ; $-\infty < \gamma < \delta \leq +\infty$, such that $x \in]\gamma, \delta[$ and the restriction of μ to the interval $]\gamma, \delta[$ is a nonnegative Borel measure with the properties:
 1. If A is a nonempty open subset of $]\gamma, \delta[$ then $\mu(A) > 0$,
 2. The measure of any compact subset of $]\gamma, \delta[$ is finite and $\mu(]\gamma, c[) = \mu(]c, \delta[) = +\infty$ for each $c \in]\gamma, \delta[$.

The proof is easy and will be omitted.

3. VECTOR FIELDS AND FLOWS ON ONE-DIMENSIONAL MANIFOLDS.

DEFINITION 3.1

Let M be a n -dimensional smooth manifold, V be a vector field and φ be a flow on M . We say that V generates the flow φ , if for any point $x \in M$ $V(x)$ is the tangent vector to the curve $t \mapsto \varphi_t(x)$ at $t=0$. The set of all flows generated by the vector field V will be denoted by $F_1(V)$.

An analogous definition may be formulated for quasi-flows. The set of all quasi-flows generated by a vector field V will be denoted by $F_1^q(V)$.

DEFINITION 3.2

Let m be a cardinal number. We say that:

- (i) a vector field V on a differential manifold has F_m -property if the cardinality of the set $F_1(V)$ is equal m .
- (ii) a flow φ on differential manifold has F_m -property if it is generated by the vector field with F_m -property.

From standard theorems of the theory of ordinary differential equations follows that any smooth and bounded vector field on finite dimensional manifold generates exactly one flow, i.e. it has F_1 -property. The following example shows that

even on a one-dimensional manifold there exist vector fields which have $F\phi$ -property.

EXAMPLE 3.5

Let C denote the Cantor set in the interval $[0,1]$, and let $\gamma: \mathbb{R} \rightarrow \mathbb{R}$ be a nonnegative continuous function on \mathbb{R} with period 1 such that $\{x \in [0,1] : \gamma(x)=0\}=C$. It is easy to see that the map

$$\sigma: \mathbb{R} \rightarrow \mathbb{R}, \quad \sigma(t) = \int_0^t \gamma(s) ds$$

is a homeomorphism of \mathbb{R} onto \mathbb{R} . It is also obvious, that the velocity of the flow $\varphi = \sigma[1]$, i.e. $\varphi(t, x) = \sigma(t + \sigma^{-1}(x))$ for $t \in \mathbb{R}$ and $x \in \mathbb{R}$, is equal

$$V_{\varphi}(x) = \lim_{\varepsilon \rightarrow 0} \frac{\varphi(\varepsilon, x) - x}{\varepsilon} = \frac{d\varphi}{dt}(0, x) = \gamma(\sigma^{-1}(x))$$

for any point $x \in \mathbb{R}$. We shall show that the vector field V_{φ} generates uncountable many flows on \mathbb{R} . Let μ_{φ} denote the time measure of the flow φ . The set $Z(V_{\varphi})$ of critical points of V_{φ} is equal:

$$Z(V_{\varphi}) = \{x \in \mathbb{R} : \gamma(\sigma^{-1}(x)) = 0\} = \{x \in \mathbb{R} : \sigma^{-1}(x) \in \bigcup_{k \in \mathbb{Z}} k + C\} = \sigma(\bigcup_{k \in \mathbb{Z}} k + C)$$

Let ν be a measure concentrated on the set $\sigma^{-1}(\bigcup_{k \in \mathbb{Z}} k + C)$ of critical points of V_{φ} .

One can take for example the measure generated by the Cantor function as follows. First, let us define a function $f: C \rightarrow [0,1]$ in the following way: if $r = \sum_{i=1}^{+\infty} r_i \cdot 3^{-i}$, where $r_i \in \{0,2\}$, then $f(r) = \sum_{i=1}^{+\infty} r_i \cdot 2^{-i}$. It can be shown that f is a continuous surjection, moreover, it has a unique extension to a continuous function F on $[0,1]$. The derivative of this function vanishes on $[0,1] - C$, and does not exist on C . Let F^* denote the following function: $F^*(x+k) = F(x) + k$ for all $x \in [0,1]$ and $k \in \mathbb{Z}$. It is obvious that the measure ν defined for any compact, nonempty interval $[\alpha, \beta]$ as:

$$\nu([\alpha, \beta]) = F^*(\sigma(\beta)) - F^*(\sigma(\alpha))$$

is a continuous nonnegative Borel measure concentrated on the set of critical points of the vector field V_{φ} . Let us return to general considerations of this example. It is easy to see that for any positive number τ , the measure $\mu = \mu_{\varphi} + \tau \cdot \nu$ satisfies the conditions (i), (ii) of proposition 2.9 and therefore is the time measure of a flow $\bar{\varphi}$ on \mathbb{R} . Now it is sufficient to show that this flow (i.e. the flow with the time measure μ) is generated by the same vector field V_{φ} i.e. the flows φ and $\bar{\varphi}$ have the same velocity V_{φ} .

- 1) The set $\mathbb{R} - Z(V_{\varphi})$ is open. Consider any of its points x and take a positive number $r > 0$ such that $[x-2r, x+2r] \subset \mathbb{R} - Z(V_{\varphi})$. It means that $\mu_{\varphi}([x, x+\varepsilon] \cup [x+\varepsilon, x])$ is equal $\mu_{\varphi}([x, x+\varepsilon] \cup [x+\varepsilon, x])$ for $\varepsilon < 2r$, and this fact implies the equality $\bar{\varphi}(t, x) = \varphi(t, x)$ in a neighbourhood of the point $t=0$. It means that the flows φ and $\bar{\varphi}$ have the same velocities at the point x .
- 2) If $x \in Z(V_{\varphi})$ then $V_{\varphi}(x) = 0$, and $\lim_{\varepsilon \rightarrow 0} \frac{\mu_{\varphi}([x, x+\varepsilon] \cup [x+\varepsilon, x])}{\varepsilon} = 0$

It follows from definition of the measure μ that

$$\lim_{\epsilon \rightarrow 0} \frac{(\mu + \tau \cdot v)([x, x+\epsilon] \cup [x-\epsilon, x])}{|\epsilon|} = +\infty$$

which is equivalent that the velocity of the flow $\bar{\varphi}$ at the point x vanishes. Therefore the vector field $V_{\bar{\varphi}} = \gamma(\sigma^{-1})$ generates the flows $\bar{\varphi}$ and φ .

In Section 4 we shall use vector fields with compact sets of critical points and similar properties.

EXAMPLE 3.4

Let γ' be any any smooth nonnegative function $\gamma': \mathbb{R} \rightarrow \mathbb{R}$, such that $\gamma' = 1$ on $\mathbb{R} - [-1/2, 3/2]$ and $\gamma' = \gamma$ on $[0, 1]$, where γ denotes the function from the last example. It is clear that the vector field $V' = \gamma'(\sigma'^{-1})$, where $\sigma'(x) = \int_0^x \gamma'(s) ds$, has the Fc-property.

Example 3.3 suggests that there is a one to one correspondence between flows generated by a given vector field on \mathbb{R} and some measures concentrated on the set of critical points of this field.

In further considerations, let V be a continuous, bounded vector field on \mathbb{R} , and let $Z(V)$ denote the set of its critical points. First note that if x_0 is not a critical point of the vector field V on \mathbb{R} , then there exists a neighborhood U of x_0 such that the Cauchy problem

$$\frac{dy}{dt} = V(y), \quad y(0) = x_0 \quad (3.1)$$

has exactly one solution in U .

Indeed, if U is a connected component of $\mathbb{R} - Z(V)$ containing x_0 , then we can write (3.1) on U as: $t = t_0 + \int_{x_0}^y \frac{ds}{V(s)}$, $y \in U$, obtaining exactly one function $t = t(\cdot, x_0)$ and equivalently one function $y = y(\cdot, t_0)$, which is the solution of (3.1).

We shall consider now the set of flows generated by a given vector field V on \mathbb{R} . Let V be a continuous, bounded vector field on \mathbb{R} and let φ be a flow generated by this field with the time measure μ_{φ} and the set $S(\varphi)$ of φ -fixed points. For any closed set $\Omega \subseteq Z(V)$ let $\{P_{\alpha}\}_{\alpha \in Q}$ be the family of connected components of the set $\mathbb{R} - \Omega$.

For further considerations we introduce, for a given Ω and φ , the set $K_{\varphi}(\Omega)$, elements of which are all (not necessarily positive) Borel measures m on \mathbb{R} satisfying the following conditions:

1. m is concentrated on $Z(V)$,
2. $\mu_{\varphi} + m$ is nonnegative and if A is any open nonempty subset of \mathbb{R} then $(m + \mu_{\varphi})(A) > 0$,
3. if A is a compact subset of any P_{α} then $m(A) < +\infty$, and $(\mu_{\varphi} + m)(P_{\alpha}) = +\infty$,
4. for any $x \in Z(V)$ $\lim_{\epsilon \rightarrow 0} \frac{\epsilon}{(\mu_{\varphi} + m)([x, x+\epsilon] \cup [x-\epsilon, x])} = 0$,
5. if $x \in \Omega$ then $m(\{x\}) = +\infty$, for $y \in \Omega$ $\mu_{\varphi}(\{y\}) = 0$.

A topology in $K_{\varphi}(\Omega)$ may be introduced by the following

DEFINITION 3.5

We say that a sequence $m_n \in K_{\Phi}(\Omega)$ converges to a measure m if:

$$\lim_{n \rightarrow \infty} \sup_{x \in \Omega} (|m_n(B) - m(B)| : B \text{ is any compact subset of } P_{\alpha}) = 0$$

The set $F_1(V)$ has the standard uniform convergence topology.

THEOREM 3.6

Let V be a continuous, bounded vector field on \mathbb{R} . If the set $F_1(V)$ is not empty, then there exists a flow $\Phi \in F_1(V)$ such that the mapping

$$\lambda: \bigcup_{\Omega} K_{\Phi}(\Omega) \rightarrow F_1(V), \quad \lambda(m) = \psi_m$$

which assigns to a measure m the flow ψ_m , which has the time measure $\mu_{\Phi} + m$, where μ_{Φ} denotes the time measure of Φ , is a continuous bijection.

Proof. Since $F_1(V) \neq \emptyset$ we can choose $\varphi \in F_1(V)$. For a closed set $\Omega \subseteq \mathbb{R}$ we denote by $F_1(V, \Omega)$ the subset of $F_1(V)$ consisting of all flows having Ω as the set of fixed points. It is clear that for $\Omega = S(\varphi)$ the mapping λ gives a continuous bijection between $K_{\varphi}(\Omega)$ and $F_1(V, \Omega)$. To complete the proof it suffices to show that there exists a flow $\Phi \in F_1(V)$ with a minimal set of fixed points i.e. $S(\Phi) \subseteq S(\varphi)$ for any $\varphi \in F_1(V)$. We remark first that if $\varphi, \psi \in F_1(V)$ and $\mu_{\varphi}, \mu_{\psi}$ are their time measures then $\mu = \min(\mu_{\varphi}, \mu_{\psi})$ is the time measure of a flow belonging to $F_1(V)$. We denote that flow by $\min(\varphi, \psi)$. Since $x \in S(\varphi)$ if and only if $\mu_{\varphi}(\{x\}) = \alpha$ we have $S(\min(\varphi, \psi)) = S(\varphi) \cap S(\psi)$. One can check that a decreasing net of time measures of flows in $F_1(V)$ converges to the time measure of a flow in $F_1(V)$. Using Kuratowski-Zorn lemma we get the existence of a flow $\Phi \in F_1(V)$ with the minimal set of fixed points. Q.E.D.

An analogous result holds when \mathbb{R} is replaced by \mathbb{S}^1 . For manifolds of larger dimension than 1 one can conclude the following

COROLLARY 3.7

Suppose that φ and ψ are different flows on a manifold M , and each ψ -orbit is contained in a φ -orbit. If φ and ψ are generated by the same vector field V then V has at least one critical point.

4. VECTOR FIELDS AND FLOWS ON HIGHER DIMENSIONAL MANIFOLDS.

This part starts from the following definition describing singular points of a given vector field V on finite-dimensional manifold.

DEFINITION 4.1

Let V be a vector field on a differential manifold M .

- (a) A point $x \in M$ will be called a V -strong branched point if there exist quasi-flows φ and ψ generated by V and a positive number $\epsilon > 0$ such that the set $R(\varphi, \psi) = \{t \in \mathbb{R} : \varphi(t, x) \neq \psi(t, x)\}$ contains at least one of the intervals $(0, \epsilon)$, $(-\epsilon, 0)$. The set of V -strong branched points will be denoted by $\mathcal{R}(V)$.
- (b) A point $x \in M$ will be called a V -weak branched point if there exist two solutions $y_i(\cdot, 0, x)$, $i=1, 2$ of Cauchy Problem:

$$y' = V(y), \quad y(0) = x, \quad (4.1)$$

such that $t=0$ is a cluster point of the set $\{t \in \mathbb{R} : v_1(t, 0, x) \neq v_2(t, 0, x)\}$. The set of weak V -branched points will be denoted by $\mathcal{R}_w(V)$.

It is obvious that each strong branched point of a given vector field V is a V -weak branched point, and if x is a V -weak branched point, then the Cauchy problem (4.1) has uncountable many solutions (see for example [5], theorem 4.1).

DEFINITION 4.2

If, for a given subset A of a manifold M , there exists a vector field V on M such that $\mathcal{R}(V)=A$ ($\mathcal{R}_w(V)=A$), then A will be called a *strong branched set* (resp. *weak branched set*).

The following fact follows easily from Theorem 4.6:

PROPOSITION 4.3

Suppose that V is a continuous bounded vector field on \mathbb{R} or S^1 , then $\mathcal{R}(V)$ is a nowhere-dense subset of this manifold, contained in the set of critical points of the field V .

It follows from the above proposition that an open subset of \mathbb{R} or S^1 can never be a weak branched set. Now we shall show that there exist strong branched sets dense in higher dimensional manifolds, especially in \mathbb{R}^2 . The construction is a continuation of M. Lavrentiev's and P. Hartman's ideas (see [6] and [7]), who have constructed examples of ordinary differential equations with locally non-unique solution of Cauchy Problem in any point of an open subset of \mathbb{R}^2 . It is equivalent to proving that \mathbb{R}^2 is a weak branched set i.e. there exists a continuous vector field V on \mathbb{R}^2 such that $\mathcal{R}_w(V)=\mathbb{R}^2$. We are going to show more, namely, that \mathbb{R}^2 is a strong branched set. First in the example 4.4 we construct a vector field V such that $\mathcal{R}(V)$ is dense in \mathbb{R}^2 .

EXAMPLE 4.4

We construct an example of a continuous vector field V on \mathbb{R}^2 , with a dense set of strong branched points. The construction is divided into two steps. In first of them we define an operation called MOD-operation and analyse its properties. The second step included a proper description of the construction.

STEP 1

Let v be a continuous vector field on \mathbb{R}^2 and $\mathcal{R}(v)$ be the set of its strong branched points. Suppose that for a connected open subset U of the set $\mathbb{R}^2 - \mathcal{R}(v)$ there exists a smooth local coordinate system (λ, U) in which v has the coordinates $\langle 1, 0 \rangle$.

Let Ω denote a nonempty rectangle $[a, b] \times [c, d] \subset \lambda(U)$. We choose real functions $\alpha(\cdot)$ and $h(\cdot)$ such that $\alpha: \mathbb{R} \rightarrow \mathbb{R}$ is continuous in $[a, b]$, and $\alpha \equiv 0$ on $\mathbb{R} - [a, b]$, $h: \mathbb{R} \rightarrow [c, d]$ is a diffeomorphism such that $\frac{dh}{dx} \rightarrow 0$ when $x \rightarrow \pm\infty$. We assume, moreover, that

$$\int_a^b \alpha(s) ds = 0, \sup |\alpha| > 6\pi(b-a)^{-1} \text{ and } \sup \left| \frac{dh}{dx} \right| = \eta \quad (4.2)$$

where η is a positive real number.

Let u denote the vector field constructed in Example 3.3. We can define the following vector field on \mathbb{R}^2 : $f(x, y) = \langle 1, \bar{u}(y) \cdot \alpha(x) \rangle$, where:

$$\bar{u}(y) = \begin{cases} h'(y)u(h^{-1}(y)) & \text{for } y \in [c, d], \\ 0 & \text{for } y \in \mathbb{R} - [c, d]. \end{cases}$$

To assure that each orbit of any flow from $F^1(f)$ crosses the set of branched points of f , we assume additionally that there exist a', b' such that

$$\int_{a'}^{b'} \alpha(s) ds > \sigma^{-1}(1) \quad (4.3)$$

where σ is defined in Example 3.3.

DEFINITION 4.5

We say that a vector field v_1 is a *modification* of v_0 and write $v_1 = \text{MOD}(U, \mathcal{R})v_0$ if there exist $\alpha, h, \lambda, \mathcal{R}$ as above such that $v_1 = v_0$ outside U and $v_1 = f$ in U in local coordinates given by λ .

To analyze properties of v_1 note that f has F^1 -property: for any $\phi \in F^1(u)$, it generates the flow

$$\psi(t, (x, y)) = (x + t, \phi(\int_0^t \alpha(x+s) ds, y)). \quad (4.4)$$

Vector fields $\langle 1, 0 \rangle$ and f are different only inside the rectangle \mathcal{R} . It is easy to see that $\mathcal{R} = [a, b] \times \mathcal{R}(\bar{u})$ is the set of f -strong branched points. Condition (4.3) implies that the orbits of any flow generated by f , which pass across \mathcal{R} have nonempty intersection with the set \mathcal{R} . It is easy to see that flows generated by f which are of the form (4.4) are conjugate with the unit flow $\mathbf{1}_2$ by the following homeomorphism Λ :

$$\Lambda(x, y) = (x, \phi(\int_{-\alpha}^x \alpha(x+s) ds, y)), \text{ for } (x, y) \in \mathbb{R}^2.$$

DEFINITION 4.6

We say that a vector field V has α -property if for any point $p \in \overline{\mathcal{R}(v)}$ there exists a connected neighborhood $U(p)$ and a local map $\lambda: U(p) \rightarrow \lambda(U(p)) \subset \mathbb{R}^2$ such that V has the coordinates $\langle 1, 0 \rangle$ in the map λ .

Observe that if a vector field V has the α -property, p and $U(p)$ are as in the above definition, then $\text{MOD}(U(p), \mathcal{R})V$ has the α -property. The operation $\text{MOD}(U(p), \mathcal{R})$ depends on the choice of the local map λ . We can choose λ in such a way that $p \in \overline{\mathcal{R}(\text{MOD}(U(p), \mathcal{R})V)}$. In the following we shall always choose such a λ .

STEP 2

We choose a countable dense set $P = \{p_n : n \in \mathbb{N}\}$ in \mathbb{R}^2 and start from the vector field V_1 with coordinates $\langle 1, 0 \rangle$ which obviously has α -property. Let V_n denote the vector field obtained in n -th iteration. As the next iteration we take $V_{n+1} = \text{MOD}(U(p_n), \mathcal{R}_n)V_n$ if $p_n \in \overline{\mathcal{R}(V_n)}$ and $V_{n+1} = V_n$ otherwise.

LEMMA 4.7

Parameters \mathcal{R}_n of the MOD -operation can be chosen small enough, so that

- the sequence (V_n) converges uniformly on \mathbb{R}^2 (in the sense of the uniform convergence of coordinates in the canonical map in \mathbb{R}^2).
- each of V_n has the F^1 -property.

c) $\mathcal{R}(V_{n+1}) \supseteq \mathcal{R}(V_n)$.

Proof. The distance d_n between v_n and v_{n+1} , defined as $\sup\{\|v_n(x) - v_{n+1}(x)\| : x \in \mathbb{R}^2\}$, is proportional to η_n (with a constant depending on n), so it is possible to make the series $\sum d_n$ convergent. b) and c) follow immediately from the construction. Q.E.D.

If n is such that $v_{n+1} \neq v_n$ then for any flow $\Phi_n \in \mathcal{F}^1(V_n)$ we can construct, using (4.3), continuum of flows $\Phi_{n+1} \in \mathcal{F}^1(V_{n+1})$ which are equal to Φ_n outside $U(p_n)$ (in the sense of definition 4.4). The distance between Φ_n and Φ_{n+1} defined as $\sup\{\|\Phi_n(t, p) - \Phi_{n+1}(t, p)\| : t \in \mathbb{R}, p \in \mathbb{R}^2\}$ can be bounded by constant (depending on n) proportional to η_n .

LEMMA 4.8

Parameters η_n of the MOD-operation can be chosen small enough, so that any sequence (Φ_n) chosen as above converges uniformly to a flow on \mathbb{R}^2 .

The proof follows from the above remarks.

PROPOSITION 4.9

If η_n are chosen according to Lemmas 4.7 and 4.8 then the vector field $V = \lim V_n$ has a dense set of strong branched points. The limit of any sequence (Φ_n) as in Lemma 4.8 is a flow generated by V .

Proof. Lemma 4.8 guarantees that $\lim \Phi_n$ is a flow on \mathbb{R}^2 . From standard theorems about differential equations (see [5] Theorem 4.4) follows that $\lim \Phi_n$ is generated by V . It is easy to see from the construction, that using different flows from $\mathcal{F}^1(\bar{U})$ in any MOD($U(p_n), \eta_n$)-operation we obtain different limits $\lim \Phi_n$. It implies that V has Fc-property and any point of P is a V -strong branched point. Q.E.D.

It can be proved that if the parameters η_n are sufficiently small the set $\mathcal{R}(V)$ is dense but different from \mathbb{R}^2 . This completes Example 4.4.

REMARK 4.10

If the vector field u used in MOD-operation, is replaced by the vector field constructed in Example 3.4 we can obtain a vector field V^* such that $\mathcal{R}(V^*) = \mathbb{R}^2$, i.e. \mathbb{R}^2 is a strong branched set.

5. GENERAL PROPERTIES OF VECTOR FIELDS AND FLOWS WITH Fc-PROPERTY

In this part we investigate sets of vector fields and flows having Fc-property. Those sets will be regarded as subsets of the spaces $\mathcal{F}^0(M)$ and $\mathcal{F}^1(M)$ respectively, which consist of all continuous vector fields and flows on a finite-dimensional manifold M . In order to avoid problems with non-integrable vector fields and with definition of the uniform convergence of vector fields we assume that all considered manifolds are compact.

PROPOSITION 5.1

If M is an n -dimensional manifold and $n \geq 2$, then any continuous vector field on M can be uniformly approximated by vector fields with Fc-property.

Proof: If the given vector field is smooth and does not vanish one can apply MOD-operation with a sufficiently small parameter η . If the field vanishes in an

open set one can add to it any field with F_c -property, multiplied by a sufficiently small constant. A continuous vector field can be uniformly approximated by smooth fields. Q.E.D.

Let k be any cardinal number; the set of continuous, bounded vector fields on M with the F_k -property will be denoted by $G(M,k)$. The same way, if we consider the set $G_{int}(M)$ of vector fields with nonempty, open sets of strong branched points (for instance $G_{int}(\mathbb{R}^2) \neq \emptyset$ - see Remark 4.10), it can be similarly proved that $G_{int}(M)$ is dense in $F^\circ(M)$. The sets $G(M,0)$ and $G(M,1)$ are also dense in $F^\circ(M)$, it can be shown much more: $G(M,1)$ is the set of second Baire category (see [3] pp.119-121).

The space $F^\circ(M)$ may be in a natural way represented as a sum of disjoint sets $G(M,k)$, for $k \geq 0$. Unfortunately, for many dimensional ($\dim M \geq 2$) manifolds authors do not know anything about the sets $G(M,k)$ for $k \geq 2$ and $k \leq \aleph_0$; are these sets nonempty?

It was shown that on the n -dimensional manifolds, $n \geq 2$, there exist vector fields having open sets of strong branched points. For an arbitrary open set, approximating it by sums of rectangles, we get

PROPOSITION 5.2

Every open subset $\bigcup \mathbb{R}^n$, where $n > 1$, is a branched set.

It is easy to see that the stronger result occurs: every open subset of the n -dimensional manifold, where $n > 1$, is a branched set.

DEFINITION 5.3

For a compact manifold M and given cardinal number k we define the following sets:

$$A_q(M,k) = \bigcup_{f \in G(M,k)} \{f\} \times F_{\leq q}(f)$$

$$B_q(M) = \bigcup_{f \in F^\circ(M)} \{f\} \times F_{\leq q}(f)$$

Since M is metrizable, then using the metric p on M we define a topology of above sets by the following metric:

$$d((f_1, \varphi_1), (f_2, \varphi_2)) = \|f_1 - f_2\|_\infty + p_\infty(\varphi_1, \varphi_2)$$

where $\|\cdot\|_\infty$ is the norm of the uniform convergence in $F^\circ(M)$, the set of all continuous vector fields on M , and p_∞ denotes the norm:

$$p_\infty(\varphi_1, \varphi_2) = \sup_{t \in \mathbb{R}} \sup_{x \in M} p(\varphi_1(t, x), \varphi_2(t, x)).$$

In what follows the topology defined by d in those sets will be called the C° -topology.

Before formulating results of this section, we give a short description of MOD1-operation which assigns to a quasi-flow ψ a quasi-flow φ having the F_c -property such that the $p_\infty(\varphi, \psi)$ can be arbitrarily small. Let V be a continuous vector field on a manifold M with the set $Z(V)$ of critical points differ-

rent from M . Suppose that $F_{1_q}(V)$ is not empty and let ψ be a quasi-flow generated by V . It is obvious that for any $x \in Z(V)$ there exists a map (λ, U) of a neighbourhood U of x such that ψ is conjugate to the unit flow 1_n on $U \cap \psi(x)$ (in the sense of definition 2.5). Without loss of generality we can assume that (U) contains the cube $[0,1]^n$. Let α be a nonnegative continuous function on \mathbb{R}^n such that

- (a) $\alpha(\cdot) - 1$ vanishes beyond a neighborhood of the Cantor set C on interval $[0,1]$,
- (b) $\int_0^1 (\alpha(s) - 1) ds < \delta$, where δ denotes a positive number,
- (c) for any $x \in C$, $\alpha(x) = 1$.

Let ψ' be any flow which in the map (λ, U) has the following coordinates:

$$\psi(t, x) = (pr_1(x) + \int_{a_1}^{b_1} \alpha(s) ds, pr_2(x), \dots, pr_n(x)),$$

where $x \in \mathbb{R}^n$. Let φ denote the quasi-flow which is equal ψ on $M \cup \psi(x)$, and ψ' on $U \cap \psi(x)$.

DEFINITION 5.4

We say that a map $\varphi_1 : \mathbb{R} \times M \rightarrow M$ is a modification of quasi-flow ψ and write $\varphi_1 = \text{MOD1}(U, \delta)\psi$ if there exist $\alpha, \lambda, \varepsilon, U, \psi, \psi'$ as above such that $\varphi_1 = \psi$ outside $U \cap \psi(x)$ and $\varphi_1 = \psi'$ in $U \cap \psi(x)$.

It is easy to see that the map: φ is a quasi-flow with a F_c -property, sufficiently close to ψ : $\rho_0(\varphi, \psi) \leq \varepsilon$, where $\varepsilon = S\delta$ and S denotes a constant depending on λ, α and U .

THEOREM 5.5

Let M be a compact smooth manifold with Euler characteristic $\chi(M)$ different from zero. Suppose that V is continuous vector field on M and $\varphi_n \in F_{1_q}(V)$. Then there exists a sequence of vector fields V_n having the F_c -property and quasi-flows $\varphi_n \in F_{1_q}(V_n)$ which uniformly converge to V and φ respectively; i.e. the set $A_q(M, c)$ is dense in $B_q(M)$ in C^0 -topology.

Proof. Let V be a continuous vector field on M with nonempty set $F_{1_q}(V)$. If V vanishes in an open subset of M theorem is obvious. In the other case let us consider an integer number n and any quasi-flow $\varphi \in F_{1_q}(V)$. Since $\chi(M) \neq 0$, it follows from Poincaré-Hopf theorem (see [8] p.69), that V has nonempty set $Z(V)$ of critical points. Let us choose a point $x \in Z(V)$ and open set U such that $\varphi(x) \cap U$ is disjoint from $Z(V)$.

Performing the MOD1 -operation we obtain quasi-flow $\psi = \text{MOD1}(U, 1/n)\varphi$ which satisfies the condition

$$\|V' - V\|_0 + \rho_0(\psi, \varphi) \leq n^{-1},$$

where V' denotes the vector field which generates the quasi-flow ψ . Since n can be chosen arbitrarily large, the proof is complete. Q.E.D.

Since Euler characteristic $\chi(S^k)$ of k -dimensional sphere is equal $1 + (-1)^k$, it follows immediately from theorem 5.5 that the set $A_q(S^{2n}, c)$ is dense in $B_q(S^{2n})$.

in C^0 -topology.

Theorem 5.5 does not, in general, hold for manifolds with Euler characteristic zero. For example, since any vector field on S^1 with Fc -property has uncountable many of critical points (see Section 3); this fact implies that $A_q(S^1, c)$ is not dense in $B_q(S^1)$. There exist also higher dimensional manifolds for which Theorem 5.5 does not hold. Let us consider $T^2 = S^1 \times S^1 = \{\exp(2\pi i \alpha) : \alpha \in \mathbb{R}\} \times \{\exp(2\pi i \beta) : \beta \in \mathbb{R}\}$, and the flow Φ having the form $\Phi(t, (\alpha, \beta)) = (\alpha + bt + at \cos \beta, \beta)$, where a and b are positive numbers. Let V_Φ denote the vector field which generates Φ . It can be checked that the pair (V_Φ, Φ) can not be uniformly approximated by the vector fields and flows from $A_q(M, c)$. It appears, however, that V_Φ and Φ separately can be uniformly approximated by vector fields and flows with Fc -property. Note that this fact is more general:

PROPOSITION 5.6

If M is a compact smooth manifold, then the set $\bigcup_{V \in F^0(M)} F^0_q(V)$ is a dense subset of $\bigcup_{V \in F^0(M)} F^0_q(V)$ in the C^0 topology, i.e. any quasi-flow on M can be uniformly approximated by quasi-flows with Fc -property.

The proof is analogous to the proof of Theorem 5.5 and will be omitted.

We can easily construct an analogous example of a vector field and a flow on Klein bottle K^2 . Since for any two-dimensional compact manifold apart from S^2 and K^2 Euler characteristic is negative, we can formulate:

COROLLARY 5.7

Suppose that M is a two-dimensional compact manifold. Then the following conditions are equivalent:

1. Euler characteristic $\chi(M)$ is different from zero,
2. The set $A_q(M, c)$ is dense in $B_q(M)$ in C^0 topology,
3. $M \neq S^2$ and $M \neq K^2$.

REFERENCES

1. BATTY, C.J.K. Delays to flows on the real line. Preprint, 1979, unpub
2. BECK, A. Continuous Flows in the Plane, Springer-Verlag, Berlin, Heidelberg, New York, 1974.
3. CHOQUET, G. Lectures on Analysis, W.A. Benjamin, Mathematical Lecture Series, 24.
4. ENGELKING, R., SIEKLICKI, K. Geometry and topology, PWN, Warsaw, 1981
5. HARTMAN, P. Ordinary Differential Equations, John Wiley and Sons, N.Y., London, Sydney, 1964.
6. HARTMAN, P. A differential equation with nonunique solutions, Amer. Ma Monthly, 70 (1963), pp. 255-259.
7. LAVRENTIEV, M. Sur une équation différentielle du premier ordre, Mat Zeit. 23 (1925), pp. 197-209.
8. SZLENK, W. Smooth dynamical systems, PWN Warsaw, 1982.

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk