

A STABILITY RESULT FOR PARAMETER IDENTIFICATION PROBLEMS IN NONLINEAR PARABOLIC PROBLEMS

STANISLAW MIGÓRSKI

Institute for Information Sciences
Jagellonian University
ul. Nawojki 11, 30072 Cracow
Poland

(Received March 15, 1993)

ABSTRACT. A sequence of identification problems of coefficients in the parabolic equation with nonlinear boundary conditions is considered. The parameter (index of an element of the sequence) appears in the cost functionals as well as boundary data. It is proved that the optimal solutions exist and that under some continuous convergence of the cost functionals and the convergence of the data, the sets of optimal solutions converge in some sense to the set of optimal solutions of the limit problem.

KEY WORDS AND PHRASES. Inverse problem, system identification, stability of solutions, parabolic differential equation, nonlinear boundary condition.

1992 AMS SUBJECT CLASSIFICATION. 35R30, 35K60.

1. INTRODUCTION.

In the recent years there has been an increasing interest in the parameter identification (or inverse) problems involving differential equation constraints. Such problems arise in particular in the coefficient estimation for partial differential equations (for example in [2-3], [14], [17-18]) as well as in the theory of structural optimization. The identification problems consist in determining of unknown parameters (coefficients) from known observations of the modelled processes.

In this paper we investigate a class of identification problems for the second order nonlinear parabolic system:

$$u' = -\mathcal{A}(t)u \quad \text{in } \Omega \times (0, T), \quad (1.1)$$

$$\frac{\partial u}{\partial \nu_{\mathcal{A}(t)}} + \beta(u) \ni g \quad \text{in } \Gamma \times (0, T), \quad (1.2)$$

$$u(0) = \varphi \quad \text{in } \Omega, \quad (1.3)$$

where $\Omega \subset \mathbf{R}^n$ with boundary Γ , $0 < T < +\infty$, β is a maximal monotone graph in $\mathbf{R} \times \mathbf{R}$, the operator $\mathcal{A}(t)$ has the form

$$\mathcal{A}(t) = -\frac{\partial}{\partial x_i} (a_{ij}(x, t) \frac{\partial}{\partial x_j})$$

and

$$\frac{\partial u}{\partial \nu_{\mathcal{A}(t)}} = a_{ij}(x, t) \frac{\partial u}{\partial x_i} \nu_j$$

is the conormal derivative associated to $\mathcal{A}(t)$. Above, ν is the unit outward normal vector to Γ .

Given the set of admissible parameters \mathcal{M} and the cost functional \mathcal{J} defined on the space

\mathcal{W} (see Notation) of the solutions to (1.1) - (1.3), we are interested in the following parameter identification problem:

$$(\mathcal{P}) \quad \left\{ \begin{array}{l} \text{find the parameters } a^* = \{a_{ij}^*(x, t)\} \text{ in } \mathcal{M} \text{ so that} \\ \mathcal{J}(u(a^*, g, \varphi)) \leq \mathcal{J}(u(a, g, \varphi)) \text{ for all } a \in \mathcal{M}, \end{array} \right.$$

where $u(a, g, \varphi)$ denotes the weak solution in \mathcal{W} corresponding to the data a , g and φ . Here and in what follows, we suppose that the graph β is fixed.

Our aim is to prove the existence of optimal solutions (i.e. an element which realizes the minimum) to (\mathcal{P}) and to show the stability of optimal solutions under perturbations of the data g and φ as well as of the cost functional \mathcal{J} . As indicated in [1], [4-5] and [10-11] the stability of this kind plays an important role in applications.

It should be noticed that a compactness of admissible subset of parameters is the crucial assumption in the identification problems (compare e.g. [5]). Since the cost functional is not convex in general, the uniqueness of optimal solutions is not guaranteed. Therefore the stability is understood in the sense of continuity of multivalued mapping.

We note that the widely known approach to the parameter identification problems used also in numerical methods (see [4-5], [14]) is the output least squares formulation ([10-11], [13]). In this approach the cost functional to be minimized has the form

$$\|\mathcal{C}u(a, g, \varphi) - z_d\|_{\mathcal{Z}}^2, \quad (1.4)$$

where $\mathcal{C}: \mathcal{W} \rightarrow \mathcal{Z}$ is an observation operator defined on the space of solutions, z_d is the desired element (target) in the space of observations \mathcal{Z} . Such cases are also included in the frame of the paper. Next, it should be underlined that the identification of coefficients in partial differential equations is, in general, an unstable problem ([16-17], [13]). This is due to the theory of homogenization ([8], [18]) which shows that operators with highly oscillatory coefficients can be "replaced" by very different ones and still giving the same response.

Finally, we point out that the problems of the type (1.1) - (1.3) occur in many mathematical models of phenomena studied in physics. For example, equation (1.1) describes the change of pressure during the flow of viscous fluids in porous media or it governs the heat distribution in a body occupying the volume Ω . It is natural to consider such problems not only with the classical (Dirichlet and/or Neumann) boundary conditions, but also with the more general ones. The boundary condition (1.2) includes some particular cases e.g. the Signorini condition, the Stefan-Boltzmann heat radiation law, the Newton's law, the natural convection, the Michaelis-Menten law. For these and other important examples of the condition (1.2) which appears in mechanics, biology and chemistry, we refer to [12], [7], [9] and the bibliography in them. We also recall that the evolution variational inequalities can be formulated in the form (1.1) - (1.3) (see [6], [9], [12], [15]). For the general identification theory presented in an abstract manner we refer to [13] and [1].

The remainder of this note is divided in three parts. In Section 2 we give a result on the continuous dependence of solution to boundary value problem (1.1) - (1.3) on the data. With this background, in Section 3, we show that the problem (\mathcal{P}) has a solution. The last section is devoted to the stability of optimal solutions with respect to variations in the given data and the cost functional.

Notation. Let Ω be a bounded open subset of \mathbf{R}^n with Lipschitz continuous boundary Γ . For a fixed real interval $[0, T]$, we introduce $Q = \Omega \times (0, T)$, $\Sigma = \Gamma \times (0, T)$. Putting $V = H^1(\Omega)$, $H = L^2(\Omega)$, we denote by $\|\cdot\|$, $|\cdot|$, the norms in V and H , respectively. By V' we denote the dual to V . Following e.g. [15], we define the Banach spaces: $\mathcal{V} = L^2(0, T; V)$, $\mathcal{H} = L^2(0, T; H)$, $\mathcal{V}' = L^2(0, T; V')$ (the spaces of the square summable functions defined on $(0, T)$ with the values,

resp., in V , H , V') and $\mathcal{W} = \{v \in \mathcal{V} : v' \in \mathcal{V}'\}$. Here and subsequently the partial derivative with respect to t , is understood in the distributional sense and dv/dt will be denoted by v' . The pairing of V and V' and also the inner product on H is denoted by $\langle \cdot, \cdot \rangle$. The symbol $\langle \cdot, \cdot \rangle_\Gamma$ stands for the inner product on $L^2(\Gamma)$. Given closed sets A and B in a Banach space X , we define the distance function by $d(x, A) = \inf\{\|x - a\|_X : a \in A\}$ and the separation of a set A from a set B by

$$h^*(A, B) = \sup\{d(a, B) : a \in A\}. \quad (1.5)$$

Throughout this paper a summation convention over repeated subscripts is adopted.

2. CONTINUOUS DEPENDENCE ON THE DATA.

The goal of this section is to study the question of continuous dependence of solutions to (1.1) - (1.3) on the data a_{ij} , g and φ . First we give the existence result on the solutions to this problem. To this end, we adopt the following

DEFINITION 2.1. (see [7]) A function $u \in \mathcal{W}$ is a weak solution to (1.1) - (1.3) if and only if there exists a function $w \in L^2(\Sigma)$ such that

$$w(\sigma, t) \in \beta(u(\sigma, t)) \quad \text{a.e. on } \Sigma, \quad (2.1)$$

and

$$\langle u'(t), v \rangle + a(t; u(t), v) + \langle w(t), v \rangle_\Gamma = \langle g(t), v \rangle_\Gamma \quad \forall v \in V, \text{a.e. on } (0, T),$$

$$u(0) = \varphi,$$

where we have set

$$a(t; z, v) = \int_{\Omega} a_{ij}(x, t) \frac{\partial z}{\partial x_i} \frac{\partial v}{\partial x_j} dx, \quad \forall z, v \in V, \quad \text{a.e. } t \in (0, T). \quad (2.2)$$

We need the following hypotheses on the data of the problem (1.1) - (1.3):

(H_1) the coefficients $\{a_{ij}\}$, $i, j = 1, \dots, n$ are functions from $C(\overline{Q})$ such that

$$\alpha \xi_i \xi_i \leq a_{ij}(x, t) \xi_i \xi_j, \quad \forall \xi \in \mathbf{R}^n \quad (2.3)$$

a.e. in Q , for some constant $\alpha > 0$,

(H_2) β is a maximal (multivalued) monotone graph in $\mathbf{R} \times \mathbf{R}$ which satisfies the condition $0 \in \beta(0)$,

(H_3) $g \in L^2(\Sigma)$, $\varphi \in H$.

It is well known (see e.g. [6]) that β is a subdifferential of a proper, convex, l.s.c. function $j: \mathbf{R} \rightarrow \mathbf{R} \cup \{+\infty\}$, i.e. $\beta = \partial j$.

PROPOSITION 2.1. In addition to the hypotheses $(H_1) - (H_3)$ we assume that $j(\varphi) \in L^1(\Omega)$. Then the problem (1.1) - (1.3) has a unique weak solution $u \in \mathcal{W}$. Moreover, the following estimate holds:

$$\|u\|_{\mathcal{W}} + \|w\|_{L^2(\Sigma)} \leq c(1 + \|g\|_{L^2(\Sigma)} + |\varphi|), \quad (2.4)$$

where $w \in L^2(\Sigma)$ is a selection of β which appears in (2.1) and $c \in \mathbf{R}$ is independent of g and φ .

The proof of this result can be found in [7], Proposition 1, where the overall hypotheses on the coefficients a_{ij} were more restrictive. A careful look in that proof can convince the reader that it is still true for the case of coefficients satisfying (H_1) .

Let $\{a_{ij}^k\}$, $k \in \mathbf{N}$, be a sequence in $C(\overline{Q})$ satisfying (2.3) uniformly with respect to k and let $\{(g_k, \varphi_k)\}$ be a sequence in $L^2(\Sigma) \times H$. Denoting by $u_k = u(a_{ij}^k, g_k, \varphi_k)$ the solution to (1.1) - (1.3) corresponding to $\{a_{ij}^k\}$, $\{(g_k, \varphi_k)\}$, we have

LEMMA 2.1. Under the above notations, let us assume that β satisfies (H_2) and $j(\varphi_k) \in L^1(\Omega)$. If

$$a_{ij}^k \rightarrow a_{ij} \quad \text{in } C(\overline{Q}), \quad \forall i, j,$$

$$(g_k, \varphi_k) \rightarrow (g, \varphi) \quad \text{in} \quad L^2(\Sigma) \times H,$$

as $k \rightarrow +\infty$, then

$$u_k \rightarrow u \quad \text{in} \quad \mathcal{V} \cap C(0, T; H) \text{ and weakly in } \mathcal{W}, \quad (2.5)$$

as $k \rightarrow +\infty$, where $u = u(a_{ij}, g, \varphi)$ is a unique solution (still in the sense of Definition 2.1) to (1.1) - (1.3) corresponding to a_{ij} , g and φ .

PROOF. Note that by lower semicontinuity of j , it follows that $j(\varphi) \in L^1(\Omega)$. Therefore, from Proposition 2.1, we know that there exists a unique solution to (1.1) - (1.3) corresponding to a_{ij} , g and φ . Subtracting two equations which are satisfied by u_k and u , we obtain

$$\begin{aligned} & < u'_k(s) - u'(s), v > + a_k(s; u_k(s), v) - a(s; u(s), v) + \\ & + < w_k(s) - w(s), v >_\Gamma = < g_k(s) - g(s), v >_\Gamma \end{aligned}$$

for all $v \in V$, a.e. $s \in (0, T)$, where $w, w_k \in L^2(\Sigma)$ are such that

$$w(\sigma, s) \in \beta(u(\sigma, s)), \quad w_k(\sigma, s) \in \beta(u_k(\sigma, s)) \quad \text{a.e. on } \Sigma$$

and $a(\cdot, \cdot, \cdot), a_k(\cdot, \cdot, \cdot)$ are of the form (2.2) with the coefficients a_{ij}, a_{ij}^k , respectively.

Taking $u_k(s) - u(s)$ as the function v and integrating both sides we get:

$$\begin{aligned} & \frac{1}{2} |u_k(t) - u(t)|^2 + \int_0^t a_k(s; u_k(s) - u(s), u_k(s) - u(s)) ds + \\ & + \int_0^t < w_k(s) - w(s), u_k(s) - u(s) >_\Gamma ds = \frac{1}{2} |\varphi_k - \varphi|^2 + \int_0^t (a(s; u(s), u_k(s) - u(s)) - \\ & - a_k(s; u(s), u_k(s) - u(s))) ds + \int_0^t < g_k(s) - g(s), u_k(s) - u(s) >_\Gamma ds. \end{aligned}$$

Hence, from (H_1) and (H_2) we obtain:

$$\begin{aligned} & |u_k(t) - u(t)|^2 + 2\alpha \int_0^t \|u_k(s) - u(s)\|^2 ds \leq |\varphi_k - \varphi|^2 + \\ & + \int_0^t \sum_{j=1}^n \left| \sum_{i=1}^n (a_{ij}^k - a_{ij}) \frac{\partial u(s)}{\partial x_i} \right| \left| \frac{\partial u_k(s)}{\partial x_j} - \frac{\partial u(s)}{\partial x_j} \right| ds + \\ & + 2 \int_0^t |g_k(s) - g(s)|_{L^2(\Gamma)} |u_k(s) - u(s)|_{L^2(\Gamma)} ds. \end{aligned}$$

Using the continuity of the trace operator from V to $L^2(\Gamma)$ and the inequality $2ab \leq \frac{2}{\alpha}a^2 + \frac{\alpha}{2}b^2$, we have

$$\begin{aligned} & |u_k(t) - u(t)|^2 + \alpha \int_0^t \|u_k(s) - u(s)\|^2 ds \leq |\varphi_k - \varphi|^2 + \\ & + c_1 \|u\|_{\mathcal{V}}^2 \sum_{i,j=1}^n \|a_{ij}^k - a_{ij}\|_{C(\bar{Q})} + c_2 \int_0^t |g_k(s) - g(s)|_{L^2(\Gamma)} ds, \end{aligned}$$

where $c_i, i = 1, 2$ are positive constants independent of k . From this, (2.4) and from the hypotheses we deduce that

$$u_k \rightarrow u \quad \text{in} \quad \mathcal{V} \cap C(0, T; H), \quad \text{as} \quad k \rightarrow +\infty.$$

On the other hand, owing to the estimate (2.4), which is uniform with respect to k , we obtain

$$u_k \rightarrow u \quad \text{weakly in } \mathcal{W}, \quad \text{as} \quad k \rightarrow +\infty.$$

The uniqueness of solutions of problem (1.1) - (1.3) implies that the whole sequence $\{u_k\}$ converges to u in the sense of (2.5). This completes the proof of the lemma.

3. EXISTENCE RESULT IN IDENTIFICATION.

Using the result of Lemma 2.1, we can prove the existence of solution of the identification problem (\mathcal{P}) . This is given in the following

THEOREM 3.1. Let the assumptions of Proposition 2.1 hold. We suppose that the set of admissible parameters satisfies

(H_4) the set \mathcal{M} is a compact subset of $(C(\overline{Q}))^{n^2}$ such that $\alpha \xi_i \xi_i \leq a_{ij}(x, t) \xi_i \xi_j$ for each $\xi \in \mathbf{R}^n$ and for all $a = \{a_{ij}(x, t)\}$ in \mathcal{M} .

Let the cost functional $\mathcal{J}: \mathcal{W} \rightarrow \mathbf{R}$ be weakly sequentially lower semicontinuous on \mathcal{W} . Then the problem (\mathcal{P}) has a solution.

PROOF. We apply the direct method of calculus of variations (see e.g. [1]). Let g, φ satisfy (H_3) and let $\{a_k\}$ be a minimizing sequence from \mathcal{M} such that

$$\lim_{k \rightarrow \infty} \mathcal{J}(u(a_k, g, \varphi)) = \inf \{\mathcal{J}(u(a, g, \varphi)) : a \in \mathcal{M}\} = m.$$

Since \mathcal{M} is compact, there exists a subsequence of the sequence $\{a_k\}$, relabeled again as $\{a_k\}$, and a $a_0 \in \mathcal{M}$ such that $a_k \rightarrow a_0$. It follows from Lemma 2.1 that in particular

$$u(a_k, g, \varphi) \rightarrow u(a_0, g, \varphi) \text{ weakly in } \mathcal{W}, \text{ as } k \rightarrow +\infty.$$

Therefore, by lower semicontinuity of \mathcal{J} we have

$$m \leq \mathcal{J}(u(a_0, g, \varphi)) \leq \liminf_{k \rightarrow \infty} \mathcal{J}(u(a_k, g, \varphi)) = \lim_{k \rightarrow \infty} \mathcal{J}(u(a_k, g, \varphi)) = m.$$

Hence the result follows.

REMARK 3.1. In general, without convexity assumptions, we do not expect uniqueness of the optimal solution in identification ([1], [16]).

4. STABILITY RESULT.

In this section we give the main result of this paper on the dependence (hence also the stability) of the optimal elements for the problem (\mathcal{P}) on the data as well as on the cost functional.

We consider the sequence (indexed by the parameter $k \in \mathbf{N}$) of the identification problems:

$$(\mathcal{P}_k) \begin{cases} \text{find } a^* = \{a_{ij}^*\} \text{ in } \mathcal{M} \text{ so that} \\ \mathcal{J}_k(u(a^*, g_k, \varphi_k)) \leq \mathcal{J}_k(u(a, g_k, \varphi_k)) \text{ for all } a \in \mathcal{M}, \end{cases}$$

where $u(a, g_k, \varphi_k)$ are the solutions in \mathcal{W} to (1.1) - (1.3) corresponding to the perturbed data g_k, φ_k and \mathcal{J}_k are the perturbed functionals. We show that the set of optimal solutions to (\mathcal{P}_k) converges in some sense, to the set of optimal solution to (\mathcal{P}) .

We need the following continuous convergence of functionals. Let (\mathcal{X}, τ) be a topological space and $\mathcal{J}_k: \mathcal{X} \rightarrow \mathbf{R}$.

DEFINITION 4.1. We say that a sequence of functionals $\{\mathcal{J}_k\}$, $k \in \mathbf{N}$, is sequentially continuously convergent (shortly, C_{seq} -converges) to \mathcal{J} , and we write $\mathcal{J} = C_{seq}(\tau - \mathcal{X}) \lim_{k \rightarrow \infty} \mathcal{J}_k$, if for every $x \in \mathcal{X}$ and for every $\{x_k\} \subset \mathcal{X}$ which τ -converges to x , the sequence $\mathcal{J}_k(x_k)$ converges to $\mathcal{J}(x)$.

For each $k \in \mathbf{N}$, we denote by $\mathcal{S}, \mathcal{S}_k$ the sets of optimal elements to the problem $(\mathcal{P}), (\mathcal{P}_k)$, respectively, i.e.

$$\mathcal{S} = \mathcal{S}(g, \varphi, \mathcal{J}) = \{a^* \in \mathcal{M} : \mathcal{J}(u(a^*, g, \varphi)) \leq \mathcal{J}(u(a, g, \varphi)), \forall a \in \mathcal{M}\},$$

$$\mathcal{S}_k = \mathcal{S}(g_k, \varphi_k, \mathcal{J}_k) = \{a^* \in \mathcal{M} : \mathcal{J}_k(u(a^*, g_k, \varphi_k)) \leq \mathcal{J}_k(u(a, g_k, \varphi_k)), \forall a \in \mathcal{M}\}.$$

With the above notation we have

THEOREM 4.1. In addition to the hypotheses (H_2) and (H_4) , we assume that $\mathcal{J}, \mathcal{J}_k: \mathcal{W} \rightarrow \mathbf{R}$ are given weakly sequentially lower semicontinuous functionals. Let $g, g_k \in L^2(\Sigma)$ and $\varphi, \varphi_k \in H$ be such that $j(\varphi_k) \in L^1(\Omega)$. If

$$\{(g_k, \varphi_k)\} \rightarrow (g, \varphi) \text{ in } L^2(\Sigma) \times H, \text{ as } k \rightarrow +\infty, \quad (4.1)$$

$$\mathcal{J} = C_{seq}(w - \mathcal{W}) \lim_{k \rightarrow \infty} \mathcal{J}_k, \quad (4.2)$$

(where $w - \mathcal{W}$ stands for the weak topology in \mathcal{W}), then

$$\lim_{k \rightarrow \infty} h^*(\mathcal{S}_k, \mathcal{S}) = 0, \quad (4.3)$$

where $h^*(\cdot, \cdot)$ is defined in (1.5).

PROOF. We argue by contradiction. If (4.3) is not verified there exist $\bar{\varepsilon} > 0$ and a sequence $\{k_\nu\}$, $k_\nu \rightarrow +\infty$ such that

$$\bar{\varepsilon} < h^*(\mathcal{S}_{k_\nu}, \mathcal{S}), \forall k_\nu \in \mathbf{N}.$$

Clearly, there exist $a_{k_\nu}^* \in \mathcal{S}_{k_\nu}$ such that

$$\bar{\varepsilon} < d(a_{k_\nu}^*, \mathcal{S}), \forall k_\nu \in \mathbf{N}. \quad (4.4)$$

In view of compactness of \mathcal{M} , we deduce that there exists a subsequence of $\{a_{k_\nu}^*\}$, that we will denote in the same way, such that

$$a_{k_\nu}^* \rightarrow a^* \text{ in } (C(\bar{Q}))^2, \text{ as } k_\nu \rightarrow +\infty, \quad (4.5)$$

for some $a^* \in \mathcal{M}$.

Let now $u_{k_\nu}^* = u(a_{k_\nu}^*, g_{k_\nu}, \varphi_{k_\nu})$ and $u^* = u(a^*, g, \varphi)$ denote the solutions to (1.1) - (1.3) which correspond to the triples $(a_{k_\nu}^*, g_{k_\nu}, \varphi_{k_\nu})$ and (a^*, g, φ) , respectively. From (4.1) and (4.5) Lemma 2.1 gives

$$u_{k_\nu}^* \rightarrow u^* \text{ weakly in } \mathcal{W}, \text{ as } k_\nu \rightarrow +\infty.$$

Since the functionals \mathcal{J}_k C_{seq} -converges to \mathcal{J} , we get

$$\mathcal{J}(u^*) = \lim \mathcal{J}_{k_\nu}(u_{k_\nu}^*). \quad (4.6)$$

Let us fix an arbitrary $a \in \mathcal{M}$. Let $u_{k_\nu} = u(a, g_{k_\nu}, \varphi_{k_\nu})$ and $u = u(a, g, \varphi)$ be the solutions to (1.1) - (1.3) with the indicated data. From the continuous dependence on the data, we conclude that

$$u_{k_\nu} \rightarrow u \text{ weakly in } \mathcal{W}, \text{ as } k_\nu \rightarrow +\infty.$$

Hence, as above, we have

$$\mathcal{J}(u) = \lim \mathcal{J}_{k_\nu}(u_{k_\nu}). \quad (4.7)$$

Since $a_{k_\nu}^*$ are in \mathcal{S}_{k_ν} , we have

$$\mathcal{J}_{k_\nu}(u_{k_\nu}^*) \leq \mathcal{J}_{k_\nu}(u_{k_\nu}).$$

In the limit, as $k_\nu \rightarrow +\infty$, one gets from (4.6), (4.7) :

$$\mathcal{J}(u(a^*, g, \varphi)) \leq \mathcal{J}(u(a, g, \varphi)), \forall a \in \mathcal{M}.$$

From the arbitrariness of $a \in \mathcal{M}$, we have $a^* \in \mathcal{S}$ and this implies that

$$d(a_{k_\nu}^*, \mathcal{S}) \leq \|a_{k_\nu}^* - a^*\|. \quad (4.8)$$

But now from (4.5), we obtain that (4.8) contradicts (4.4). This proves (4.3) and completes the proof of the theorem.

The convergence (4.2) holds, for instance, for the sequence of functionals of the form (1.4). Namely, the following two cases can be considered:

(i) Let $\mathcal{Z} = \mathcal{H}$. Given the functionals $\mathcal{J}_k(u) = \|u - z_d^k\|_{\mathcal{H}}^2$, where $\{z_d^k\}$ is a sequence of perturbed distributed observations in \mathcal{H} converging to z_d , we have that (4.2) is satisfied with $\mathcal{J}(u) = \|u - z_d\|_{\mathcal{H}}^2$. This follows directly, by the definition of C_{seq} -limit, from the fact that $\mathcal{W} \subset \mathcal{H}$ and this embedding is compact (compare [15]).

(ii) Let $\mathcal{Z} = L^2(\Sigma)$. We consider the functionals $\mathcal{J}_k(u) = \|u - z_d^k\|_{L^2(\Sigma)}^2$, where $\{z_d^k\}$ is a sequence of perturbed measurements (observations) performed on Σ . If $z_d^k \rightarrow z_d$ in $L^2(\Sigma)$, as $k \rightarrow +\infty$, then using the compactness of the trace embedding from \mathcal{W} into $L^2(\Sigma)$, we easily get the convergence (4.2), where $\mathcal{J}(u) = \|u - z_d\|_{L^2(\Sigma)}^2$.

REMARK 4.1. One can generalize the results presented above to the case when $\mathcal{A}(t)$ is a differential operator of the form

$$\mathcal{A}(t) = -\frac{\partial}{\partial x_i}(a_{ij}(x, t)\frac{\partial}{\partial x_j}) + a(x, t)u.$$

Our theory, with some minor changes, can handle inverse problems involving the identification of any of the parameters a_{ij} , a , or initial data or boundary function.

REMARK 4.2. A further generalization of our results can be obtained considering the identification problems for (1.1) - (1.3) with the mixed boundary conditions

$$\frac{\partial u}{\partial \nu_{\mathcal{A}(t)}} + \beta_i(u) \ni g_i \text{ on } \Sigma_i, \quad i = 1, 2,$$

instead of (1.2), where $\Sigma_i = \Gamma_i \times (0, T)$ and Γ_i are the disjoint parts of Γ . The exact formulation of the results with obvious modifications is left to the interested reader.

ACKNOWLEDGEMENT. The paper was written while the author was visiting the Scuola Normale Superiore di Pisa and was supported by the Istituto Nazionale di Alta Matematica Francesco Severi, Rome, Italy. This work was partially funded by a National Research Project in Mathematics "Equazioni Differenziali e Calcolo delle Variazioni" of the Ministero dell'Università e della Ricerca Scientifica e Tecnologia, 40% contracts.

REFERENCES

1. AHMED, N.U. Optimization and Identification of Systems Governed by Evolution Equations on Banach Space, Pitman Research Notes in Mathematics 184, Longman, Boston - London - Melbourne, 1988.
2. BALAKRISHNAN, A.V. Identification of distributed parameter systems: non-computational aspects, Proceedings IFIP Working Conference, Rome, 1976, A.Ruberti, Ed., Lecture Notes in Control and Information Sciences, Vol.1, Springer-Verlag, Berlin, 1978, 1-10.
3. BAMBERGER, A., CHAVENT, G. and LAILLY, P. About the stability of the inverse problem in 1-D wave equations - applications to the interpretation of seismic profiles, Appl. Math. Optim. 5 (1979), 1-47.
4. BANKS, H.T. On a variational approach to some parameter estimation problems, ICASE Rep. No. 85-32, NASA Langley Res. Ctr., Hampton, VA, June, 1985.
5. BANKS, H.T. and ILES, D.W. On compactness of admissible parameter sets, Proceedings IFIP Working Conference, Gainesville, 1986, I.Lasiecka, R.Triggiani Eds., Lecture Notes in Control and Information Sciences, Vol.97, Springer-Verlag, Berlin, 1987, 130-142.

6. BARBU, V. Nonlinear Semigroups and Differential Equations in Banach Spaces, Noordhoff International Publishing, Leyden, The Netherlands, 1976.
7. BARBU, V. Boundary control problems with nonlinear state equation, SIAM J. Control Optim. 20 (1982), 125-143.
8. BENSOUSSAN, A., LIONS, J.L. and PAPANICOLAOU, G. Asymptotic Analysis for Periodic Structures, Studies in Mathematics and its Applications Ser., Vol.5, North-Holland, Amsterdam, 1978.
9. BREZIS, H. Problèmes unilatéraux, J. Math. Pures Appl. 51 (1972), 1-168.
10. COLONIUS, F. and KUNISCH, K. Output least squares stability for estimation of the diffusion coefficient in an elliptic equation, Proceedings IFIP Working Conference, Gainesville, 1980, I.Lasiecka, R.Triggiani Eds., Lecture Notes in Control and Information Sciences, Vol.97, Springer-Verlag, Berlin, 1987, 185-195.
11. COLONIUS, F. and KUNISCH, K. Stability for Parameter Estimation in Two Point Boundary Value Problems, J. Reine Angew. Math. 370 (1986), 1-29.
12. DUVAUT, G. and LIONS, J.L. Les Inéquations en Mécanique et en Physique, Dunod, Paris, 1972.
13. KRAVARIS, C. and SEINFELD, J.H. Identification of parameters in distributed parameter systems by regularization, SIAM J. Control Optim. 23 (1985), 217-241.
14. KUNISCH, K. Identification and Estimation of Parameters in Abstract Cauchy Problems, Mathematical Control Theory, Banach Center Publication, Vol.14., PWN, Warsaw, 1985, 277-298.
15. LIONS, J.L. Quelques méthodes de résolution des problèmes aux limites non linéaires, Dunod-Gauthier Villars, Paris, 1969.
16. LIONS, J.L. Some Aspects of Modelling Problems in Distributed Parameter Systems, Proceedings IFIP Working Conference, Rome, 1976, A.Ruberti, Ed., Lecture Notes in Control and Information Sciences, Vol.1, Springer-Verlag, Berlin, 1978, 11-41.
17. MURAT, F. Contre-exemples pour divers problèmes où le contrôle intervient dans les coefficients, Ann. Mat. Pura Appl. 112 (1977), 49-68.
18. MURAT, F. Control in coefficients, Systems and Control Encyclopedia, M.G.Singh Ed., Pergamon Press, 1989, 808-812.

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru