

**FIXED POINTS AND THEIR APPROXIMATIONS FOR ASYMPTOTICALLY
NONEXPANSIVE MAPPINGS IN LOCALLY CONVEX SPACES**

P. VIJAYARAJU

Department of Mathematics
Anna University
Madras 600 025
India

(Received March 19, 1992)

ABSTRACT. We construct an example that the class of asymptotically nonexpansive mappings include properly the class of nonexpansive mappings in locally convex spaces, prove a theorem on the existence of fixed points, and the convergence of the sequence of iterates to a fixed point for asymptotically nonexpansive mappings in locally convex spaces.

KEY WORDS AND PHRASES. Fixed points, asymptotically nonexpansive, uniformly asymptotically regular maps.

1992 AMS SUBJECT CLASSIFICATION CODES. 47H10, 46006.

1. INTRODUCTION

Some results concerning fixed point theorems for nonexpansive mappings on locally convex spaces have been obtained by Taylor [5], Su and Sehgal [6], Tarafdar [4] and others. Su and Sehgal [6] have extended a theorem of Taylor [5] for nonexpansive self-mapping of a compact star-shaped subset K of a locally convex space X , to nonexpansive non-self mapping T of K into X with $T(\partial K) \subset K$, where ∂K denotes the boundary of K . In 1972, Goebel and Kirk [2] have introduced the notion of asymptotically nonexpansive mappings in Banach spaces and they have proved fixed point theorems for such mappings in uniformly convex Banach spaces.

The author [7] has introduced in 1988 the notion of asymptotically nonexpansive mappings (see Def. 1.1 (iii)) and uniformly asymptotically regular mappings (see Def. 1.1 (iv)) in locally convex spaces X and showed in [7] that if K is a weakly compact star-shaped subset of X and $T: K \rightarrow K$ is asymptotically nonexpansive, uniformly asymptotically regular and $I - T$ is demiclosed, then T has a fixed point in K , where I denotes the identity map. In the second section of this paper, we prove that the

condition $T: K \rightarrow K$ in [7] may be weakened to $T: K \rightarrow X$ with $T^n(\partial K) \subset K$ for every positive integer n .

In the third section, we prove the convergence of the sequence of iterates to a fixed point for asymptotically regular, asymptotically nonexpansive self-mapping in a locally convex space. This result extends those of Theorem 3.3 of Taylor [5] for asymptotically regular, nonexpansive self-mappings.

Here and later, let X denote a locally convex Hausdorff linear topological space with a family $(p_\alpha)_{\alpha \in J}$ of seminorms which defines the topology on X , where J is any index set.

We recall the following definition.

DEFINITION 1.1. Let K be a nonempty subset of X . If T maps K into X , we say that

i) T is contractive (i.e., p_α -contractive) [6] if

$$\begin{aligned} p_\alpha(Tx - Ty) &< p_\alpha(x - y) & \text{if } p_\alpha(x - y) \neq 0 \\ &= 0 & \text{if } p_\alpha(x - y) = 0 \end{aligned}$$

for each $x, y \in K$ and for each $\alpha \in J$;

ii) T is nonexpansive (i.e., p_α -nonexpansive) [6] if

$p_\alpha(Tx - Ty) \leq p_\alpha(x - y)$ for each $x, y \in K$ and for each $\alpha \in J$;

iii) T is asymptotically nonexpansive [7] if

$$p_\alpha(T^n x - T^n y) \leq k_n p_\alpha(x - y)$$

for each $x, y \in K$, for each n and for each $\alpha \in J$, where $\{k_n\}$ is a sequence of real numbers such that $k_n \rightarrow 1$ as $n \rightarrow \infty$.

It is assumed that $k_n \geq 1$ and $k_n \geq k_{n+1}$ for $n = 1, 2, \dots$;

iv) T is uniformly asymptotically regular [7] if for each α in J and $\eta > 0$, there exists a $N(\alpha, \eta)$ such that

$$p_\alpha(T^n x - T^{n+1} x) < \eta$$

for all $n \geq N(\alpha, \eta)$ and for all $x \in K$; and

v) T is asymptotically regular on K [4] if, for each $x \in K$ and $\alpha \in J$,

$$\lim_{n \rightarrow \infty} p_\alpha(T^n x - T^{n+1} x) = 0.$$

DEFINITION 1.2. A mapping T from K to X is said to be demiclosed [5] if, for every net (x_β) in K such that (x_β) weakly converges to x in K (i.e., $x_\beta \rightarrow x$) and (Tx_β) converges to y in X (i.e., $Tx_\beta \rightarrow y$) we have $Tx = y$.

The following example shows that the class of asymptotically nonexpansive mappings is wider than the class of nonexpansive mappings in locally convex spaces.

EXAMPLE 1.1. Let $X = \text{space}(s)$, the space of all sequences of complex numbers whose topology is defined by the family of seminorms p_n defined by

$$p_n(x) = \max_{1 \leq i \leq n} |\xi_i| \text{ for } x = (\xi_1, \xi_2, \dots) \in X \text{ and } n = 1, 2, \dots$$

Let $K = \{x = (\xi_1, \xi_2, \dots) \in X: |\xi_1| \leq 1/2 \text{ and } |\xi_j| \leq 1 \text{ for } j = 2, \dots\}$. Define a map T from K to K by

$$Tx = (0, 2\xi_1, A_2\xi_2, \dots, A_k\xi_k, \dots)$$

for all $x = (\xi_1, \xi_2, \dots, \xi_k, \dots) \in K$, where $\{A_i\}$ is a sequence of real numbers in $(0, 1)$ such that $\prod_{i=2}^{\infty} A_i = 1/2$.

Let $a = (1/2, 0, \dots)$, $b = (0, \dots) \in K$. Then we have

$$p_2(Ta - Tb) = 1 > 1/2 = p_2(a - b)$$

and hence T is not nonexpansive.

Now, let $x = (\xi_1, \xi_2, \dots, \xi_k, \dots)$, $y = (\eta_1, \eta_2, \dots, \eta_k, \dots) \in K$.

Then $p_n(Tx - Ty) \leq 2 p_n(x - y)$ for $n = 1, 2, \dots$ and

$$T^m(x) = (0, \dots, 0, 2 \prod_{i=2}^m A_i \xi_1, \prod_{i=2}^{m+1} A_i \xi_2, \dots, \prod_{i=k}^{m+k-1} A_i \xi_k, \dots).$$

Therefore $p_n(T^m(x) - T^m(y)) = 0$ for $m \geq n$.

If $m < n$, then $m = n - k$, where $k > 0$ and $n > k$

and therefore $p_n(T^m(x) - T^m(y))$

$$\begin{aligned} &= \max \left[\left[2 \prod_{i=2}^m A_i \right] |\xi_1 - \eta_1|, \left[\prod_{i=2}^{m+1} A_i \right] |\xi_2 - \eta_2|, \dots, \left[\prod_{i=k}^{m+k-1} A_i \right] |\xi_k - \eta_k| \right] \\ &\leq \max \left[2 \prod_{i=2}^m A_i, \prod_{i=2}^{m+1} A_i, \prod_{i=3}^{m+2} A_i, \dots, \prod_{i=k}^{m+k-1} A_i \right] p_k(x - y) \\ &\leq 2 \prod_{i=2}^m A_i p_n(x - y) = k_m p_n(x - y), \end{aligned}$$

where $k_m = 2 \prod_{i=2}^m A_i \rightarrow 1$ as $m \rightarrow \infty$.

Hence T is asymptotically nonexpansive. Also T is uniformly asymptotically regular on K .

The following example due to the author in [7] shows that the uniform asymptotic regularity is stronger than asymptotic regularity.

EXAMPLE 1.2. Let $X = \ell^p$, $1 < p < \infty$. Let K denote the unit ball in X . Define a map $T: K \rightarrow K$ by

$$Tx = (\xi_2, \xi_3, \dots) \text{ for all } x = (\xi_1, \xi_2, \dots) \in K.$$

Then T is asymptotically regular but not uniformly asymptotically regular on K . Also T is nonexpansive and hence T is asymptotically nonexpansive.

DEFINITION 1.3. A nonempty subset K of X is said to be star-shaped [1] provided that there is at least one element x in K such that if y is any element of K and $t \in (0, 1)$, then $(1-t)x + ty \in K$. Such a point x is

called a star-center of K . Every convex set is a star-shaped set but the converse is not true.

MAIN RESULTS

2. FIXED POINTS OF ASYMPTOTICALLY NONEXPANSIVE MAPPINGS

For the proof of Theorem 2.1, we need the following lemma due to Su and Sehgal [6, Theorem 2].

LEMMA 2.1. Let K be a nonempty compact subset of X . Let T be a contractive mapping of K into X such that $T(\partial K) \subset K$. Then T has a unique fixed point in K .

Taylor [5] has proved a result on the existence of fixed points for nonexpansive self-mapping T of a nonempty compact star-shaped subset K of a locally convex space X . This result was extended by Su and Sehgal [6] to nonexpansive non-self mapping T of K into X by assuming the condition that $T(\partial K) \subset K$. We extend the corresponding theorem for asymptotically nonexpansive, uniformly asymptotically regular mappings. The following theorem is new even in the case of Banach spaces.

THEOREM 2.1. Let T be a mapping of X into itself. Let K be a nonempty compact star-shaped subset of X . Let T be an asymptotically nonexpansive, uniformly asymptotically regular mapping of K into X such that $T^n(\partial K) \subset K$ for every $n = 1, 2, \dots$. Then T has a fixed point in K .

PROOF. Let y be a star center of K . Define a map T_n from K to X by

$$T_n x = a_n T^n x + (1-a_n) y \text{ for all } x \in K, n = 1, 2, \dots,$$

where $a_n = (1 - (1/n)) / k_n$ and $\{k_n\}$ is as in Definition 1.1 (iii). Then each T_n clearly maps K into X .

If $x, z \in K$, then since T is asymptotically nonexpansive, we have

$$p_\alpha(T_n x - T_n z) = a_n p_\alpha(T^n x - T^n z) \leq (1 - (1/n)) p_\alpha(x - z).$$

Therefore T_n is a contraction of K into X and hence a contractive mapping of K into X .

Since $T^n(\partial K) \subset K$ and K is star-shaped, $T_n(\partial K) \subset K$.

Therefore by Lemma 2.1, T_n has a unique fixed point, say, x_n in K .

Therefore, $x_n - T^n x_n = (1 - a_n) (y - T^n x_n) \rightarrow 0$ as $n \rightarrow \infty$, since K is bounded and $a_n \rightarrow 1$ as $n \rightarrow \infty$. (2.1)

Since T is uniformly asymptotically regular, it follows that

$$T^n x_n - T^{n+1} x_n \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (2.2)$$

From (2.1) and (2.2) we obtain

$$T^{n+1} x_n - x_n \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (2.3)$$

Now

$$\begin{aligned} p_\alpha(T x_n - x_n) &\leq p_\alpha(T x_n - T^{n+1} x_n) + p_\alpha(T^{n+1} x_n - x_n) \\ &\leq k_1 p_\alpha(x_n - T^n x_n) + p_\alpha(T^{n+1} x_n - x_n). \end{aligned} \quad (2.4)$$

Using (2.1) and (2.3) in (2.4) we get

$$Tx_n - x_n \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (2.5)$$

Since K is compact and $\{x_n\} \subset K$, there is a subnet (x_β) of the sequence $\{x_n\}$ such that $x_\beta \xrightarrow{\beta} x \in K$.

Therefore $(I - T)(x_\beta) \xrightarrow{\beta} (I - T)x$ and by (2.5), $(I - T)(x_\beta) \xrightarrow{\beta} 0$. Since X is Hausdorff, it follows that $(I - T)x = 0$. Thus x is a fixed point of T in K .

3. CONVERGENCE OF ITERATES OF ASYMPTOTICALLY NONEXPANSIVE MAPPING

Taylor [5] has proved that the sequence of iterates converges to a fixed point for nonexpansive self-mapping in a locally convex space. This result is extended below to asymptotically nonexpansive self-mapping.

We use the following definition to prove our Theorem 3.2.

DEFINITION 3.1. A point x in a topological space X is called a cluster point [3] of a net S if and only if S is frequently in every neighbourhood of x .

THEOREM 3.2. Let K be a nonempty closed bounded subset of X . Let T be a continuous, asymptotically regular self-mapping of K . Assume that $I - T$ maps closed subsets of K into closed subsets of X , where I denotes the identity mapping. Then, for each $x \in K$, the sequence of iterates $\{T^n x\}$ clusters at a fixed point of T and each such cluster point is fixed by T . If, in addition, T is an asymptotically nonexpansive self-mapping of K , then every sequence $\{T^n x\}$ converges to a fixed point of T .

PROOF. Let T be a continuous, asymptotically regular self-mapping of K . Let $x \in K$ and M denote the closure of $\{T^n x\}$. Since T is asymptotically regular, it follows that

$$T^n x - T^{n+1} x \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore 0 lies in the closure of $(I - T)(M)$. Since M is closed and $I - T$ maps closed subsets of K into closed subsets of X , it follows that $(I - T)(M)$ is closed. Therefore $0 \in (I - T)(M)$ and hence there is a point y in M such that $(I - T)(y) = 0$.

Since $y \in M$, either $y \in \{T^n x\}$ or y is a cluster point of $\{T^n x\}$.

If $y = T^m x$ for some m , then

$$T^{n+m}(x) = T^n(T^m x) = T^n y = y \quad \text{for } n = 1, 2, \dots$$

Therefore $T^k x = y$ if $k > m$. Hence y is a cluster point of $\{T^n x\}$.

Let z be any cluster point of $\{T^n x\}$.

We know that a point b in a topological space X is a cluster point of a net S if and only if some subnet of S converges to b [3].

Therefore there is a subnet $(T^\beta x)$ of $\{T^n x\}$ such that $T^\beta x \xrightarrow{\beta} z$.

$$\text{Hence } (I - T)z = (I - T)\lim_{\beta} T^{\beta}x = \lim_{\beta} (I - T)(T^{\beta}x),$$

since $(I - T)$ is continuous.

$= 0$, since T is asymptotically regular on K .

Thus z is a fixed point of T .

Assume further that T is an asymptotically nonexpansive self-mapping of K . We already know that y is a cluster point of $\{T^n x\}$.

Therefore for each $\alpha \in J$ and $\delta > 0$, there exists an integer m such that

$$p_{\alpha}(T^m x - y) < \delta. \quad (3.1)$$

Since T is asymptotically nonexpansive, it follows that

$$p_{\alpha}(T^{n-m}(T^m x) - T^{n-m}y) \leq k_{n-m} p_{\alpha}(T^m x - y) \text{ for } n \geq m.$$

From (3.1) we obtain

$$p_{\alpha}(T^n x - y) < k_{n-m} \delta.$$

Therefore

$$\limsup_{n \rightarrow \infty} p_{\alpha}(T^n x - y) \leq \limsup_{n \rightarrow \infty} k_{n-m} \delta \leq \delta.$$

$$\text{Hence } \lim_{n \rightarrow \infty} p_{\alpha}(T^n x - y) = 0.$$

That is, the sequence $\{T^n x\}$ converges to a fixed point of T .

ACKNOWLEDGEMENT. The author would like to thank Prof.T.R.Dhanapalan for his guidance and encouragement in the preparation of this paper. The author wishes to thank the referees for suggestions that lead to a better presentation of this paper.

REFERENCES

1. DOTSON, W.G. (JR). Fixed point theorems for nonexpansive mappings on star-shaped subsets of Banach spaces, J. London Math. Soc. **4** (1972), 408-410.
2. GOEBEL, K. and W.A.KIRK. A fixed point theorem for asymptotically nonexpansive mappings, Proc. Amer. Math. Soc. **35** (1972), 171-174.
3. KELLEY, J.L. General Topology, Van Nostrand, Princeton, NJ, 1955.
4. TARAFDAR, E. An approach to fixed point theorems on uniform spaces, Trans. Amer. Math. Soc. **191** (1974), 209 - 225.
5. TAYLOR, W.W. Fixed point theorems for nonexpansive mappings in linear topological spaces, J. Math. Anal. Appl. **40** (1972), 164-173.
6. SU.C.H. and V.M.SEHGAL. Some fixed point theorems for nonexpansive mappings in locally convex spaces, Boll. Un. Mat. Ital. **10** (1974), 598-601.
7. VIJAYARAJU, P. Fixed point theorems for asymptotically nonexpansive mappings, Bull. Calcutta Math. Soc. **80** (1988), 133-136.

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru