

ON CHAINS AND POSETS WITHIN THE POWER SET OF A CONTINUUM

P. T. MATTHEWS and T. B. M. McMASTER

Pure Mathematics Department
Queen's University
Belfast BT7 1NN, Northern Ireland

(Received May 6, 1993 and in revised form October 27, 1993)

ABSTRACT. Transfinite induction is employed to construct a copy of an arbitrary partially-ordered set of cardinality at most c within the power set (quasi-ordered by sub-chain embeddability) of the real line.

KEY WORDS AND PHRASES. Partially-ordered set, sub-chain embeddability, real line.

1991 AMS SUBJECT CLASSIFICATION CODES. 06A05, 06A06.

1. INTRODUCTION.

One way to explore the structure of a quasi-ordered set X is to seek subsets of it which, under the induced order, are partially- or totally-ordered: for instance the behavior of chains within X is closely related, through a variant of Zorn's lemma, to the existence of elements that are in some sense [4] maximal or minimal in the quasi-order. In her doctoral thesis [2] Matier employed ideas of Stephen Watson to carry out one such investigation on the power set of \mathbb{R} ordered not by set-inclusion but by sub-chain embeddability. She demonstrated that this quasi-ordered set contains an infinite antichain, and hence deduced that the family of posets on c points or fewer (ordered by sub-poset embeddability) contains an infinite decreasing sequence. This finding has relevance to the behavior of the total negation operation, defined for topological spaces by Bankston [1], when it is applied to partially-ordered topological spaces (see [3] for a brief account).

This note makes use of a modification of the Watson-Matier argument to establish a stronger conclusion about the set of subsets of \mathbb{R} ; namely, that it contains not only infinite antichains and chains, but also copies of every partially-ordered set whose cardinality does not exceed c . An initial examination is also presented of the circumstances (in terms of set-theoretic axioms assumed) in which analogous results may be obtained for higher cardinals.

LEMMA A. Let C be an arbitrary chain, A a non-empty subset of C and $f: A \rightarrow C$ a strictly increasing mapping. If every open interval in C contains a fixed point for f then f is id_A .

PROOF. Suppose that there exists $x \in A$ such that $x \neq f(x)$. Then either $x < f(x)$ or $x > f(x)$. In the first case, $y \in (x, f(x))$ implies $x < y$ giving $f(x) < f(y)$, so $f(y) \notin (x, f(x))$ which in turn implies $y \neq f(y)$: thus $(x, f(x))$ contains no fixed point for f . In the second case, $(f(x), x)$ contains no such point.

DEFINITION. Let us call an infinite cardinal α *continuum-like* if

- (i) $\alpha = 2^\beta$ for some (infinite) $\beta < \alpha$ and

- (ii) there exists a chain C of cardinality α with the following properties:
 - (a) each open interval in C has cardinality α and
 - (b) there is a subset Q of C such that $\text{card}(Q) = \beta$ and every open interval in C intersects Q .

Clearly, c itself is a continuum-like cardinal. We shall address the question of the existence of other continuum-like cardinals later in this article.

Given a strictly increasing function $f:Q \rightarrow C$ (where Q and C are as above) and an element x of $C \setminus Q$, consider the set:

$$A := \{\bar{f}(x) : \bar{f} \text{ is a strictly increasing extension of } f \text{ over } Q \cup \{x\}\}.$$

Whenever this set is a singleton, we shall use the notation $f!(x)$ for its unique element. We make the following definitions:

- (i) x is a *non-extension point* for f if $A = \emptyset$,
- (ii) x is a *trivial-extension point* for f if $\text{card}(A) = 1$ and $f!(x) \in Q$,
- (iii) x is a *unique-extension point* for f if $\text{card}(A) = 1$ and $f!(x) \in C \setminus Q$,
- (iv) x is a *multi-extension point* for f if $\text{card}(A) > 1$.

It is clear that the four classes of points defined here partition $C \setminus Q$ and we note that there are at most β trivial-extension points for f (for otherwise there would exist x, y in $C \setminus Q$ with $x < y$ and $f!(x) = f!(y) \in Q$, contradicting the strictly increasing nature of f). By considering the example $C = \mathbb{R}, Q = \mathbb{Q}, f(x) = x\sqrt{2}$ it is apparent that the number of trivial-extension points can be as high as β . It can also be as low as zero, as in the case $C = \mathbb{R}, Q = \mathbb{Q}, f(x) = x$. Somewhat less obvious is the observation that the number of multi-extension points is likewise constrained to lie between 0 and β :

LEMMA B. Let α be a continuum-like cardinal, C and Q as described in the definition. A given strictly increasing function $f:Q \rightarrow C$ has at most β multi-extension points.

PROOF. For each multi-extension point y for f we can choose elements t_y^1, t_y^2 of C such that $t_y^1 < t_y^2$ and that

$$\bar{f}_y^1(x) = \begin{cases} f(x) & \text{if } x \in Q, \\ t_y^1 & \text{if } x = y, \end{cases}$$

$$\bar{f}_y^2(x) = \begin{cases} f(x) & \text{if } x \in Q, \\ t_y^2 & \text{if } x = y \end{cases}$$

define two distinct strictly increasing extensions \bar{f}_y^1, \bar{f}_y^2 of f over $Q \cup \{y\}$. Let I_y denote the interval (t_y^1, t_y^2) , and note that the family

$$\{I_y : y \text{ is a multi-extension point for } f\}$$

is pairwise-disjoint: for if z and y are two multi-extension points for f with $z < y$, we can choose $q \in Q$ with $z < q < y$ and observe that for any $a \in I_z, b \in I_y$:

$$\begin{aligned} a < t_z^2 &= \bar{f}_z^2(z) < \bar{f}_z^2(q) = f(q) \\ &= \bar{f}_y^1(q) < \bar{f}_y^1(y) = t_y^1 < b \end{aligned}$$

so $a \neq b$. Since each of the I_y contains a point of Q and $\text{card}(Q) = \beta$, the result follows.

COROLLARY. In the same notation, every open interval in C contains either α non-extension points for f or α unique-extension points for f .

Let $\mathcal{P}_Q(C)$ denote the set of all those subsets of C which contain Q and consider it as a quasi-ordered set (poset) under subchain embeddability: that is, given $A, B \in \mathcal{P}_Q(C)$ we write $A \leq B$ if and only if A is order-isomorphic to a subset of B (where A and B inherit the order on C).

THEOREM. Let S be a given partially-ordered set of cardinality α . There is a subset of $\mathcal{P}_Q(C)$ which is isomorphic to S .

PROOF. Denote by \mathfrak{F} the set of strictly increasing functions from Q into C . Since $\text{card}(\mathfrak{F}) \leq \alpha^\beta = \alpha$, $\mathfrak{F} \times S$ has cardinality α and can be expressed as the range of an α -sequence:

$$\mathfrak{F} \times S = \{(f_t, s_t) : t \in \alpha\}$$

where we are viewing α as an ordinal. Make an arbitrary choice of $q_0 \in Q$. Transfinite induction will now serve to construct three α -sequences $(x_\delta, \delta \in \alpha), (y_\delta, \delta \in \alpha), (z_\delta, \delta \in \alpha)$ in the set $(C \setminus Q) \cup \{q_0\}$.

Let $\gamma \in \alpha$ and suppose that we have already chosen, for each $\delta < \gamma$ in α , elements $x_\delta, y_\delta, z_\delta$ of C such that

- (i) $x_\delta, y_\delta \in (C \setminus Q) \cup \{q_0\}, z_\delta \in C \setminus Q$,
- (ii) all choices are distinct except for repetitions of q_0 ,
- (iii) whenever $f_\delta = \text{id}_Q$ then $x_\delta = y_\delta = q_0$,
- (iv) whenever $f_\delta \neq \text{id}_Q$ then
 - either x_δ is a unique-extension point for f_δ and $y_\delta = f_\delta!(x_\delta)$
 - or x_δ is a non-extension point for f_δ and $y_\delta = q_0$.

Now if $f_\gamma = \text{id}_Q$ choose $x_\gamma = q_0, y_\gamma = q_0$ and, bearing in mind that the cardinality of $C \setminus Q$ exceeds that of the set of all previously-made choices, select z_γ in $C \setminus Q$ distinct from all the x_δ, y_δ and z_δ for $\delta < \gamma$. On the other hand, suppose $f_\gamma \neq \text{id}_Q$. If f_γ possesses α non-extension points then choose one which is different from all preceding choices, denoting it by x_γ , put $y_\gamma = q_0$ and assign to z_γ any value in $C \setminus Q$ distinct from all previous selections. If not, then f_γ must have a strictly increasing extension f_γ^* over a subset D of C such that $C \setminus D$ has cardinality less than α ; since each interval in C has α elements, this D will therefore be order-dense. An appeal to Lemma A and the Corollary guarantees the existence of an open interval J_γ in C which is free from fixed points of f_γ^* and contains α unique-extension points for f_γ . Once again, since fewer than α points have previously been identified we can select one of these α unique-extension points x_γ in such a way that x_γ and $f_\gamma^*(x_\gamma)$ differ from all preceding choices, and note that $f_\gamma^*(x_\gamma) \neq x_\gamma$ since $x_\gamma \in J_\gamma$; pick also $z_\gamma \in C \setminus Q$ distinct from all other chosen elements. This completes the inductive step, and we are accordingly assured of the existence of α -sequences $(x_\delta), (y_\delta), (z_\delta)$ satisfying the above conditions (i) to (iv) for every δ in α .

For each s in the poset S (order denoted by \leq) we now define

$$I_s = \{x_\delta, z_\delta : s_\delta \leq s\}.$$

It is immediate from the definition that $r \leq s$ implies $I_r \subseteq I_s$ and therefore $Q \cup I_r$ trivially embeddable into $Q \cup I_s$.

Supposing now that $r \not\leq s$ in S , consider the hypothesis that $Q \cup I_r$ could be embedded in $Q \cup I_s$. Then we could find a strictly increasing function

$$j: Q \cup I_r \rightarrow Q \cup I_s.$$

The pair $(j|_Q, r)$ belongs to $\mathfrak{F} \times S$ and is therefore listed as (f_δ, s_δ) for some $\delta \in \alpha$. Two possibilities must be considered.

(I) $j|_Q = id_Q$. Here Lemma A implies that j is the identity map on $Q \cup I_r$, giving $Q \cup I_r \subseteq Q \cup I_s$. Yet since $s_\delta = r \notin s, z_\delta \in I_r$ but $z_\delta \notin I_s$, yielding a contradiction.

(II) $j|_Q \neq id_Q$. This time, x_δ will be either a unique-extension point or a non-extension point for f_δ . In the first case, since $x_\delta \in I_r$ and $j|_{Q \cup \{x_\delta\}}$ is strictly increasing,

$$j(x_\delta) = j|_{Q \cup \{x_\delta\}}(x_\delta) = f_\delta!(x_\delta) = y_\delta$$

forces y_δ to belong to $Q \cup I_s$, contrary to the definitions. In the second, no strictly increasing extension of $j|_Q$ over $Q \cup \{x_\delta\}$ could exist: and yet, as we saw in the discussion of the first case, $j|_{Q \cup \{x_\delta\}}$ is such an extension.

We conclude that, when $r \notin s$, no order-embedding of $Q \cup I_r$ into $Q \cup I_s$ can be obtained. Thus the map

$$\theta: S \rightarrow \mathcal{P}_Q(C)$$

defined by $\theta(s) = Q \cup I_s$ satisfies the condition

$$r \leq s \text{ if and only if } \theta(r) \leq \theta(s)$$

and establishes an order-isomorphism between S and the sub-poset $\theta(S)$ of the poset $\mathcal{P}_Q(C)$.

COROLLARY 1. Given any poset S with $\text{card}(S) \leq \alpha$, we can find a subset of $\mathcal{P}_Q(C)$ which is isomorphic to S .

PROOF. Extend S in any fashion to yield a poset S^* of cardinality α . By the theorem there is an embedding

$$\theta: S^* \rightarrow \theta(S^*) \subseteq \mathcal{P}_Q(C).$$

The restriction of θ to S now embeds the latter into $\mathcal{P}_Q(C)$ as required.

COROLLARY 2. Any poset of cardinality not exceeding c can be embedded in $\mathcal{P}_Q(\mathbb{R})$.

NOTE. The question of which infinite cardinals are continuum-like appears difficult to resolve fully, and will certainly depend to some extent on the axiom system adopted. For instance, if we assume the negation of the continuum hypothesis (*CH*), so that $\aleph_1 < c$, it will evidently be impossible to express \aleph_1 in the form 2^β , whence \aleph_1 will not be continuum-like: this contrasts with its status when *CH* itself is assumed. One positive result is fairly easy to obtain: it will follow from the generalized continuum hypothesis (*GCH*) that every successor cardinal α is continuum-like. For let β be the immediate cardinal predecessor of α , so that *GCH* implies $\alpha = 2^\beta$, and define

$$A = \{\text{all } \beta\text{-sequences of 0s and 1s that are ultimately constant at 0}\}.$$

Then

$$\begin{aligned} \text{card}(A) &\leq \sum_{\delta < \beta} 2^\delta \leq \sum_{\delta < \beta} \beta \text{ [using GCH again]} \\ &= \beta^2 = \beta < \alpha. \end{aligned}$$

Next put $C = \{\text{all } \beta\text{-sequences of 0s and 1s}\} \setminus A$ and impose on C the lexicographic ordering, converting it into a chain with α elements.

Let $x = (x_\gamma, \gamma \in \beta)$ and $y = (y_\gamma, \gamma \in \beta)$ be any elements of C for which $x < y$. Then there

must exist $\delta < \beta$ for which $x_\delta = 0$ and $y_\delta = 1$; let ζ denote the least cardinal for which $y_\zeta = 1$ and $\zeta > \delta$ (recalling that $y \notin A$). Now any $z = (z_\gamma, \gamma \in \beta)$ for which

$$\begin{aligned} z_\gamma &= y_\gamma & \text{if } \gamma < \zeta, \\ z_\zeta &= 0 \end{aligned}$$

will lie in the open interval (x, y) of C . There are 2^β such sequences z , so every interval in C has cardinality α as required.

Now if $Q = \{\text{all } \beta\text{-sequences of 0s and 1s that are ultimately constant at 1}\}$ we again have $\text{card}(Q) \leq \beta$; indeed, equality occurs here since it is easy to exhibit β distinct elements of Q (those consisting simply of a ‘block’ of 0s followed by a ‘block’ of 1s). In the previous paragraph, the z constructed to lie between x and y could have been chosen to have $z_\eta = 1$ for all $\eta > \zeta$, therefore belonging to Q : this verifies that Q is order-dense in C and concludes the demonstration.

REFERENCES

1. BANKSTON, P., The total negation of a topological property, *Illinois J. Math.* **23** (1979), 241-252.
2. MATIER, J., Total negation in general topology and in ordered topological spaces, Ph.D. thesis, Queen’s University Belfast, 1991.
3. MATIER, J. and McMASTER, T.B.M., Iteration of the ‘anti’ operation in ordered topological spaces and in other categorical contexts, *Boll. Un. Mat. Ital.*, to appear.
4. MATTHEWS, P.T., and McMASTER, T.B.M., A viewpoint on minimality in topology, *Bull. Irish Math. Soc.*, submitted.

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk