

WEAKLY COMPACTLY GENERATED FRECHET SPACES

SURJIT SINGH KHURANA

Department of Mathematics
The University of Iowa
Iowa City, Iowa 52242
U.S.A.

(Received on March 12, 1979 and in revised form July 13, 1979)

ABSTRACT. It is proved that a weakly compact generated Frechet space is Lindelöf in the weak topology. As a corollary it is proved that for a finite measure space every weakly measurable function into a weakly compactly generated Frechet space is weakly equivalent to a strongly measurable function.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES. Primary 28A40, 60B05, Secondary 28A20, 46A99, 54E50.

KEYWORDS AND PHRASES. Lindelöf spaces, weakly measurable functions, strongly measurable functions, universally measurable functions, Frechet spaces, weakly compact sets.

1. INTRODUCTION.

If E is a weakly compactly generated Banach space then it is proved in [7] that E , with weak topology, is Lindelöf. (A topological space is said to be Lindelöf if its every open covering has a countable subcovering.) In this note we extend this result to the case when E is a weakly compactly generated

Frechet space. Also, some consequences are obtained. All locally convex spaces are taken over the field of real numbers. By a Frechet space we mean a Hausdorff, metrizable, complete locally convex space; we use the notations of [4] for locally convex spaces. E' will always denote the topological dual of a locally convex space E . A locally convex space is said to be weakly compactly generated if there exists an increasing sequence of $\sigma(E, E')$ -compact subsets of E whose union is dense in E .

THEOREM 1. Let E be a weakly compactly generated Frechet space. Then $(E, \sigma(E, E'))$ is a Lindelöf space and E is a Borel subset of $(E', \sigma(E', E'))$, E' being the bidual of E .

PROOF. Let $\{V_n\}$ be a sequence of 0-nbd. base having the properties:

- (i) each V_n is absolutely convex and closed,
- (ii) $(n+1)V_{n+1} \subset V_n$, for every n .

We take $\{A_n\}$ for an increasing sequence of weakly compact, absolutely convex subsets of E such that $\bigcup_{n=1}^{\infty} A_n = H$ is dense in E . We identify $(E, \sigma(E, E'))$ as a subspace of $R^{E'}$, with product topology. $R^{E'}$ is a subset of the compact Hausdorff space $\overline{R}^{E'}$, where $\overline{R} = [-\infty, \infty]$. For an $x \in R^{E'}$ and $y \in \overline{R}^{E'}$, $x+y \in \overline{R}^{E'}$ has the natural meaning. For a compact set $A \subset R^{E'}$ and a compact set $B \subset \overline{R}^{E'}$, $A+B$ is compact. Thus $A_k + \overline{V}_n$ is a compact subset of $\overline{R}^{E'}$ for each k and n , \overline{V}_n being the closure of V_n in $\overline{R}^{E'}$. We claim that $\bigcap_{n=1}^{\infty} (H + \overline{V}_n) = E$. Since H is dense in E and V_n is a 0-nbd., $H + V_n \supset E$ for every n and so $\bigcap_{n=1}^{\infty} (H + \overline{V}_n) \supset E$. Conversely, take an $x \in \bigcap_{n=1}^{\infty} (H + \overline{V}_n)$. This means there exists a sequence $\{h_n\} \subset H$ and a sequence $\{z_n\}$ with $z_n \in \overline{V}_n$ for each n , such that $x = h_n + z_n$ for each n . Fix $n_0 \in \mathbb{N}$ and $\epsilon > 0$. Choose an $n_1 > \max(n_0, \frac{1}{\epsilon})$ and take an $n > n_1$. Since $V_{n_0} \supset nV_n$, $|f(z_n)| \leq \frac{1}{n} < \frac{1}{n_1} < \epsilon$, for every $f \in \overset{\circ}{V}_{n_0}$ the polar of V_{n_0} ([4]). Thus $f(x - h_n) \rightarrow 0$, uniformly for $f \in \overset{\circ}{V}_{n_0}$. From this it follows that $\{h_n\}$ is Cauchy in E which is complete. If $h_n \rightarrow y$ in E it

is easy to verify that, as elements of $\overline{R}^{E'}$, $x=y$. This proves the claim. Thus, in weak topology, $E = \bigcap_{n=1}^{\infty} \bigcup_{k=1}^{\infty} (A_k + \overline{V}_n)$ is analytic and so is Lindelöf ([6]). Also $(E'', \sigma(E'', E'))$ can be considered as a subspace of $R^{E'}$. Since $(A_k + \overline{V}_n)$ is compact in $\overline{R}^{E'}$, $(A_k + \overline{V}_n) \cap E''$ is closed in $(E'', \sigma(E'', E'))$ and so $(H + \overline{V}_n) \cap E''$ is Borel in $(E'', \sigma(E'', E'))$. Since $E = \bigcap_{n=1}^{\infty} (H + \overline{V}_n) \cap E''$, it follows that E is Borel in $(E'', \sigma(E'', E'))$.

REMARK. Similar results for Banach spaces are proved in [2, Cor. 3.2].

In the following result, some results and notations of ([3]) are used. Let (X, \mathfrak{A}, μ) be a finite measure space, E a Hausdorff locally convex space. A function $f: X \rightarrow E$ is called weakly measurable if $h \circ f$ is μ -measurable for every $h \in E'$. It is proved in ([2]) that if $f: X \rightarrow E$ is weakly measurable that the image measure $\nu: \mathcal{B} \rightarrow R$, $\nu(B) = \mu(f^{-1}(B))$, is a Baire measure on $(E, \sigma(E, E'))$, \mathcal{B} being the class of all Baire subsets of $(E', \sigma(E, E'))$ ([2], [8]). Two weakly measurable functions $f_i: X \rightarrow E$, $i=1,2$ are said to be weakly equivalent if $h \circ f_1 = h \circ f_2$ a.e. $[\mu]$, for every $h \in E'$. If E is Frechet then $f: X \rightarrow E$ is called strongly measurable if there exists a sequence $\{f_n\}$ of \mathfrak{A} -simple functions, $f_n: X \rightarrow E$, such that $f_n \rightarrow f$, pointwise a.e. $[\mu]$.

COROLLARY 2. Let (X, \mathfrak{A}, μ) be a finite measure space, E a weakly compactly generated Frechet space, and $f: X \rightarrow E$ a weakly measurable function. Then f is weakly equivalent to a strongly measurable function.

PROOF. By ([3], Cor. 5) it is enough to show that image Baire measure on $(E, \sigma(E, E'))$ is tight (cf. [2]). Since $(E, \sigma(E, E'))$ is Lindelöf, Baire measures are τ -additive (normal in the terminology of [5],[8]). By ([5], Theorems 3.3, 3.4) every Frechet space is universally measurable and so every τ -smooth measure is tight. This proves the result.

REMARK. In case E is a Banach space, this result is implicit in ([2], p. 88(4), Theorem 5.4); if in addition f is bounded this is proved in ([1], p. 88).

REFERENCES

1. Diestel, J. and J. J. Uhl Jr. Vector Measures, Amer. Math. Soc. Math. Surveys, Number 15(1977).
2. Edgar, E. Measurability in Banach Spaces, Indiana Univ. Math. J. 26(1977), 663-677.
3. Khurana, S. S. Strong Measurability in Frechet Spaces, Indian J. Pure Appl. Math. (to appear).
4. Schaefer, H. H. Topological Vector Spaces, Macmillan: New York, 1971.
5. Schwartz, L. Certaines Propriétés des Mesures sur les Espaces de Banach, Sem. Maurey-Schwartz, 1975, Ecole Polytechnique, No. 23.
6. Sion, S. On Analytic Sets in Topological Spaces, Trans. Amer. Math. Soc. 96(1960), 341-354.
7. Talagrand. Sur Une Conjecture de H. H. Corson, Bull. Sci. Math. (2), 99 (1975), 211-212.
8. Varadarajan, V. S. Measures on Topological Spaces, Amer. Math. Soc. Transl. 48(1965), 161-228.

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mssklai@cityu.edu.hk