

ON AN INVERSE TO THE HÖLDER INEQUALITY

J. PEČARIĆ

Faculty of Textile Technology
Zagreb University
Zagreb

C.E.M. PEARCE

Applied Mathematics Department
University of Adelaide
Adelaide 5005
South Australia

(Received October 5, 1994)

ABSTRACT. An extension is given for the inverse to Hölder's inequality obtained recently by Zhuang.

KEY WORDS AND PHRASES. Inverse Hölder inequality.

1991 AMS SUBJECT CLASSIFICATION CODE. 26D15.

Recently Zhuang [1] proved the following inverse of the arithmetico-geometric inequality.

THEOREM A. Let $0 < a \leq x \leq A$, $0 < b \leq y \leq B$, $1/p + 1/q = 1$, $p > 1$; then

$$\frac{x}{p} + \frac{y}{q} \leq \max \left\{ \frac{A/p + b/q}{A^{1/p}b^{1/q}}, \frac{a/p + B/q}{a^{1/p}B^{1/q}} \right\} x^{1/p}y^{1/q}, \quad (1)$$

or

$$x + y \leq \max \left\{ \frac{A + b}{A^{1/p}b^{1/q}}, \frac{a + B}{a^{1/p}B^{1/q}} \right\} x^{1/p}y^{1/q}, \quad (2)$$

the sign of equality in (1) and (2) holds if and only if either $(x, y) = (a, B)$ or $(x, y) = (A, b)$.

Moreover, if $a \geq B$, then

$$\frac{a/p + B/q}{a^{1/p}B^{1/q}} x^{1/p}y^{1/q} \leq \frac{x}{p} + \frac{y}{q} \leq \frac{A/p + b/q}{A^{1/p}b^{1/q}} x^{1/p}y^{1/q}, \quad (3)$$

the sign of equality on the right-hand side of (3) holds if and only if $(x, y) = (A, b)$, and the sign of equality on the left-hand side of (3) holds if and only if $(x, y) = (a, B)$. The sign of inequality in (3) is reversed if $b \geq A$.

This enables us to formulate the following theorem.

THEOREM 1. Suppose x, y, a, b, A, B, p, q are as in Theorem A and $\alpha, \beta > 0$. Then

$$\alpha x^p + \beta y^q \leq \max(C, D)xy, \quad (4)$$

where

$$C = (\alpha A^p + \beta b^q)/(Ab), \quad D = (\alpha a^p + \beta B^q)/(aB). \quad (5)$$

Equality occurs if and only if either $(x, y) = (a, B)$ or $(x, y) = (A, b)$. Moreover, if $\alpha p A^p \geq \beta q B^q$, then

$$Cxy \leq \alpha x^p + \beta y^q \leq Dxy, \quad (6)$$

with equality on the right-hand side if and only if $(x, y) = (A, b)$ and on the left if and only if $(x, y) = (a, B)$. The inequalities in (6) are reversed if $\alpha p A^p \leq \beta q B^q$.

PROOF. Inequalities (4) and (6) follow from (1) and (3) under the substitutions

$$x \rightarrow \alpha p x^p, \quad y \rightarrow \beta q y^q, \quad a \rightarrow \alpha p a^b, \quad b \rightarrow \beta q b^q, \quad A \rightarrow \alpha p A^p, \quad B \rightarrow \beta q B^q.$$

REMARK. Theorem 1 gives (1) and (2) together, (1) resulting from the substitutions $\alpha = 1/p$, $x \rightarrow x^{1/p}$, $A = A^{1/p}$, $a \rightarrow a^{1/p}$ and corresponding relations for β, y etc. with q in place of p , while (2) results from similar substitutions with $\alpha = 1 = \beta$.

The following result now gives an extension of the inverse to Hölder's inequality obtained in [1]. We suppose that all the integrals involved exist.

THEOREM 2. Let the functions f, g satisfy $0 < a \leq f(x) \leq A$, $0 < b \leq g(x) \leq B$ for almost all $x \in X$ with respect to a measure μ . Suppose $\alpha, \beta, p, q, C, D$ are as in Theorem 1. Then

$$\left(\int_X f^p d\mu\right)^{1/p} \left(\int_X g^q d\mu\right)^{1/q} \leq (\alpha\beta)^{-1/p} (\beta q)^{-1/q} \max(C, D) \int_X fg d\mu \tag{7}$$

and equality holds if and only if

$$\mu(E_1 \cup F_1) = \mu(X)$$

and

$$\mu(E_1) = \frac{(\alpha p A^p - \beta q b^q) \mu(X)}{\alpha p (A^p - a^p) + \beta q (B^q - b^q)},$$

where

$$E_1 = \{x \in X : f(x) = a, g(x) = B\},$$

$$F_1 = \{x \in X : f(x) = A, g(x) = b\}.$$

Moreover, if $\alpha p a^p \geq \beta q B^q$, then

$$\left(\int_X f^p d\mu\right)^{1/p} \left(\int_X g^q d\mu\right)^{1/q} \leq (\alpha p)^{-1/p} (\beta q)^{-1/q} D \int_X fg d\mu, \tag{8}$$

with equality only if $(f, g) = (a, B)$ a.e. on X and $\alpha p a^p = \beta q B^q$, and if $\alpha p A^p \leq \beta q b^q$, then

$$\left(\int_X f^p d\mu\right)^{1/p} \left(\int_X g^q d\mu\right)^{1/q} \leq (\alpha p)^{-1/p} (\beta q)^{-1/q} C \int_X fg d\mu, \tag{9}$$

with equality only if $(f, g) = (A, b)$ a.e. on X and $\alpha p A^p = \beta q b^q$.

PROOF. The first statement was proved in [1]. A simple proof of the remainder of the theorem was given for the case $\alpha = 1/p$, $\beta = 1/q$ in [2]. We give a similar simple proof for the general case.

$$\begin{aligned} \max(C, D) \int_X fg d\mu &= \int_X \max(C, D) fg d\mu \\ &\geq \int_X (\alpha f^p + \beta g^q) d\mu \\ &= \frac{1}{p} (\alpha p) \int_X f^p d\mu + \frac{1}{q} (\beta q) \int_X g^q d\mu \\ &\geq (\alpha p)^{1/p} (\beta q)^{1/q} \left(\int_X f^p d\mu\right)^{1/p} \left(\int_X g^q d\mu\right)^{1/q}, \end{aligned}$$

by the arithmetico-geometric inequality.

The equality conditions result from those in Theorem 1 and the arithmetico-geometric inequality.

Similarly we can prove (8). Using the second inequality in (6) we have

$$\begin{aligned} D \int_X fg \, d\mu &= \int_X Dfg \, d\mu \\ &\geq \int_X (\alpha f^p + \beta g^q) d\mu \\ &= \frac{1}{p} (\alpha p) \int_X f^p d\mu + \frac{1}{q} (\beta q) \int_X g^q d\mu \\ &\geq (\alpha p)^{1/p} (\beta q)^{1/q} \left(\int_X f^p d\mu \right)^{1/p} \left(\int_X g^q d\mu \right)^{1/q}. \end{aligned}$$

Relation (9) follows similarly.

REMARK. The simplest cases of (8) and (9) occur for $\alpha = 1/p, \beta = 1/q$. Then we have that if $a^p \geq B^q$, then

$$\left(\int_X f^p d\mu \right)^{1/p} \left(\int_X g^q d\mu \right)^{1/q} \leq D_1 \int_X fg \, d\mu$$

and if $A^p \leq b^q$, then

$$\left(\int_X f^p d\mu \right)^{1/p} \left(\int_X g^q d\mu \right)^{1/q} \leq C_1 \int_X fg \, d\mu,$$

where

$$D_1 = \left(\frac{1}{p} a^p + \frac{1}{q} B^q \right) / (aB),$$

$$C_1 = \left(\frac{1}{p} A^p + \frac{1}{q} b^q \right) / (Ab).$$

REFERENCES

1. ZHUANG, YA-DONG. On inverses of the Hölder inequality, J. Math. Anal. Applic., **161** (1991), 566-575.
2. MOND, B. and PEČARIĆ, J.E. Remark on a recent converse of Hölder's inequality, J. Math. Anal. Applic., **181** (1994), 280-281.

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru