

ON FUZZY SUBBUNDLES OF VECTOR BUNDLES

V. MURALI and G. LUBCZONOK

Received 27 March 2002

This paper considers fuzzy subbundles of a vector bundle. We define the operations sum, product, tensor product, Hom, and intersection of fuzzy subbundles and in each case, we characterize the corresponding flag of vector subbundles. We then propose two alternative definitions of integrability on fuzzy subbundles of a given type and discuss their naturality, merits, and shortcomings. We do these here with a view to introduce and study integrable fuzzy subbundles of tangent bundles on manifolds and foliations in further papers.

2000 Mathematics Subject Classification: 14F05, 14M15, 03E72, 14A20.

1. Introduction. A fuzzy subspace of a vector space can be characterized as a flag with weights, which are real numbers taken from the unit interval, attached at each component of the “nested” subspaces of the flag [7]. Similar ideas have been employed to study fuzzy subgroups of finite Abelian groups by Murali and Makamba [14]. A number of authors [1, 2, 5, 6, 8, 12] have considered fuzzy subspaces of vector spaces in other contexts. On the other hand, flags and flag manifolds are familiar objects of study for algebraists, geometers, topologists, and others [3, 4, 11, 13]. The present work was suggested by analogy with basic constructions of vector bundles. A fuzzy subbundle is a complex mathematical construction. It consists of a chain of subbundles and weights. Chains of subbundles or foliations are considered and investigated in foliation theory, for example, see [10]. In this context, we will have to deal with weights which are new components in the case of fuzzy subbundles. It is worthwhile to investigate the role and possible applications of weights. Therefore, a large part of this paper is devoted to the formulation and elucidation of definitions and their simple consequences. Further, we propose some integrability conditions on fuzzy subbundles, basically from an algebraic point of view. In so doing, we hope to lay the foundations for the study of foliations and other geometric aspects of fuzzy subbundles. We will omit the standard topological considerations in this paper, but concentrate largely on algebraic aspects.

2. Preliminaries

2.1. Definitions. In this paper, V denotes a vector space over the field of real numbers of dimension $n < \infty$. Further, suppose that the triple $\xi = (E, M, p)$ is a vector bundle, with $p : E \rightarrow M$ being the projection of the total space E onto

the base space M . For every $x \in M$, the fiber $E_x = p^{-1}(x)$ is a vector space over F isomorphic to V . For the purposes of formal constructions, it is not necessary to impose local triviality condition on the vector bundle, and as such, we have not assumed any such condition in general, unless otherwise stated in a particular context. We use $I = [0, 1]$, the real unit interval as a chain with the usual ordering in which \wedge stands for infimum (or intersection) and \vee stands for supremum (or union).

DEFINITION 2.1. A *fuzzy subbundle* of ξ is a fuzzy subset $\mu : E \rightarrow I$ such that the following holds. For $x \in M$, the restriction $\mu_x : E_x \rightarrow I$ of μ to E_x is a fuzzy subspace of E_x , which means that for every $x \in M$ and $u, v \in E_x$, $\mu_x(\alpha u + \beta v) \geq \mu_x(u) \wedge \mu_x(v)$, for $\alpha, \beta \in F$.

Without loss of generality, throughout this paper we tacitly assume that $\mu_x(0) = 1$ for every fuzzy subspace μ_x , $x \in M$. We will use the following form of Lowen's representation theorem [6].

THEOREM 2.2. A fuzzy subset μ of V is a fuzzy subspace of V if and only if there exist an integer $k \leq n$ and weights $1 = \alpha_0 > \alpha_1 > \dots > \alpha_k \geq 0$, and a strict flag $V^{\alpha_0} \subset V^{\alpha_1} \subset \dots \subset V^{\alpha_k} = V$ such that $\mu|V^{\alpha_j} \setminus V^{\alpha_{j-1}} = \alpha_j$ for $1 \leq j \leq k$.

Therefore, for every $x \in M$, we have a strict flag

$$V_x^{\alpha_0(x)} \subset V_x^{\alpha_1(x)} \subset \dots \subset V_x^{\alpha_k(x)} \subset E_x, \quad (2.1)$$

where k may depend on x and

$$1 = \alpha_0(x) > \alpha_1(x) > \dots > \alpha_k(x) \geq 0, \quad k \leq n, \quad (2.2)$$

are weights. Denote

$$C_x^{\alpha_j(x)} = V_x^{\alpha_j(x)} \setminus V_x^{\alpha_{j-1}(x)}, \quad j = 1, \dots, k, \quad x \in M. \quad (2.3)$$

Clearly,

$$C_x^{\alpha_j(x)} = \mu_x^{-1}(\alpha_j(x)), \quad j = 1, \dots, k, \quad x \in M. \quad (2.4)$$

REMARK 2.3. In this paper, we assume that the number of subspaces k in the flag at $x \in M$ is independent of x . Thus each weight $\alpha_j(x)$, $j = 0, 1, \dots, k$, defines a fuzzy subset of M .

From the definition of fuzzy subbundle and the Lowen's representation theorem as stated above, we have the following consequences, which we state as a proposition.

PROPOSITION 2.4. Suppose $\xi = (E, p, M)$ is a vector bundle. A fuzzy subset $\mu : E \rightarrow I$ is a fuzzy subbundle if and only if there exist

- (i) a flag on E_x for every $x \in M$, associated with μ_x ,

(ii) a finite chain of fuzzy subsets defined on M in the form of

$$1 = \alpha_0(x) > \alpha_1(x) > \dots > \alpha_k(x) \geq 0, \quad k \leq n, \quad x \in M. \quad (2.5)$$

In fuzzy set theory, it is not uncommon to discuss definitions in terms of the so-called levels. That is, the c -cut levels for $0 \leq c \leq 1$ of a fuzzy subbundle is a crisp subbundle. The following example demonstrates that if c -cuts are crisp subbundles, then the weights α 's are constants.

EXAMPLE 2.5. Let $\xi' = (E', p', M)$ be a subbundle of $\xi = (E, p, M)$. Thus $E'_x \subset E_x$ is a subspace, $\dim E'_x = m$, $m < n$. Consider a fuzzy subbundle, with the flag

$$\{0\} \subset E'_x \subset E_x \quad (2.6)$$

and the weights

$$1 > \alpha(x) > 0, \quad x \in M. \quad (2.7)$$

Thus, given $c \in I$, the c -cut consists of E'_x if $\alpha(x) \geq c$ and $\{0\}$ if $\alpha(x) < c$. We see that every c -cut of μ is a subbundle of ξ only if $\alpha(x)$ is a constant function on M .

For this reason, we proposed the more general definition as given in [Definition 2.1](#), from which the following is, however, immediate.

For every j , the set of all $v \in E$ such that $\mu(v) \geq \alpha_j(p(v))$ is a subbundle of ξ if $\dim V_x^{\alpha_j(x)}$ is constant for $x \in M$.

2.2. Morphisms. For the sake of completeness, we recall the following.

(i) If $\phi : V \rightarrow W$ is a linear map and if μ is a fuzzy subspace of V , then by the *image* of the fuzzy subspace μ , we mean the fuzzy subset $\phi(\mu)$ of W defined by $(\phi(\mu))(y) = \sup\{\mu(x) : \phi(x) = y\}$ if $\phi^{-1}(y) \neq \emptyset$, and $(\phi(\mu))(y) = 0$ if $\phi^{-1}(y) = \emptyset$. It is easy to check that $\phi(\mu)$ is a fuzzy subspace of W whenever μ is a fuzzy subspace of V .

(ii) A pair (ϕ, f) , where

$$\phi : E \rightarrow E', \quad f : M \rightarrow M', \quad (2.8)$$

is a *morphism* between two vector bundles $\xi = (E, p, M)$ and $\xi' = (E', p', M')$ if $f \circ p = p' \circ \phi$ and the restriction ϕ_x of ϕ to $E_x \rightarrow E'_{f(x)}$ is linear for every $x \in M$. For further results, we refer the reader to [\[4\]](#).

Given a fuzzy subbundle ν of ξ' , we can induce a fuzzy subbundle μ of ξ by the composition

$$\mu = \phi \circ \nu. \quad (2.9)$$

Clearly it is a pullback operation, and for every $x \in M$,

$$\mu_x = \phi_x \circ \nu_{f(x)}. \quad (2.10)$$

We can also define a push-forward operation (which is not possible for the case of crisp subbundles) on fuzzy subbundles as follows.

Let μ be a fuzzy subbundle of ξ . Denote by $\phi_x(\mu_x)$ the image of the fuzzy subspace μ_x . It is a fuzzy subspace of $E'_{f(x)}$ [8]. Given $y \in M'$, we define

$$\nu_y = \inf \{\phi_x(\mu_x) : x \in f^{-1}(y)\}. \quad (2.11)$$

Then the collection $\nu = \{\nu_y : y \in M'\}$ of fuzzy subspaces defines a fuzzy subbundle of ξ' . We call this subbundle ν the *image* of μ by the morphism (ϕ, f) .

2.3. Fuzzy subbundles as sections. We will give a description of fuzzy subbundle as a section of crisp bundle. Recall that E_x is vector space isomorphic to a given vector space V over a field F , for $x \in M$. Denote by $F(V)$ the set of all fuzzy subspaces of V . Then we construct the bundle $F(\xi)$ over M with the total space $F(E) = \bigcup\{F(E_x) : x \in M\}$ and the projection $F(p) : F(E) \rightarrow M$, where $F(p)(\mu) = x$ if $\mu \in F(E_x)$. Any section μ of $F(\xi)$ defines a fuzzy subbundle of ξ with $\mu_x = \mu(x) \in F(E_x)$, $x \in M$. This construction becomes more explicit for a weighted flag of a given type of a vector space of finite dimension over the field F , where F is the field of real or complex numbers. We will now explain the concept of “type” as follows.

Let $l = (l_1, l_2, \dots, l_k)$ be a finite sequence of positive integers such that $0 < l_1 < l_2 < \dots < l_k$. A fuzzy subspace μ is said to be of *type* l if its flag contains k subspaces V_1, V_2, \dots, V_k with $\dim V_p = l_p$, $p = 1, 2, \dots, k$. Denote by $F_l V(l)$ the set of all fuzzy subspaces of V of type l . We often identify type l of a flag simply with $l = (l_1, l_2, \dots, l_k)$ where the various l_i ’s are the dimensions of the various subspaces in V . If we identify V with F^n , then clearly the orthogonal group $\mathrm{SO}(n, \mathbb{R})$, $(\mathrm{SU}(n)$ in the case of complex vector space) acts transitively on the flags of the type l . Hence the set of all flags of type l is a homogeneous manifold denoted by $FL(l, F^n)$.

To discuss the weights attached to these subspaces of a flag associated with a fuzzy subspace μ , consider the open simplex in $(0, 1)^k$, denoted by Δ_k of all finite sequences $\alpha_1, \alpha_2, \dots, \alpha_k$ of real numbers of the form $1 > \alpha_1 > \alpha_2 > \dots > \alpha_k > 0$. With these notations, we can identify

$$FV(l) = FL(l, F^n) \times \Delta_k \quad (2.12)$$

as follows (see (2.1)). Every μ can be identified with

$$\{0\} \subset V_1^{\alpha_1} \subset \dots \subset V_k^{\alpha_k} \subset V, \quad (2.13)$$

which gives rise to a flag

$$\{0\} \subset V_1 \subset \cdots \subset V_k \subset V, \quad (2.14)$$

and the sequence $\alpha_1, \alpha_2, \dots, \alpha_k$ in

$$1 > \alpha_1 > \alpha_2 > \cdots > \alpha_k > 0, \quad (2.15)$$

and vice versa.

Let, as before, $\xi = (E, M, p)$ be a vector bundle.

Suppose

$$F_l(E) = \bigcup_{x \in M} FE_x(l), \quad (2.16)$$

where $FE_x(l)$ is the set of all fuzzy subspaces of E_x of type l . Now, we can construct a new fiber bundle $F\xi = (F_l(E), M, F(p))$ of fuzzy subspaces of a given type l and $F(p) : F_l(E) \rightarrow M$ is the fiber bundle projection. Then, we can regard any fuzzy subbundle of ξ as a section of the fiber bundle $F\xi$, that is, a mapping $\mu : M \rightarrow F_l(E)$ such that $\mu \circ F(p) = \text{identity on } M$.

Suppose now $\xi = (E, M, p)$ is locally trivial. Let $\phi_U : E|_U \rightarrow U \times F^n$ be a local trivialization of ξ , where $U \subset M$ is open, $E|_U = p^{-1}(U)$. Then $\mu_U = \mu \circ \phi_U^{-1}$ is a fuzzy subbundle of the trivial bundle $U \times F^n$. If $\phi_V : E|_V \rightarrow V \times F^n$ is another local trivialization of ξ , then $\phi_V \circ \phi_U^{-1}$ restricted to $U \cap V$ defines an isomorphism of the trivial bundle $(U \cap V) \times F^n$, and therefore

$$\phi_V \circ \phi_U^{-1}(x, v) = (x, \phi_{VU}(x)v), \quad x \in U \cap V, v \in F^n, \quad (2.17)$$

where $\phi_{VU} : U \cap V \rightarrow \text{GL}(F^n)$ is the transition function [4]. We have $\mu_U = \mu_V \circ \phi_{VU}$ on $U \cap V$. We note that if E is locally trivial, then so is $F\xi$. Given a local trivialization $E|_U \rightarrow U \times F^n$, we have the induced trivialization $F_l(E)|_U \rightarrow U \times FV(l)$, where $V = F^n$. The μ -section induces on every trivialization a corresponding section $\mu_U : U \rightarrow FV(l)$, where $V = F^n$. This point of view will be utilized later.

2.4. Continuity of fuzzy subbundles. In this section, we make a few observations about continuity of fuzzy subbundles. Given E_x , $x \in M$, $\mu_x : E_x \rightarrow I$ is a step function with finite number of values in I and is only right continuous. We want to define a continuous dependence of μ_x on $x \in M$. Suppose E, M are C^∞ manifolds and $p : E \rightarrow M$ is a C^∞ map. We consider the example of trivial bundle case in the next proposition.

PROPOSITION 2.6. *Suppose $p : U \times \mathbb{R}^n \rightarrow U$, where U is an open connected subset in \mathbb{R}^m and p is the trivial bundle projection and a fuzzy subbundle $\mu : U \times \mathbb{R}^n \rightarrow I$ is of a given type l . Given a fixed $v \in \mathbb{R}^n$, if $x \rightarrow \mu(x, v)$ is continuous on U , then the flag for μ is constant and the weights are continuous functions on U .*

Here we understand the flag for μ being constant in the sense that for every $x \in U$, $V_j^{\alpha_j(x)} = V_j$, $j = 1, 2, \dots, k$, where V_j is a fixed subspace of \mathbb{R}^n .

PROOF. Let $\alpha_1(x_0) < \alpha_2(x_0) < \dots < \alpha_k(x_0)$ be the weights at $x_0 \in U$. Consider a family of disjoint neighborhoods I_1, I_2, \dots, I_k of $\alpha_1(x_0), \alpha_2(x_0), \dots, \alpha_k(x_0)$, respectively. Given j , choose any $v_j \in \mathbb{R}^n$ such that

$$\mu(x_0, v_j) = \alpha_j(x_0), \quad \text{for } j = 1, 2, \dots, k. \quad (2.18)$$

By assumption,

$$\mu(x, v_j) \rightarrow \mu(x_0, v_j) = \alpha'_j(x_0), \quad \text{for } j = 1, 2, \dots, k, \text{ as } x \rightarrow x_0. \quad (2.19)$$

Hence there is a neighborhood N of x_0 such that $\mu(x, v_j) \in I_j$ for $j = 1, 2, \dots, k$. Since $\mu(x, \cdot)$ has exactly k values and I_j 's are disjoint, we must have

$$\mu(x, v_j) = \alpha_j(x), \quad \text{for } j = 1, 2, \dots, k. \quad (2.20)$$

This shows that $\alpha_j(x) \rightarrow \alpha_j(x_0)$ as $x \rightarrow x_0$. In other words, $\alpha_j(x)$ is continuous for $j = 1, 2, \dots, k$. Now we will demonstrate that

$$V_x^{\alpha_j(x)} = V_j \subset \mathbb{R}^n, \quad \text{for } j = 1, 2, \dots, k. \quad (2.21)$$

Consider a fixed point $x_0 \in U$ and choose a $v \in \mathbb{R}^n$. Then $\mu(x_0, v) = \alpha_i(x_0)$ for some $1 \leq i \leq k$. From the proof of continuity of the weights α 's, we conclude that there is a neighborhood N of x_0 in which $\mu(x, v) = \alpha_i(x)$. Now let $v \in V_{x_0}^{\alpha_j(x_0)}$, that is, $\mu(x_0, v) = \alpha_i(x_0)$ for some $i \geq j$. Therefore, $\mu(x, v) = \alpha_i(x) \geq \alpha_j(x)$. In other words, $v \in V_x^{\alpha_j(x)}$ in some neighborhood of x_0 . Since the role x and x_0 can be reversed, we have

$$V_x^{\alpha_j(x)} = V_{x_0}^{\alpha_j(x_0)}, \quad \text{for } x \in U \quad (2.22)$$

in some neighborhood of x_0 and $j = 1, 2, \dots, k$. It shows that the set of x satisfying (2.22) is open. It is trivial that this set is also closed from the assumption of continuity of μ and proven continuity of weights. Since U is connected,

$$V_x^{\alpha_j(x)} = V_{x_0}^{\alpha_j(x_0)} = V_j, \quad \forall x \in U, \quad j = 1, 2, \dots, k. \quad (2.23)$$

This completes the proof. \square

DEFINITION 2.7. A fuzzy subbundle of a given type $l = (l_1, l_2, \dots, l_k)$ is continuous or differentiable (C^∞) if the corresponding section $s_\mu : M \rightarrow F\xi$ is continuous or differentiable (C^∞).

Clearly this definition means the following.

- (i) Every distribution of the flag for μ is continuous, or differentiable (C^∞).
- (ii) Every weight is a continuous, or differentiable function (C^∞) on M .

3. Operations on fuzzy subbundles. All the operations on fuzzy subbundles are done pointwise, that is, we perform the corresponding operations on fuzzy subspaces of E_x at every $x \in M$. Thus the properties of the operations on fuzzy subbundles follow from the corresponding properties of operations on fuzzy subspaces [7]. We would like to point out that the resulting fuzzy subbundles in various cases may produce flags with the number of subspaces varying with x even in the cases of fuzzy subbundles of constant type. This anomaly exists even in the classical cases as most of the operations do not give rise to further subbundles although they do give subsheaves. The actual calculations of the number of weights and the weights themselves in various cases are given in [7].

3.1. Intersection, sum, and dual of fuzzy subbundles. Let μ and ν be fuzzy subbundles of a vector bundle $\xi = (E, p, M)$. Let

$$\begin{aligned} \{0_x\} &\subset V_x^{\alpha_1(x)} \subset \cdots \subset V_x^{\alpha_k(x)} \subset E_x, \\ \{0_x\} &\subset W_x^{\beta_1(x)} \subset \cdots \subset W_x^{\beta_l(x)} \subset E_x \end{aligned} \quad (3.1)$$

be the flags and weights of μ_x and ν_x , respectively, for a given $x \in M$.

Fuzzy intersection is given by the equation $(\mu \wedge \nu)_x = \mu_x \wedge \nu_x$, $x \in M$.

A number $\gamma^*(x) = \min\{\alpha_r(x), \beta_s(x)\}$ is a weight for $(\mu \wedge \nu)_x$ if

$$\begin{aligned} \alpha_{r-1}(x) &> \gamma^*(x) \geq \alpha_r(x), & \beta_{s-1}(x) &> \gamma^*(x) \geq \beta_s(x), \\ U^{\gamma^*(x)} &= V^{\alpha_r(x)} \cap W^{\beta_s(x)} \neq \{0_x\}, & x \in M. \end{aligned} \quad (3.2)$$

Fuzzy sum is defined by

$$(\mu + \nu)_x(\nu) = (\mu_x + \nu_x)(\nu) = \bigvee \{\mu_x(\nu - t) \wedge \nu_x(t) : t \in E_x\}. \quad (3.3)$$

A number $\gamma^*(x) = \min\{\alpha_r(x), \beta_s(x)\}$ is a weight for $(\mu + \nu)_x$ if

$$\alpha_{r-1}(x) > \gamma^*(x) \geq \alpha_r(x), \quad \beta_{s-1}(x) > \gamma^*(x) \geq \beta_s(x). \quad (3.4)$$

The corresponding subspace of E_x for $x \in M$ is

$$U^{\gamma^*(x)} = V^{\alpha_r(x)} + W^{\beta_s(x)}. \quad (3.5)$$

Dual subbundle μ^* is a fuzzy subbundle of the dual bundle ξ^* , the fiber at $x \in M$ is $(\mu^*)_x = (\mu_x)^*$, where for $f \in (E_x)^*$,

$$((\mu_x)^*)(f) = \begin{cases} 1, & f = 0, \\ 1 - \sup \{\mu_x(t) : t \in E_x, f(t) \neq 0\}, & f \neq 0. \end{cases} \quad (3.6)$$

Then the weights for μ^* are $\alpha_j^*(x) = 1 - \alpha_j(x)$, with the corresponding subspaces

$$V^{\alpha_j^*(x)} = (V^{\alpha_j(x)})^0 = \text{the annihilator of } V^{\alpha_j(x)}, \quad x \in M. \quad (3.7)$$

3.2. Whitney sum and tensor product. Let $\xi = (E, p, M)$ and $\xi' = (E', p', M)$ be two vector bundles over M . Let μ and ν be two fuzzy subbundles of ξ and ξ' , respectively.

Consider the Whitney sum $\xi \oplus \xi'$, where the total space of $\xi \oplus \xi'$ is the union of $E_x \oplus E'_x$, $x \in M$. The Whitney sum $\mu \oplus \nu$ is a fuzzy subbundle of $\xi \oplus \xi'$, with $(\mu \oplus \nu)_x = \mu_x \oplus \nu_x$, $x \in M$, where $\mu_x \oplus \nu_x$ is the fuzzy subspace of $E_x \oplus E'_x$ defined by $(\mu_x \oplus \nu_x)(t, t') = \mu_x(t) \wedge \nu_x(t')$ for $t \in E_x$, $t' \in E'_x$. Accordingly, the number $\gamma^*(x) = \min\{\alpha_r(x), \beta_s(x)\}$ is a weight for $(\mu \oplus \nu)_x$ if and only if the rectangle

$$[\alpha_r(x), \alpha_{r-1}(x)] \times [\beta_s(x), \beta_{s-1}(x)] \quad (3.8)$$

has nonempty intersection with the diagonal $I \times I$. The corresponding subspace is

$$V^{\gamma^*(x)} \oplus W^{\gamma^*(x)} \subset E_x \oplus E'_x. \quad (3.9)$$

Similarly, for the tensor product, $(\mu \otimes \nu)_x = \mu_x \otimes \nu_x$ is a fuzzy subspace of $E_x \otimes E'_x$, for $x \in M$ see [7, Section 4.2, page 204]. Accordingly, the number $\gamma^*(x) = \min\{\alpha_r(x), \beta_s(x)\}$ is a weight for $(\mu \otimes \nu)_x$ if and only if the rectangle

$$[\alpha_r(x), \alpha_{r-1}(x)] \times [\beta_s(x), \beta_{s-1}(x)] \quad (3.10)$$

has nonempty intersection with the diagonal $I \times I$.

3.3. Bundle of bilinear forms and Hom bundles. Let $\xi = (E, p, M)$ and $\xi' = (E', p', M)$ be two vector bundles over M . Let μ and ν be two fuzzy subbundles of ξ and ξ' , respectively. Consider the bundle $B(\xi, \xi') = (B(E, E'), \tilde{p}, M)$, where $B(E, E') = \bigcup(B(E_x, E'_x) : x \in M)$ with the fiber being the space of all bilinear maps from $E_x \times E'_x \rightarrow F$. Given two fuzzy subbundles μ , ν of ξ and ξ' , respectively, we define a fuzzy subbundle of $B(\xi, \xi')$ as follows:

$$(\mu \cdot \nu)_x = \mu_x \cdot \nu_x, \quad x \in M \quad (3.11)$$

as given in [7, Section 5.1, page 205]. The corresponding flag for $\mu_x \cdot \nu_x$ is given by

$$B^\alpha(E_x, E'_x) = \{\phi \in B(E_x, E'_x) : \phi(u, v) = 0 \text{ for } (u, v) \in E_x^{1-\alpha} \times E'_x^{1-\alpha}\}. \quad (3.12)$$

Similarly, for the bundle $\text{Hom}(\xi, \xi')$ with the fibers,

$$\text{Hom}_x(\xi, \xi') = \text{Hom}(E_x, E'_x), \quad x \in M. \quad (3.13)$$

We have $\text{Hom}(\mu, \nu)$ a fuzzy subbundle of $\text{Hom}(\xi, \xi')$ with the subspaces

$$(E_x^{1-\alpha})^\perp \otimes (E'_x)^\alpha, \quad x \in M. \quad (3.14)$$

Accordingly, a number $\gamma^*(x) = \min\{\alpha_r(x), \beta_s(x)\}$ is a weight for $\text{Hom}(\mu_x, \nu_x)$ if and only if the rectangle

$$[\alpha_r(x), \alpha_{r-1}(x)] \times [\beta_s(x), \beta_{s-1}(x)] \quad (3.15)$$

has nonempty intersection with the diagonal $I \times I$. Then

$$\text{Hom}^{\gamma^*}(\mu_x, \nu_x) = (E_x^{1-\alpha_r(x)})^\perp \otimes (E'_x)^{\beta_s(x)}, \quad x \in M. \quad (3.16)$$

4. Subbundles of tangent bundles. In this section, we consider fuzzy subbundles of tangent bundle on a manifold and we propose two different definitions of integrability of fuzzy subbundles. These two definitions apply to fuzzy subbundles μ of TM of a given type l , where TM is the tangent bundle of a manifold M .

4.1. Integrability I. We firstly recall that by a distribution of a tangent bundle TM of a manifold M , we mean a field of subspaces $\{D_x\}$, where $D_x \subset T_x M$ for $x \in M$ and $\text{Dim}(D_x) = k$, where k is independent of $x \in M$ [10]. It can be regarded as a section of the Grassmannian bundle $G_k(M)$, [10]. In terms of section of Grassmannian bundle, integrability means existence of local coordinates in which the section is constant. In the context of fuzzy subbundle of type l , it works as follows.

Let μ be a fuzzy subbundle of the tangent bundle $T = (TM, M, p)$ where $p : TM \rightarrow M$ is the bundle projection. It defines a section μ of the bundle $F(T) = \{FL(TM), M, F(p)\}$. Every local coordinate about $x \in M$ defines a trivialization of $TM|_U \rightarrow U \times \mathbb{R}^n$ and the corresponding trivialization of $FL(TM)|_U \rightarrow U \times F\mathbb{R}^n(l)$. This allows us to consider the induced section $\mu|_U : U \rightarrow F\mathbb{R}^n(l)$ as discussed in the end of [Section 2](#). We say that μ is *integrable* if about every $x \in M$ there is a neighborhood U of x and coordinates in U such that the induced section $\mu|_U$ is constant.

This definition, among other things, implies that

- (i) the flag for μ is integrable or all its distributions of subspaces are integrable,
- (ii) weights are constant on U .

We observe that the above definition, though a natural one, is too restrictive because we would like to have a definition of integrability that would be equally applicable to fuzzy subbundles with variable weights. This is the content of next two subsections. These concepts of integrability depend on what we mean by isomorphism of two fuzzy subspaces.

4.2. Integrability II. Let (U, μ) and (V, ν) be two fuzzy subspaces. We recall that a linear isomorphism of vector spaces $L : U \rightarrow V$ is an isomorphism of

fuzzy subspaces if

$$\nu(L(u)) = \mu(u), \quad u \in U. \quad (4.1)$$

Suppose μ is a fuzzy subbundle of the tangent bundle TM . A diffeomorphism $f : M \rightarrow M$ induces a diffeomorphism $df : TM \rightarrow TM$ through the differential df of f . We say that f preserves μ if

$$\mu(\xi) = \mu(df(\xi)), \quad \text{for } \xi \in TM. \quad (4.2)$$

Equivalently, if $x \in M$ and $\xi \in T_x M$, then

$$\mu_x(\xi) = \mu_{f(x)}(df(x)(\xi)), \quad (4.3)$$

which means that the differential

$$df(x) : T_x M \rightarrow T_{f(x)} M \quad (4.4)$$

is an isomorphism of fuzzy subspaces μ_x and $\mu_{f(x)}$. Further, we notice that

$$\alpha_j(x) = \alpha_j(f(x)), \quad \forall x \in M, \quad j = 1, 2, \dots, k, \quad (4.5)$$

$$df(x)(V_x^{\alpha_j(x)}) = V_{f(x)}^{\alpha_j(f(x))} = V_{f(x)}^{\alpha_j(x)} \quad (4.6)$$

since (4.5) is true.

Conversely, we have the following immediately straightforward result.

PROPOSITION 4.1. *A diffeomorphism $f : M \rightarrow M$ preserves a fuzzy subbundle $\mu : TM \rightarrow I$ if and only if f satisfies (4.5) and (4.6).*

Similarly, if X is a vector field on M and $\{\phi_t\}$, the corresponding flow for X , then we have the flow

$$\tilde{\phi}_t = d\phi_t : TM \rightarrow TM, \quad (4.7)$$

and we say X preserves μ if $\{\phi_t\}$ preserves μ for all $t \in \mathbb{R}$.

From (4.5) and (4.6),

$$\alpha_j(x) = \alpha_j(\phi_t(x)) \quad (4.8)$$

(or each weight α_j is a first integral of X),

$$\tilde{\phi}_t(V_x^{\alpha_j(x)}) = V_{\phi_t(x)}^{\alpha_j(x)}, \quad \forall t \in \mathbb{R}; \quad j = 1, 2, \dots, k. \quad (4.9)$$

We also have the following proposition.

PROPOSITION 4.2. *A vector field X on M preserves μ if and only if (4.8) and (4.9) are satisfied, or, equivalently, each weight α_j is a first integral of X and each distribution $\{V_x^{\alpha_j(x)}\}$, for $x \in M$, $j = 1, 2, \dots, k$, is preserved by X .*

According to foliation theory [10], a field X preserving a foliation is a foliated field.

Suppose a fuzzy subbundle $\mu : TM \rightarrow I$ is C^∞ (or smooth). We propose the following definition.

DEFINITION 4.3. The fuzzy subbundle μ is *integrable* if

- (i) $\{V_x^{\alpha_j(x)}\}$ is involutive (or a foliation) $x \in M$, $j = 1, 2, \dots, k$,
- (ii) the weight α_j is a basic function for the foliation $\{V_x^{\alpha_j(x)}\}$ $x \in M$.

Recall that $\alpha : M \rightarrow \mathbb{R}$ is a basic function for a foliation $\mathcal{F} = \{F_x\}$, $x \in M$, if for any vector field $X \in \mathcal{F}$ or $(X(x) \in F_x \ x \in M)$, $\mathcal{L}_X \alpha = 0$ or α is a first integral of X . It means that α is constant on the leaves of \mathcal{F} [10].

From this definition, we observe that, through $x \in M$ we have a chain of leaves

$$L_1 \subset L_2 \subset \dots \subset L_k \subset M, \quad (4.10)$$

where L_j is the leaf of the foliation $V_x^{\alpha_j(x)}$ and α_j is constant on L_j . We can use the μ function to express the above the fact.

REMARK 4.4. It is well known that there are foliations which do not admit any nonconstant basic function [10]. The definition II of integrability ties up closely the weights and the distribution of a fuzzy subspace. This can be relaxed if there is a need for it.

4.3. An extension to integrability II. Finally, we discuss a more general concept of isomorphism of fuzzy subspaces [9]. A possible third integrability condition (INT III) could be based on such an isomorphism (see [Remark 4.7](#) below), and integrability II will then be a special case of INT III.

Let (U, μ) and (V, ν) be two fuzzy subspaces. A pair (L, θ) , where $L : U \rightarrow V$ is an isomorphism and $\theta : I \rightarrow I$ is an increasing and onto function, is called an isomorphism of μ and ν if

$$\nu(L(u)) = \theta(\mu(u)), \quad u \in U. \quad (4.11)$$

Let $\{U^{\alpha_i}\}$, $i = 1, 2, \dots, k$, and $\{V^{\beta_j}\}$, $j = 1, 2, \dots, l$, be the two flags with weights corresponding μ and ν , respectively. From (4.11), we get

$$k = l, \quad (4.12a)$$

$$\theta(\alpha_i) = \beta_i, \quad i = 1, 2, \dots, k, \quad (4.12b)$$

$$L(U^{\alpha_i}) = V^{\beta_i}. \quad (4.12c)$$

We now apply this concept of isomorphism to fuzzy subbundle μ of TM . An *isomorphism* of μ is given by the following data: a pair (f, θ) of mappings,

such that $f : M \rightarrow M$ is a diffeomorphism and

$$\theta : M \times M \times I \rightarrow I \quad (4.13)$$

is a function which satisfies

- (i) $t \mapsto \theta(x, y, t)$ is increasing and onto, for each pair $x, y \in M$,
- (ii) $\mu(df(\xi)) = \theta(\pi(\xi), f(\pi(\xi)), \mu(\xi)), \xi \in TM$,

where $\pi : TM \rightarrow M$ is the bundle projection.

Equivalently, for any $x \in M$,

$$\mu_{f(x)}(df(x)(\xi)) = \theta(x, f(x), \mu_x(\xi)), \quad \xi \in T_x M. \quad (4.14)$$

Since we only work with fuzzy subbundles of a given type, we have from (4.12b) and (4.12c) of (4.12)

$$\begin{aligned} df(x)(U^{\alpha_j(x)}) &= U^{\alpha_j(f(x))}, \\ \alpha_j(f(x)) &= \theta(x, f(x), \alpha_j(x)), \quad j = 1, 2, \dots, k. \end{aligned} \quad (4.15)$$

Now, let X be a vector field on M and let $\{\phi_t\}$, $\{\tilde{\phi}_t\} = \{d\phi_t\}$ be the corresponding flows. Suppose θ is a function as in (4.13) which satisfies (i) and (ii).

DEFINITION 4.5. A vector field X on M preserves μ with respect to a given θ if and only if

$$\mu(\tilde{\phi}_t(\xi)) = \theta(\pi(\xi), \phi_t(\pi(\xi)), \mu(\xi)), \quad (4.16)$$

for all $t \in \mathbb{R}$, $\xi \in TM$, or

$$\mu_{\phi_t(x)}(\tilde{\phi}_t(x)(\xi)) = \theta(x, \phi_t(x), \mu_x(\xi)), \quad (4.17)$$

for all $t \in \mathbb{R}$, $\xi \in T_x M$ for each $x \in M$.

For the subspaces and weights, we have the following equations:

$$\tilde{\phi}_t(x)(U^{\alpha_j(x)}) = U^{\alpha_j(\phi_t(x))}, \quad (4.18)$$

$$\alpha_j(\phi_t(x)) = \theta(x, \phi_t(x), \alpha_j(x)), \quad x \in M, j = 1, 2, \dots, k. \quad (4.19)$$

We note that the choice of a connecting function θ can be done for every $\tilde{\phi}_t$, $t \in \mathbb{R}$. In the above, we assumed that θ is the same for all t . The second identity (4.19) implies (after passing to differentials)

$$\mathcal{L}_X \alpha_j(x) = \mathcal{L}_X^Y \theta(x, x, \alpha_j(x)), \quad j = 1, 2, \dots, k, \quad (4.20)$$

where $\mathcal{L}_X^Y \theta$ is the Lie derivative of θ on the second variable, $\theta(x, y, t)$.

Collecting the above equations, we have the following proposition in terms of the flow equations.

PROPOSITION 4.6. *A vector field X on M preserves μ with respect to θ if and only if the above equations (4.18) and (4.20) are satisfied.*

REMARK 4.7. We get the second integrability condition if $\theta(x, y, t) = t$ for all $x, y \in M$, $t \in I$. The problem of integrability involving nontrivial θ function where $\theta(x, y, t)$ is not necessarily equal to t for all $x, y \in M$, $t \in I$ will be fully discussed in an another paper. Further geometric integration theory and discussions on foliations will follow.

ACKNOWLEDGMENTS. Both authors were supported by the JRC of Rhodes University and the first author was supported by the FRD of South Africa. We gratefully acknowledge the support. We are very thankful and grateful to the anonymous referee for a number of valuable suggestions which improved the paper considerably.

REFERENCES

- [1] K. S. Abdukhalkov, *The dual of a fuzzy subspace*, Fuzzy Sets and Systems **82** (1996), no. 3, 375–381.
- [2] K. S. Abdukhalkov, M. S. Tulenbaev, and U. U. Umirbaev, *On fuzzy bases of vector spaces*, Fuzzy Sets and Systems **63** (1994), no. 2, 201–206.
- [3] R. Forman, *Hodge theory and spectral sequences*, Topology **33** (1994), no. 3, 591–611.
- [4] D. Husemoller, *Fibre Bundles*, McGraw-Hill, New York, 1966.
- [5] A. K. Katsaras and D. B. Liu, *Fuzzy vector spaces and fuzzy topological vector spaces*, J. Math. Anal. Appl. **58** (1977), no. 1, 135–146.
- [6] R. Lowen, *Convex fuzzy sets*, Fuzzy Sets and Systems **3** (1980), no. 3, 291–310.
- [7] G. Lubczonok and V. Murali, *On flags and fuzzy subspaces of vector spaces*, Fuzzy Sets and Systems **125** (2002), no. 2, 201–207.
- [8] P. Lubczonok, *Fuzzy vector spaces*, Fuzzy Sets and Systems **38** (1990), no. 3, 329–343.
- [9] B. B. Makamba, *Studies in fuzzy groups*, Ph.D. thesis, Rhodes University, South Africa, 1992.
- [10] P. Molino, *Riemannian Foliations*, Progress in Mathematics, vol. 73, Birkhäuser Boston, Massachusetts, 1988.
- [11] D. Monk, *The geometry of flag manifolds*, Proc. London Math. Soc. (3) **9** (1959), 253–286.
- [12] J. N. Mordeson, *Bases of fuzzy vector spaces*, Inform. Sci. **67** (1993), no. 1-2, 87–92.
- [13] T. Morimoto, *Geometric structures on filtered manifolds*, Hokkaido Math. J. **22** (1993), no. 3, 263–347.
- [14] V. Murali and B. B. Makamba, *On an equivalence of fuzzy subgroups. I*, Fuzzy Sets and Systems **123** (2001), no. 2, 259–264.

V. Murali: Department of Mathematics (Pure and Applied), Rhodes University, Grahamstown 6140, South Africa

E-mail address: v.murali@ru.ac.za

G. Lubczonok: Department of Mathematics (Pure and Applied), Rhodes University, Grahamstown 6140, South Africa

E-mail address: g.lubczonok@ru.ac.za

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

| | |
|------------------------|------------------|
| Manuscript Due | December 1, 2008 |
| First Round of Reviews | March 1, 2009 |
| Publication Date | June 1, 2009 |

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk