

## VARIATIONAL APPROACH TO DYNAMICS OF BRIGHT SOLITONS IN LOSSY OPTICAL FIBERS

M. F. MAHMOOD and S. BROOKS

Received 2 January 2002

A variational analysis of dynamics of soliton solution of coupled nonlinear Schrödinger equations with oscillating terms is made, considering a birefringent fiber with a third-order nonlinearity in the anomalous dispersion frequency region. This theoretical model predicts optical soliton oscillations in lossy fibers.

2000 Mathematics Subject Classification: 34A25, 35A15, 35Q55, 37K40, 78A60.

**1. Introduction.** The propagation of bright solitons in birefringent optical fibers has been the subject of intensive theoretical and experimental investigations during the past two decades. The solitons are nonlinear pulses in fibers when the nonlinearity induced by the optical intensity balances the dispersion of the fiber. Studies of bright soliton propagation in fibers are demanding with reference to the development of soliton-based optical communication, generation of short pulses, and soliton lasers. The idea of exploiting these solitons as natural bits to transmit optical data motivates important research efforts towards the development of models [2, 3, 5, 6, 7, 8, 9] describing solitary wave propagation in optical fibers under different conditions. In lossless, one would not expect solitons to distort in either the time or frequency domains regardless of the distance over which they propagate. This supposition is, however, not true in the case of lossy fibers. In this paper, we follow an adiabatic approach using a variational technique [1] to study dynamics of bright solitons generated from semiconductor lasers in a lossy birefringent fiber.

**2. Variational approach to coupled nonlinear Schrödinger (CNLS) equations.** The birefringence in fibers arises from the geometric and material contributions [4]. The geometric contribution comes from the ellipticity of the core of the fiber which breaks the cylindrical symmetry. The material contribution comes from the strain within the material forming the core and cladding of the fiber. The birefringence in fibers gives rise to two orthogonal polarization modes that need to be considered. The dynamics of optical solitons in a lossy birefringent fiber is important from a theoretical point of view as well as for

the applications, and it is governed by the following CNLS equations [5, 6, 7]:

$$\begin{aligned}
 & i(u_z + \delta u_t) + \frac{1}{2} u_{tt} + (|u|^2 + B|v|^2)u + Av^2 u^* \exp(-iR\delta z) \\
 & + Du^2 v^* \exp\left(\frac{1}{2}iR\delta z\right) + D(2|u|^2 + |v|^2)v \exp\left(-\frac{1}{2}iR\delta z\right) \\
 & = -i\gamma u, \\
 & i(v_z - \delta v_t) + \frac{1}{2} v_{tt} + (B|u|^2 + |v|^2)v + Au^2 v^* \exp(iR\delta z) \\
 & + Dv^2 u^* \exp\left(-\frac{1}{2}iR\delta z\right) + D(|u|^2 + 2|v|^2)u \exp\left(\frac{1}{2}iR\delta z\right) \\
 & = -i\gamma v,
 \end{aligned} \tag{2.1}$$

where

$$B = \frac{2(1 + \varepsilon \sin^2 \theta)}{2 + \varepsilon \cos^2 \theta}, \quad A = \frac{\varepsilon \cos^2 \theta}{2 + \varepsilon \cos^2 \theta}, \quad D = \frac{\varepsilon}{2} \tan \theta \quad \text{with } \theta = 35^\circ; \tag{2.2}$$

$u$  and  $v$  are the normalized amplitudes of the fast and slow modes, respectively;  $t$  is the time coordinate in a frame moving with the average group velocity of the two modes measured in units of the modulational wavelength;  $z$  is the distance along the propagation direction;  $\delta$  is the normalized birefringence;  $\varepsilon$  denotes the relative strength of the cross-phase modulation;  $R$  is the wave vector mismatch due to modal birefringence of the fiber; and  $\gamma$  denotes the losses in the fiber. The oscillating terms in (2.1) arise from nonlinear polarization and cannot be taken off in the case of fibers with low birefringence as it causes an instability in which the slow moving partial pulse transfers energy to the fast moving partial pulse [7]. Using the transformations (see [5, 6])  $p = \sqrt{B/2}(ue^{i\alpha z} + ve^{-i\alpha z})$  and  $q = \sqrt{B/2}(ue^{i\alpha z} - ve^{-i\alpha z})$ , we can write the CNLS equations (2.1) as

$$\begin{aligned}
 & i(p_z + \delta q_t) + \frac{1}{2} p_{tt} + \alpha q + p(f|p|^2 + h|q|^2) + gp^*q^2 + i\gamma p = 0, \\
 & i(q_z + \delta p_t) + \frac{1}{2} q_{tt} + \alpha p + q(f'|q|^2 + h|p|^2) + gq^*p^2 + i\gamma q = 0,
 \end{aligned} \tag{2.3}$$

where  $\alpha = (1/4)R\delta$ ,  $f = (1/2B)(1 + A + B + 4D)$ ,  $f' = (1/2B)(1 + A + B - 4D)$ ,  $h = (1/B)(1 - A)$ , and  $g = (1/2B)(1 + A - B)$ .

Since losses in the fiber lead to exponential decrease of soliton amplitude, we use the transformations  $p \rightarrow p'e^{-\gamma z}$  and  $q \rightarrow q'e^{-\gamma z}$  to write (2.3) in the form

$$\begin{aligned}
 & i(p'_z + \delta q'_t) + \alpha q' + \frac{1}{2} p'_{tt} + p'(f'|p'|^2 + h|q'|^2)e^{-2\gamma z} + gp'^*q'^2e^{-2\gamma z} = 0, \\
 & i(q'_z + \delta p'_t) + \alpha p' + \frac{1}{2} q'_{tt} + q'(f'|q'|^2 + h|p'|^2)e^{-2\gamma z} + gq'^*p'^2e^{-2\gamma z} = 0.
 \end{aligned} \tag{2.4}$$

The spatiotemporal evolution of the wave amplitudes in the case of bright solitons is governed by the Lagrangian density

$$\begin{aligned}
L = \int_{-\infty}^{\infty} & \left[ \frac{i}{2} (P_z P'^* - p_z'^* p') + \frac{i}{2} (q_z q'^* - q_z'^* q') + \frac{i\delta}{2} (p_t q'^* - q_t'^* p') \right. \\
& + \frac{i\delta}{2} (q_t p'^* - p_t'^* q') + \alpha (p'^* q' + p' q'^*) + h |p'|^2 |q'|^2 e^{-2yz} \\
& + \frac{1}{2} (f |p'|^4 e^{-2yz} - |p_t'|^2) + \frac{1}{2} (f' |q'|^4 e^{-2yz} - |q_t'|^2) \\
& \left. + \frac{1}{2} g (p'^2 q'^* + q'^2 p'^*) e^{-2yz} \right] dt,
\end{aligned} \tag{2.5}$$

where  $p'_z = \partial p'/\partial z$ ,  $p'_t = \partial p'/\partial t$ , and so on.

The bright soliton solutions of (2.4) using a variational technique is based upon assuming the trial function [6]

$$\begin{aligned}
\begin{bmatrix} p' \\ q' \end{bmatrix} = & 2\eta_r \exp \left[ 2iV_r(t - \zeta_r) - iC \tanh [2\eta_r(t - \zeta_r)](t - \zeta_r)^2 + iD_r \right] \\
& \times \operatorname{sech} [2\eta_r(t - \zeta_r)],
\end{aligned} \tag{2.6}$$

that describes the temporal form of the soliton pulses. The evolution parameters  $\eta_r$ ,  $\zeta_r$ ,  $V_r$ ,  $D_r$  ( $r = 1, 2$  correspond to  $p'$  and  $q'$  solitons, respectively), and  $C$  represent amplitude, central position, velocity of soliton's central position as it propagates along the fibre, phase, and initial frequency chirp of the soliton, respectively. We substitute (2.6) into the Lagrangian density (2.5) and use Euler-Lagrange equations to obtain the following system of coupled ordinary differential equations (ODEs) for the evolution of soliton parameters:

$$\frac{d}{dz} (\eta_r V_r) = \frac{1}{8} \frac{\partial L_1}{\partial \zeta_r} + \frac{1}{8} \frac{\partial L_2}{\partial \zeta_r} + \frac{1}{8} \frac{\partial L_3}{\partial \zeta_r}, \tag{2.7}$$

$$8\eta_r \frac{d\zeta_r}{dz} = 16\eta_r V_r - 2C\alpha_1 - 4C - \frac{\partial L_1}{\partial V_r} - \frac{\partial L_3}{\partial V_r}, \tag{2.8}$$

$$4 \frac{d\eta_r}{dz} = (-1)^r \frac{\partial L_1}{\partial \Lambda} + (-1)^r \frac{\partial L_3}{\partial \Lambda}, \tag{2.9}$$

$$\begin{aligned}
\frac{dD_1}{dz} = & 2V_1 \frac{d\zeta_1}{dz} + 4f\eta_1^2 e^{-2yz} - 2V_1^2 - 2\eta_1^2 + \frac{C^2}{16\eta_1^2} \alpha_2 \\
& + \frac{C^2}{4\eta_1^2} \alpha_4 + \frac{C^2}{4\eta_1^2} \alpha_5 + \frac{1}{4} \frac{\partial L_1}{\partial \eta_1} + \frac{1}{4} \frac{\partial L_2}{\partial \eta_1} + \frac{1}{4} \frac{\partial L_3}{\partial \eta_1},
\end{aligned} \tag{2.10}$$

$$\begin{aligned}
\frac{dD_2}{dz} = & 2V_2 \frac{d\zeta_2}{dz} + 4f'\eta_2^2 e^{-2yz} - 2V_2^2 - 2\eta_2^2 + \frac{C^2}{16\eta_2^2} \alpha_2 \\
& + \frac{C^2}{4\eta_2^2} \alpha_4 + \frac{C^2}{4\eta_2^2} \alpha_5 + \frac{1}{4} \frac{\partial L_1}{\partial \eta_2} + \frac{1}{4} \frac{\partial L_2}{\partial \eta_2} + \frac{1}{4} \frac{\partial L_3}{\partial \eta_2},
\end{aligned} \tag{2.11}$$

where

$$\begin{aligned}
 \alpha_1 &= \left( \frac{\pi^2}{9} - \frac{2}{3} \right), & \alpha_2 &= \left( \frac{7\pi^4}{225} - \frac{\pi^2}{3} + \frac{2}{5} \right), \\
 \alpha_3 &= 1, & \alpha_4 &= \left( \frac{\pi^2}{18} + \frac{2}{3} \right), & \alpha_5 &= \left( \frac{\pi^2}{12} - \frac{1}{2} \right), \\
 L_1 &= 4\alpha\eta_2 \int_{-\infty}^{\infty} \operatorname{sech} x_1 \operatorname{sech} x_2 \cos \Lambda dx_1 = \frac{8\alpha\eta\rho \cos \Lambda}{\sinh \rho}, \\
 L_2 &= 8h\eta_1\eta_2^2 e^{-2yz} \int_{-\infty}^{\infty} \operatorname{sech}^2 x_1 \operatorname{sech}^2 (x_1 + \rho) dx_1 \\
 &= 32h\eta^3 e^{-2yz} \frac{\cosh \rho}{\sinh \rho} (\rho - \tanh \rho), \\
 L_3 &= 8g\eta_1\eta_2^2 e^{-2yz} \int_{-\infty}^{\infty} \operatorname{sech}^2 x_1 \operatorname{sech}^2 (x_1 + \rho) \cos 2\Lambda dx_1 \\
 &= 32g\eta^3 e^{-2yz} \cos \Lambda \frac{\cosh \rho}{\sinh^3 \rho} (\rho - \tanh \rho), \\
 \Lambda &= 2t(V_1 - V_2) - 2(V_1\zeta_1 - V_2\zeta_2) - (D_2 - D_1) \\
 &\quad - C \tanh [2\eta_1(t - \zeta_1)](t - \zeta_1)^2 + C \tanh [2\eta_2(t - \zeta_2)](t - \zeta_2)^2.
 \end{aligned} \tag{2.12}$$

We evaluate the above integrals for nearly equal pulse amplitudes  $\eta_1 \approx \eta_2 \approx \eta$ , relative phase  $\phi = D_2 - D_1$ , relative distance between two polarization maxima  $\rho = x_2 - x_1$ , and  $V_1 \approx V_2 \approx V$ . The relative parameters  $\eta_{12}$ ,  $V_{12}$ ,  $\phi$ , and  $\rho$  defined for  $p'$  and  $q'$  solitons are obtained as follows. Writing  $\eta_{12} = \eta_1 - \eta_2$  and using (2.9), we get

$$\begin{aligned}
 \frac{d\eta_{12}}{dz} &= -4\alpha\eta \left( \frac{\rho}{\sinh \rho} \right) \sin \left( \frac{V}{\eta} \rho + \phi \right) \\
 &\quad - 32g\eta^3 e^{-2yz} \left[ \frac{\cosh \rho}{\sinh^3 \rho} (\rho - \tanh \rho) \right] \sin 2 \left( \frac{V}{\eta} \rho + \phi \right).
 \end{aligned} \tag{2.13}$$

Similarly, writing  $V_{12} = V_1 - V_2$  and using (2.7), we get

$$\begin{aligned}
 \frac{dV_{12}}{dz} &= -\frac{4}{3}\alpha\eta\rho \cos \left( \frac{V}{\eta} \rho + \phi \right) - \frac{64}{15}h\eta^3 \rho e^{-2yz} \\
 &\quad - \frac{64}{15}g\eta^3 \rho e^{-2yz} \cos 2 \left( \frac{V}{\eta} \rho + \phi \right).
 \end{aligned} \tag{2.14}$$

Also, from  $\rho = 2\eta(\zeta_1 - \zeta_2)$  and by using (2.8), (2.13), and (2.14), we obtain

$$\begin{aligned}
 \frac{d^2\rho}{dz^2} &= -\frac{16}{3}\alpha\eta^2 \rho \cos \left( \frac{V}{\eta} \rho + \phi \right) - \frac{256}{15}h\eta^4 \rho e^{-2yz} \\
 &\quad - \frac{256}{15}g\eta^4 \rho e^{-2yz} \cos 2 \left( \frac{V}{\eta} \rho + \phi \right) \\
 &\quad - 2 \left( \frac{\pi^2}{9} + \frac{4}{3} \right) C \alpha \sin \left( \frac{V}{\eta} \rho + \phi \right) \\
 &\quad - \frac{16}{3} \left( \frac{\pi^2}{9} + \frac{4}{3} \right) C g \eta^2 e^{-2yz} \sin 2 \left( \frac{V}{\eta} \rho + \phi \right).
 \end{aligned} \tag{2.15}$$

The two solitons with opposite phases form a bound state provided that

$$\frac{V}{\eta} \rho + \phi = \Phi = \pm \pi. \quad (2.16)$$

Thus, we write (2.15) as

$$\frac{d^2\rho}{dz^2} = \frac{16}{3} \alpha \eta^2 \rho - \frac{256}{15} h \eta^4 \rho e^{-2yz} - \frac{256}{15} g \eta^4 \rho e^{-2yz} \quad (2.17)$$

or

$$\frac{d^2\rho}{dz^2} + (ae^{-2yz} - b)\rho(z) = 0, \quad (2.18)$$

where  $a = (256/15)\eta^4(h+g)$  and  $b = (16/3)\alpha\eta^2$ .

Equation (2.18) can also be written as

$$\sigma^2 \frac{d^2\rho}{d\sigma^2} + \sigma \frac{d\rho}{d\sigma} + \left( \sigma^2 - \frac{b}{\gamma^2} \right) \rho = 0, \quad (2.19)$$

where  $\sigma = (\sqrt{a}/\gamma)e^{-yz}$ .

Equation (2.19) is a Bessel equation. Its general solution is given by

$$\rho(\sigma) = XJ \frac{\sqrt{b}}{\gamma}(\sigma) + YJ - \frac{\sqrt{b}}{\gamma}(\sigma). \quad (2.20)$$

Considering  $\gamma$  to be small, we use asymptotic expansion of Bessel function to write (2.20) as

$$\rho(\sigma) \approx \left( \frac{2\gamma}{\pi\sqrt{a}e^{-yz}} \right)^{1/2} \cos \left( \frac{\sqrt{a}}{\gamma} (1 - \gamma z + o(z^2)) - \frac{\pi}{2} \frac{\sqrt{b}}{\gamma} - \frac{\pi}{4} \right), \quad (2.21)$$

and the frequency  $\omega$  of relative oscillations of soliton positions is given by

$$\omega^2 = a = \frac{256}{15} \eta^4 \left( \frac{2}{B} - 1 \right). \quad (2.22)$$

**3. Conclusion.** In this paper, we have considered CNLS equations with oscillating terms to develop a theoretical model of a birefringent optical fiber. This theoretical model demonstrates polarized bright soliton dynamics in a lossy birefringent fiber. We used a variational approach to obtain frequency of relative oscillations of soliton positions by taking into account the interaction between different polarizations in a lossy birefringent optical fiber.

## REFERENCES

- [1] D. Anderson, M. Lisak, and T. Reichel, *Approximate analytical approaches to nonlinear pulse propagation in optical fibers: a comparison*, Phys. Rev. A **38** (1988), 1618–1622.
- [2] K. J. Blow, N. J. Doran, and D. Wood, *Polarization instabilities for solitons in birefringent fibers*, Opt. Lett. **12** (1987), no. 3, 202–204.

- [3] M. N. Islam, C. D. Poole, and J. P. Gordon, *Soliton trapping in birefringent optical fibers*, Opt. Lett. **14** (1989), no. 18, 1011–1013.
- [4] I. P. Kaminow, *Polarization in optical fibers*, IEEE J. Quant. Electron. **17** (1981), no. 1, 15–22.
- [5] M. F. Mahmood, W. W. Zachary, and T. L. Gill, *Chirped solitary pulses in low birefringent optical fibers: a theoretical model*, Opt. Quant. Electron. **28** (1996), 1007–1012.
- [6] ———, *Nonlinear pulse propagation in elliptically birefringent optical fibers*, Phys. D **90** (1996), no. 3, 271–279.
- [7] C. R. Menyuk, *Pulse propagation in an elliptically birefringent Kerr medium*, IEEE J. Quant. Electron. **25** (1989), no. 12, 2674–2682.
- [8] J. M. Soto-Crespo, N. Akhmediev, and A. Ankiewicz, *Soliton propagation in optical devices with two-component fields: a comparative study*, J. Opt. Soc. Amer. B Opt. Phys. **12** (1995), no. 6, 1100–1109.
- [9] S. Trillo, S. Wabnitz, W. C. Banyai, N. Finlayson, C. T. Seaton, G. I. Stegeman, and R. H. Stolen, *Observation of ultrafast nonlinear polarization switching induced by polarization instability in a birefringent fiber rocking filter*, IEEE J. Quant. Electron. **25** (1989), no. 1, 104–112.

M. F. Mahmood: Department of Mathematics, Howard University, Washington, DC 20059, USA

*E-mail address:* [mmahmood@howard.edu](mailto:mmahmood@howard.edu)

S. Brooks: Department of Mathematics, Howard University, Washington, DC 20059, USA

## Special Issue on Intelligent Computational Methods for Financial Engineering

### Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Lean Yu**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [yulean@amss.ac.cn](mailto:yulean@amss.ac.cn)

**Shouyang Wang**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; [sywang@amss.ac.cn](mailto:sywang@amss.ac.cn)

**K. K. Lai**, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [mskklai@cityu.edu.hk](mailto:mskklai@cityu.edu.hk)