

## THE ROOTS OF THE THIRD JACKSON $q$ -BESSEL FUNCTION

L. D. ABREU, J. BUSTOZ, and J. L. CARDOSO

Received 16 July 2002

We derive analytic bounds for the zeros of the third Jackson  $q$ -Bessel function  $J_v^{(3)}(z; q)$ .

2000 Mathematics Subject Classification: 33D15, 33D67.

**1. Introduction.** There are three known  $q$ -analogs of classical Bessel functions [6, 5] that are due to Jackson [7]. Following the notation of Ismail [6, 5], these are designated by  $J_v^{(k)}(z; q)$ ,  $k = 1, 2, 3$ .

The parameter  $q$  is taken to satisfy  $0 < q < 1$ . The third Jackson  $q$ -Bessel function  $J_v^{(3)}(z; q)$  is defined as

$$J_v^{(3)}(z; q) := \frac{(q^{v+1}; q)_\infty}{(q; q)_\infty} z^v {}_1\Phi_1 \left[ \begin{matrix} 0 \\ q^{v+1} \end{matrix} ; q, qz^2 \right]. \quad (1.1)$$

This function is also known as the Hahn-Exton  $q$ -Bessel function [8, 9]. The notation  ${}_1\Phi_1$  in (1.1) is the standard in use for  $q$ -hypergeometric series [4]. The function  $J_v^{(3)}(z; q)$  satisfies a linear  $q$ -difference equation and it is known that  $J_v^{(3)}(z; q)$  has an infinite number of simple real zeros [8]. In this paper, we will give lower and upper bounds for these zeros. The roots of these functions are of interest for several reasons. Firstly, it is intrinsically interesting to provide information about the roots of a function such as (1.1), which is an entire function of order zero. Also, the roots of  $J_v^{(2)}(z; q)$  and  $J_v^{(3)}(z; q)$  figure prominently in expansions in terms of “ $q$ -Fourier series” [2, 3]. Lastly, if we denote the roots of  $J_v^{(3)}(z; q)$  by  $j_{n,v}^{(3)}$ , then the mass points of the orthogonality measure for a  $q$ -analog of Lommel polynomials are located at the points  $1/j_{n,v}^{(3)}$ . Furthermore, although the function defined in (1.1) is of a simpler character than the remaining Jackson  $q$ -Bessel functions, it is hoped that the results given here for  $J_v^{(3)}(z; q)$  may be extended in the future to  $J_v^{(k)}(z; q)$ ,  $k = 1, 2$ .

**2. The roots of  $J_v^{(3)}(z; q)$ .** We prove two lemmas stating the existence of an odd number of roots in a certain interval and then we prove that  $J_v^{(3)}(z; q)$  has only one root in such an interval.

First we apply the following transformation to (1.1):

$$(c; q)_{\infty} \Phi_1 \left[ \begin{array}{c} 0 \\ c \end{array} ; q, z \right] = (z; q)_{\infty} \Phi_1 \left[ \begin{array}{c} 0 \\ z \end{array} ; q, c \right]. \quad (2.1)$$

This produces

$$J_v^{(3)}(z; q) = \frac{(qz^2; q)_{\infty}}{(q; q)_{\infty}} z^v \Phi_1 \left[ \begin{array}{c} 0 \\ qz^2 \end{array} ; q, q^{v+1} \right]. \quad (2.2)$$

This last relation in a series form gives

$$\begin{aligned} J_v^{(3)}(z; q) &= \frac{z^v}{(q; q)_{\infty}} \sum_{k=0}^{\infty} (-1)^k \frac{(qz^2; q)_{\infty}}{(qz^2; q)_k (q; q)_k} q^{k(k+2v+1)/2} \\ &= \frac{z^v}{(q; q)_{\infty}} \sum_{k=0}^{\infty} (-1)^k \frac{(q^{k+1} z^2; q)_{\infty}}{(q; q)_k} q^{k(k+2v+1)/2}. \end{aligned} \quad (2.3)$$

This representation will be critical in the proof of the next two lemmas.

**LEMMA 2.1.** *If  $q^{v+1} < (1-q)^2$ , then  $\text{sgn}[J_v^{(3)}(q^{-m/2}; q)] = (-1)^m$ ,  $m = 1, 2, \dots$*

**PROOF.** Set  $z = q^{-m/2}$  in (2.3) to obtain

$$\frac{(q; q)_{\infty}}{q^{-mv/2}} J_v^{(3)}(q^{-m/2}; q) = \sum_{k=0}^{\infty} (-1)^k \frac{(q^{-m+k+1}; q)_{\infty}}{(q; q)_k} q^{k(k+2v+1)/2}. \quad (2.4)$$

Now observing that  $(q^{-m+k+1}; q)_{\infty} = 0$  if  $k < m$ , the series on the right-hand side of this last equality can be written as

$$\sum_{k=m}^{\infty} (-1)^k \frac{(q^{-m+k+1}; q)_{\infty}}{(q; q)_k} q^{k(k+2v+1)/2}. \quad (2.5)$$

Setting  $j = k - m$  in this last series yields

$$\frac{(q; q)_m}{q^{-mv/2}} J_v^{(3)}(q^{-m/2}; q) = (-1)^m \sum_{j=0}^{\infty} (-1)^j A_j, \quad (2.6)$$

where

$$A_j = \frac{(q^{j+1}; q)_{\infty}}{(q; q)_{j+m}} q^{(j+m)(j+m+2v+1)/2}. \quad (2.7)$$

Now we prove that  $A_{j+1} < A_j$ . A calculation shows that  $A_{j+1} < A_j$  is equivalent to  $q^{m+j+v+1} < (1-q^{m+j+1})(1-q^{j+1})$ . But the left-hand side of this inequality is decreasing in  $m$  and  $j$ , while the right-hand side is increasing in  $m$  and  $j$ . So we only need to verify the case  $j = m = 0$ , that is,  $q^{v+1} < (1-q)^2$ , but

this is the hypothesis of the lemma. Clearly, since  $A_{j+1} < A_j$ , then

$$\operatorname{sgn}(-1)^m \sum_{j=0}^{\infty} (-1)^j A_j = (-1)^m. \quad (2.8) \quad \square$$

**Lemma 2.1** states that there exist an odd number of roots in the interval  $(q^{-m/2+1/2}, q^{-m/2})$ . The next lemma refines this statement.

**LEMMA 2.2.** *Let  $q^{\nu+1} < (1-q)^2$  and define*

$$\alpha_m^{(\nu)}(q) = \frac{\log(1-q^{m+\nu}/(1-q^m))}{\log q}. \quad (2.9)$$

*Then*

$$\operatorname{sgn}\left[J_{\nu}^{(3)}\left(q^{-m/2+\alpha_m^{(\nu)}(q)/2}; q\right)\right] = (-1)^{m-1}, \quad m = 1, 2, \dots \quad (2.10)$$

**PROOF.** First observe that the function  $\alpha_m^{(\nu)}(q)$  is well defined because if  $q^{\nu+1} < (1-q)^2$ , then  $q^{\nu+1} < (1-q)$  and so, for positive integer  $m$ ,  $q^{m+\nu} < (1-q^m)$ , that is,  $1-q^{m+\nu}/(1-q^m) > 0$ . Being defined, it is clear that  $\alpha_m^{(\nu)}(q)$  is positive because  $1-q^{m+\nu}/(1-q^m) < 1$  holds for any  $q \in (0, 1)$ .

Observe also that

$$\alpha_m^{(\nu)}(q) < 1 \iff \log\left(1 - \frac{q^{m+\nu}}{1-q^m}\right) > \log q \iff q^{m+\nu} < (1-q)(1-q^m), \quad (2.11)$$

which is true if  $q^{\nu+1} < (1-q)^2$ . So, we have  $0 < \alpha_m^{(\nu)}(q) < 1$ .

Now, set  $z = q^{-m/2+\alpha_m^{(\nu)}(q)/2}$  in (2.3). The substitution gives

$$\begin{aligned} & \frac{(q;q)_{\infty}}{q^{-m\nu/2+\nu\alpha_m^{(\nu)}(q)/2}} J_{\nu}^{(3)}\left(q^{-m/2+\alpha_m^{(\nu)}(q)/2}; q\right) \\ &= \sum_{k=0}^{m-2} (-1)^k \frac{(q^{-m+\alpha_m^{(\nu)}(q)+k+1}; q)_{\infty}}{(q;q)_k} q^{k(k+2\nu+1)/2} \\ & \quad + \sum_{k=m-1}^{\infty} (-1)^k \frac{(q^{-m+\alpha_m^{(\nu)}(q)+k+1}; q)_{\infty}}{(q;q)_k} q^{k(k+2\nu+1)/2}. \end{aligned} \quad (2.12)$$

Denote the first sum above by  $S_1$  and the second by  $S_2$ . If  $0 \leq k \leq m-2$ , then  $0 < \alpha_m^{(\nu)}(q) < 1$  implies that  $\operatorname{sgn}(q^{-m+\alpha_m^{(\nu)}(q)+k+1}; q)_{\infty} = (-1)^{m-k-1}$ . Thus  $\operatorname{sgn} S_1 = (-1)^{m-1}$ ,  $m = 1, 2, \dots$

In  $S_2$ , set  $j = k - m + 1$  to obtain

$$S_2 = (-1)^{m-1} \sum_{j=0}^{\infty} (-1)^j A_j, \quad (2.13)$$

where

$$A_j = \frac{(q^{\alpha_m^{(\nu)}(q)+j}; q)_\infty}{(q; q)_{j+m-1}} q^{(j+m-1)(j+m+2\nu)/2}. \quad (2.14)$$

A calculation shows that  $A_{j+1} < A_j$  is reduced to  $q^{j+m+\nu} < (1 - q^{\alpha_m^{(\nu)}(q)+j})(1 - q^{m+j})$ , which holds because  $q^{m+\nu} = (1 - q^{\alpha_m^{(\nu)}(q)})(1 - q^m)$  and because the left-hand side of the last inequality is decreasing in  $j$  and the right-hand side is increasing in  $j$ . The infinite series is thus positive and therefore

$$\operatorname{sgn} S_2 = (-1)^{m-1} = \operatorname{sgn} S_1. \quad (2.15)$$

□

From Lemmas 2.1 and 2.2, we know that  $J_\nu^{(3)}(z; q)$  has an odd number of roots in the interval  $(q^{-m/2+\alpha_m^{(\nu)}(q)}, q^{-m/2})$ . The next theorem proves that there is exactly one root in each such interval and that there are no other roots.

**THEOREM 2.3.** *If  $q^{\nu+1} < (1 - q)^2$  and if  $w_k^{(\nu)}(q)$  are the positive roots of  $J_\nu^{(3)}(z; q)$ , ordered increasingly in  $k$ , then  $w_k^{(\nu)}(q) = q^{-k/2+\epsilon_k(\nu)}$ , with  $0 < \epsilon_k(\nu) < \alpha_k^{(\nu)}(q)$ ,  $k = 1, 2, \dots$ .*

**PROOF.** From the preceding lemmas, we know that  $J_\nu^{(3)}(z; q)$  has roots of the form  $w_k^{(\nu)} = q^{-k/2+\epsilon_k}$ , with  $0 < \epsilon_k < \alpha_k^{(\nu)}(q)$ .

To simplify the notation, we set

$$F(z) = \frac{(q; q)_\infty}{(q^{\nu+1}; q)_\infty z^\nu} J_\nu^{(3)}(z; q). \quad (2.16)$$

We prove that the only positive roots of  $F(z)$  inside the disk  $|z| < q^{-m/2}$  are  $w_k^{(\nu)}(q)$ ,  $k = 1, 2, \dots, m$ .

Suppose there are other roots  $\pm\lambda_k$ ,  $k = 1, 2, \dots, P_m$ ;  $\lambda_k > 0$ . By Jensen's theorem [1], we can write

$$\begin{aligned} & \frac{1}{2\pi} \int_0^{2\pi} \log |F(q^{-m/2} e^{i\theta})| d\theta \\ &= 2 \sum_{k=1}^m \log \frac{q^{-m/2}}{w_k} + 2 \sum_{k=1}^{P_m} \log \frac{q^{-m/2}}{\lambda_k} \\ &= 2 \sum_{k=1}^m \log q^{-m/2+k/2-\epsilon_k} + 2 \sum_{k=1}^{P_m} \log \frac{q^{-m/2}}{\lambda_k} \\ &= \frac{-m^2+m}{2} \log q - 2 \log q \sum_{k=1}^m \epsilon_k + 2 \sum_{k=1}^{P_m} \log \frac{q^{-m/2}}{\lambda_k}. \end{aligned} \quad (2.17)$$

On the other hand, by the definition of the  $q$ -Bessel function, we have

$$\begin{aligned}
 F(q^{-m/2}e^{i\theta}) &= \sum_{k=0}^{\infty} (-1)^k \frac{q^{k(k+1)/2-mk}}{(q^{\nu+1};q)_k (q;q)_k} e^{2ik\theta} \\
 &= (-1)^m \frac{q^{(-m^2+m)/2}}{(q^{\nu+1};q)_m (q;q)_m} e^{2im\theta} \\
 &\quad \times \sum_{k=-m}^{\infty} (-1)^k \frac{q^{k(k+1)/2}}{(q^{m+\nu+1};q)_k (q^{m+1};q)_k} e^{2ik\theta}.
 \end{aligned} \tag{2.18}$$

Then we have

$$\begin{aligned}
 \log |F(q^{-m/2}e^{i\theta})| &= \frac{-m^2+m}{2} \log q - \log (q^{\nu+1};q)_m - \log (q;q)_m \\
 &\quad + \log \left| \sum_{k=-m}^{\infty} (-1)^k \frac{q^{k(k+1)/2}}{(q^{m+\nu+1};q)_k (q^{m+1};q)_k} e^{2ik\theta} \right|
 \end{aligned} \tag{2.19}$$

so that

$$\begin{aligned}
 &\lim_{m \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \log |F(q^{-m/2}e^{i\theta})| d\theta \\
 &= \lim_{m \rightarrow \infty} \frac{-m^2+m}{2} \log q - \log (q^{\nu+1};q)_{\infty} - \log (q;q)_{\infty} \\
 &\quad + \lim_{m \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \log \left| \sum_{k=-m}^{\infty} (-1)^k \frac{q^{k(k+1)/2}}{(q^{m+\nu+1};q)_k (q^{m+1};q)_k} e^{2ik\theta} \right| d\theta.
 \end{aligned} \tag{2.20}$$

Now observe that

$$\lim_{m \rightarrow \infty} \sum_{k=-m}^{\infty} (-1)^k \frac{q^{k(k+1)/2}}{(q^{m+\nu+1};q)_k (q^{m+1};q)_k} e^{2ik\theta} = \sum_{k=-\infty}^{\infty} (-1)^k q^{k(k+1)/2} e^{2ik\theta}. \tag{2.21}$$

The above limit is uniform in  $\theta$ . By the Jacobi triple product identity [4],

$$\sum_{k=-\infty}^{\infty} (-q^{1/2}e^{2i\theta})^k (q^{1/2})^{k^2} = (q;qe^{2i\theta};e^{-2i\theta};q)_{\infty}. \tag{2.22}$$

Using the uniform convergence to interchange the limit and the integral,

$$\begin{aligned}
& \lim_{m \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \log \left| \sum_{k=-m}^{\infty} (-1)^k \frac{q^{k(k+1)/2}}{(q^{m+\nu+1};q)_k (q^{m+1};q)_k} e^{2ik\theta} \right| d\theta \\
&= \frac{1}{2\pi} \int_0^{2\pi} \log |(q;qe^{2i\theta};e^{-2i\theta};q)_\infty| d\theta \\
&= \log(q;q)_\infty + \frac{1}{2\pi} \int_0^{2\pi} \log |(qe^{2i\theta};q)_\infty| d\theta \\
&\quad + \frac{1}{2\pi} \int_0^{2\pi} \log |(e^{-2i\theta};q)_\infty| d\theta \\
&= \log(q;q)_\infty + \sum_{j=0}^{\infty} \frac{1}{2\pi} \int_0^{2\pi} \log |(1-q^{j+1}e^{2i\theta})_\infty| d\theta \\
&\quad + \sum_{j=0}^{\infty} \frac{1}{2\pi} \int_0^{2\pi} \log |(1-q^je^{-2i\theta})_\infty| d\theta \\
&= \log(q;q)_\infty.
\end{aligned} \tag{2.23}$$

The integrals in the third equality above vanish because of the mean value theorem for harmonic functions.

We have thus concluded that

$$\begin{aligned}
& \lim_{m \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \log |F(q^{-m/2}e^{i\theta})| d\theta \\
&= \lim_{m \rightarrow \infty} \frac{-m^2 + m}{2} \log q - \log(q^{\nu+1};q)_\infty.
\end{aligned} \tag{2.24}$$

From (2.17), we can write

$$\lim_{m \rightarrow \infty} \left( 2 \sum_{k=1}^{P_m} \log \frac{q^{-m/2}}{\lambda_k} \right) - \lim_{m \rightarrow \infty} \left( 2 \log q \sum_{k=1}^m \epsilon_k \right) = -\log(q^{\nu+1};q)_\infty. \tag{2.25}$$

But, as can be seen by the Taylor expansion of  $\alpha_k^{(\nu)}(q)$ ,  $\epsilon_k = O(q^k)$ , and this implies  $\sum_{k=1}^{\infty} \epsilon_k < \infty$ .

Also

$$\sum_{k=1}^{P_m} \log \frac{q^{-m/2}}{\lambda_k} > \log \frac{q^{-m/2}}{\lambda_1} \rightarrow \infty \quad \text{as } m \rightarrow \infty. \tag{2.26}$$

So the identity (2.25) can only hold if the first sum is empty, and the only roots are thus  $\pm w_k^{(\nu)}(q)$ ,  $k = 1, 2, \dots$ .  $\square$

**REMARK 2.4.** It follows from (2.25) that  $\sum_{k=0}^{\infty} \epsilon_k = \log(q^{\nu+1}; q)_{\infty} / 2 \log q$ .

**REMARK 2.5.** Observe that for fixed  $j$ , we can always choose  $m$  sufficiently large so that

$$\begin{aligned} q^{m+j+\nu+1} &< (1-q^{m+j+1})(1-q^{j+1}), \\ q^{j+m+\nu} &< (1-q^{\alpha_m^{\nu}(q)+j})(1-q^{m+j}). \end{aligned} \quad (2.27)$$

Thus, we have the asymptotic behaviour

$$w_k \sim q^{-m/2} \quad \text{when } m \rightarrow \infty \quad (2.28)$$

without the restriction  $q^{\nu+1} < (1-q)^2$ . In [5], Ismail conjectured that

- (1)  $\lim_{m \rightarrow \infty} q^{m/2} w_m^{(\nu)}(q) = 1$ ,
- (2)  $\lim_{m \rightarrow \infty} w_{m+1}^{(\nu)}(q^2) / w_m^{(\nu)}(q^2) = 1/q$ .

The asymptotic relation (2.28) establishes these conjectures.

**REMARK 2.6.** Lemmas 2.1 and 2.2 and Theorem 2.3 state that the roots  $w_k^{(\nu)}(q)$  satisfy the inequalities

$$q^{-m/2+\alpha_m^{(\nu)}(q)} < w_k^{(\nu)}(q) < q^{-m/2}. \quad (2.29)$$

These bounds are quite accurate. This is evident if we estimate the length of the interval containing the roots. A somewhat tedious calculation with Taylor series shows that

$$q^{-m/2} - q^{-m/2+\alpha_m^{(\nu)}(q)} = q^{m/2+\nu} O(1). \quad (2.30)$$

Clearly, for fixed  $q$  satisfying the conditions of the theorem, the bounds become increasingly accurate as either  $k$  or  $\nu$  increases.

## REFERENCES

- [1] L. V. Ahlfors, *Complex Analysis*, 3rd ed., McGraw-Hill, New York, 1979.
- [2] J. Bustoz and J. L. Cardoso, *Basic analog of Fourier series on a  $q$ -linear grid*, J. Approx. Theory **112** (2001), no. 1, 134–157.
- [3] J. Bustoz and S. K. Suslov, *Basic analog of Fourier series on a  $q$ -quadratic grid*, Methods Appl. Anal. **5** (1998), no. 1, 1–38.
- [4] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Encyclopedia of Mathematics and Its Applications, vol. 35, Cambridge University Press, Cambridge, 1990.
- [5] M. E. H. Ismail, *Some properties of Jackson's third  $q$ -Bessel function*, to appear.
- [6] ———, *The zeros of basic Bessel functions, the functions  $J_{\nu+ax}(x)$ , and associated orthogonal polynomials*, J. Math. Anal. Appl. **86** (1982), no. 1, 1–19.
- [7] F. H. Jackson, *On generalized functions of Legendre and Bessel*, Trans. Roy. Soc. Edin. **41** (1904), 1–28.
- [8] H. T. Koelink and R. F. Swarttouw, *On the zeros of the Hahn-Exton  $q$ -Bessel function and associated  $q$ -Lommel polynomials*, J. Math. Anal. Appl. **186** (1994), no. 3, 690–710.

[9] R. F. Swarttouw, *The Hahn-Exton  $q$ -Bessel function*, Ph.D. thesis, Technische Universiteit Delft, Delft, the Netherlands, 1992.

L. D. Abreu: Department of Mathematics, University of Coimbra, Coimbra 3000, Portugal

*E-mail address:* [daniel@mat.uc.pt](mailto:daniel@mat.uc.pt)

J. Bustoz: Department of Mathematics, Arizona State University, Tempe, AZ 85287-1804, USA

*E-mail address:* [bustoz@asu.edu](mailto:bustoz@asu.edu)

J. L. Cardoso: Department of Mathematics, University of Tras-os-Montes and Alto Douro, Vila Real 5000-410, Portugal

*E-mail address:* [jluis@utad.pt](mailto:jluis@utad.pt)

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)