

TWISTINGS, CROSSED COPRODUCTS, AND HOPF-GALOIS COEXTENSIONS

S. CAENEPEEL, DINGGUO WANG, and YANXIN WANG

Received 1 December 2002

Let H be a Hopf algebra. Ju and Cai (2000) introduced the notion of twisting of an H -module coalgebra. In this paper, we study the relationship between twistings, crossed coproducts, and Hopf-Galois coextensions. In particular, we show that a twisting of an H -Galois coextension remains H -Galois if the twisting is invertible.

2000 Mathematics Subject Classification: 16W30.

1. Introduction. A fundamental result in Hopf-Galois theory is the normal basis theorem, stating that, for a finitely generated cocommutative Hopf algebra H over a commutative ring k , the set of isomorphism classes of Galois H -objects that are isomorphic to H as an H -comodule is a group, and this group is isomorphic to the second Sweedler cohomology group $H^2(H, k)$ (see [14]). The Galois object corresponding to a 2-cocycle is then given by a crossed product construction. The crossed product construction can be generalized to arbitrary Hopf algebras and plays a fundamental role in the theory of extensions of Hopf algebras (see [3, 11]). Also in this more general situation, it turns out that there is a close relationship between crossed products on one side and Hopf-Galois extensions and cleft extensions on the other side (cf. [3, 4, 11]). A survey can be found in [13]. An alternative way to deform the multiplication on an H -comodule algebra A has been proposed in [1], using the so-called twisting of A , and it was shown that the crossed product construction can be viewed as a special case of the twisting construction. The relation between twistings and H -Galois extensions was studied in [2].

Now, there exists a coalgebra version of the normal basis theorem (see [6]). In this situation, one tries to deform the comultiplication on a commutative Hopf algebra H , using this time a Harrison cocycle instead of a Sweedler cocycle. Crossed coproducts, cleft coextensions, and Hopf-Galois coextensions have been introduced and studied in [8, 10]. Ju and Cai [12] have introduced the notion of twisting of an H -module coalgebra, which can be viewed as a dual version of the twistings introduced in [1]. The aim of this paper is to study the relationship between twistings, crossed coproducts, and Hopf-Galois coextensions. Our main result is the fact that the twisting of a Hopf-Galois coextension by an invertible twist map is again a Hopf-Galois coextension (and vice versa).

Our paper is set up as follows. In [Section 2.1](#), we recall the twistings introduced in [\[12\]](#), and in [Section 2.2](#) we recall the definition of a Harrison cocycle and the crossed coproduct construction from [\[8, 10\]](#). In [Section 3](#), we introduce an alternative version of 2-cocycles, called twisted 2-cocycles, and discuss the relation with Harrison cocycles ([Proposition 3.3](#)). In [Section 4](#), we introduce an equivalence relation on the set of twistings of an H -module coalgebra and we show that a twisting in an equivalence class is invertible if and only if all the other twistings in this equivalence class are invertible ([Theorem 4.4](#)). Two twistings are equivalent if and only if their corresponding crossed coproducts are isomorphic ([Proposition 2.1](#)). In [Section 5](#), the relationship between twistings and Hopf-Galois coextensions is investigated.

For the general theory of Hopf algebras, we refer to the literature (see, e.g, [\[9, 13, 15\]](#)).

2. Notation and preliminary results. We work over a field k . All maps are assumed to be k -linear. For the comultiplication on a k -coalgebra C , we use the Sweedler-Heyneman notation

$$\Delta_C(c) = c_1 \otimes c_2 \quad (2.1)$$

with the summation implicitly understood. We use a similar notation for a (right) coaction of a coalgebra on a comodule:

$$\rho(m) = m_0 \otimes m_1 \in M \otimes C. \quad (2.2)$$

Let A be a k -algebra, then $\text{Hom}(C, A)$ is also an algebra, with convolution product

$$(f * g)(c) = f(c_1)g(c_2). \quad (2.3)$$

We will denote by $\text{Reg}(C, A)$ the set of convolution invertible elements in $\text{Hom}(C, A)$, and \mathcal{M}_A^C will be the category of modules with a right A -action and a right C -coaction, such that the C -coaction is A -linear.

2.1. Twistings of a coalgebra. We recall some definitions and results from [\[12\]](#). Let H be a Hopf algebra over a field k , with bijective antipode S . The composition inverse of the antipode will be denoted by \bar{S} .

Recall that a right H -module coalgebra is a coalgebra C which is also a right H -module such that

$$\Delta(c \cdot h) = c_1 \cdot h_1 \otimes c_2 \cdot h_2, \quad \varepsilon_C(c \cdot h) = \varepsilon_C(c)\varepsilon_H(h) \quad (2.4)$$

for all $c \in C$ and $h \in H$.

Consider the category \mathcal{M}_H^C , whose objects are right H -modules and right C -comodules M such that the following compatibility relation is satisfied:

$$\rho(m \cdot h) = m_0 \cdot h_1 \otimes m_1 \cdot h_2. \quad (2.5)$$

Recall from [12] that we have the following associative multiplication on $\text{Hom}(C, H \otimes C)$:

$$\tau * \lambda = (m_H \otimes \text{id}_C) \circ (\text{id}_H \otimes \lambda) \circ \tau \quad (2.6)$$

for all $\tau, \lambda \in \text{Hom}(C, H \otimes C)$. The unit of this multiplication is the map $\sigma : C \rightarrow H \otimes C$, $\sigma(c) = 1 \otimes c$.

Remark that we have an algebra isomorphism

$$\alpha : \text{Hom}(C, H \otimes C) \longrightarrow_H \text{End}(H \otimes C)^{\text{op}}, \quad (2.7)$$

where α is defined as follows. Take $\tau : C \rightarrow H \otimes C$, and write $\tau(c) = c_{-1} \otimes c_0$ (summation is understood). Then $\alpha(\tau) = f_\tau : H \otimes C \rightarrow H \otimes C$ is given by

$$f_\tau(h \otimes c) = h\tau(c) = hc_{-1} \otimes c_0. \quad (2.8)$$

Assume that τ satisfies the following normality conditions:

$$(1 \otimes \varepsilon_C)\tau(c) = \varepsilon(c)1_H, \quad (\varepsilon_H \otimes 1)\tau(c) = c \quad (2.9)$$

or

$$c_{-1}\varepsilon_C(c_0) = \varepsilon_C(c)1_H, \quad \varepsilon_H(c_{-1})c_0 = c. \quad (2.10)$$

We can then define a new (in general noncoassociative) comultiplication Δ_τ on C as follows:

$$\Delta_\tau(c) = c_1 \cdot c_{2,-1} \otimes c_{2,0} \quad \text{or} \quad \Delta_\tau = (\psi_H \otimes \text{id}) \circ (\text{id} \otimes \tau) \circ \Delta, \quad (2.11)$$

where $\psi_H : C \otimes H \rightarrow C$ is the right H -action on C , and where we used the Sweedler-type notation $(\text{id} \otimes \tau)(\Delta c) = c_1 \otimes c_{2,-1} \otimes c_{2,0}$. Let C^τ be equal to C as a right H -module, with comultiplication $\Delta_\tau(c)$. A similar construction applies to $M \in \mathcal{M}_H^C : F_\tau(M) = M^\tau$ as a right H -comodule, with

$$\rho^\tau(m) = m_0 \cdot m_{1,-1} \otimes m_{1,0}, \quad (2.12)$$

where τ is called a twisting if and only if C^τ is a right H -module coalgebra, and $M^\tau \in \mathcal{M}_H^{C^\tau}$ for all $M \in \mathcal{M}_H^C$. It is shown in [12, Theorem 1.1] that $\tau : C \rightarrow H \otimes C$,

satisfying (2.9), is a twisting if and only if for all $h \in H$ and $c \in C$,

$$c_{-1}h_1 \otimes c_0 \cdot h_2 = h_1(c \cdot h_2)_{-1} \otimes (c \cdot h_2)_0, \quad (2.13)$$

$$c_{-1} \otimes c_{0,1} \cdot c_{0,2,-1} \otimes c_{0,2,0} = c_{1,-1}c_{2,-1,1} \otimes c_{1,0} \cdot c_{2,-1,2} \otimes c_{2,0}, \quad (2.14)$$

where (2.13) is equivalent to

$$S(h_1)c_{-1}h_2 \otimes c_0 \cdot h_3 = (c \cdot h)_{-1} \otimes (c \cdot h)_0. \quad (2.15)$$

If τ has an inverse λ , then the functor F is an equivalence of categories.

Left-hand twistings are defined in a similar way. Consider the vector space isomorphism

$$\text{Hom}(C, C \otimes H) \cong \text{End}_H(C \otimes H^{\text{op}}, C \otimes H^{\text{op}}). \quad (2.16)$$

The composition on the right-hand side is transported into the following associative multiplication on $\text{Hom}(C, C \otimes H)$:

$$\tau \times \lambda = T \circ (T \circ \lambda * T \circ \tau). \quad (2.17)$$

Here, T is the usual twist map. The unit σ' on $\text{Hom}(C, C \otimes H)$ is given by $\sigma'(c) = c \otimes 1$. If $\lambda \in \text{Hom}(C, C \otimes H)$ satisfies the normalizing conditions

$$(1 \otimes \varepsilon_H)\lambda(c) = c, \quad (\varepsilon_C \otimes 1)\lambda(c) = \varepsilon_C(c)1_H, \quad (2.18)$$

then we can twist the comultiplication on C as follows. Write $\lambda(c) = c_0 \otimes c_1$, and define ${}_{\lambda}\Delta$ by

$${}_{\lambda}\Delta(c) = c_{1,0} \otimes c_2 \cdot c_{1,1}. \quad (2.19)$$

Let ${}^{\lambda}C$ be equal to C as a right H -module, equipped with the comultiplication ${}_{\lambda}\Delta$. The C -coaction $M \in {}^C\mathcal{M}_H$ can also be twisted as follows:

$${}^{\lambda}\rho(m) = m_{-1,0} \otimes m_0 m_{-1,1}, \quad (2.20)$$

where λ is called a left-hand twisting if ${}^{\lambda}C$ is an H -module coalgebra, and ${}^{\lambda}M \in {}^{\lambda}C\mathcal{M}_H$ for every $M \in {}^C\mathcal{M}_H$. The map $\lambda : C \rightarrow C \otimes H$ satisfying (2.18) is a left-hand twisting if and only if for all $h \in H$ and $c \in C$,

$$c_0 \cdot h_1 \otimes c_1 h_2 = (c \cdot h_1)_0 \otimes h_2(c \cdot h_1)_1, \quad (2.21)$$

$$c_{0,1,0} \otimes c_{0,2} \cdot c_{0,1,1} \otimes c_1 = c_{1,0} \otimes c_{2,0} \cdot c_{1,1,1} \otimes c_{2,1}c_{1,1,2}. \quad (2.22)$$

Equation (2.21) is equivalent to

$$c_0 \cdot h_1 \otimes \bar{S}(h_3)c_1h_2 = \sum (c \cdot h)_0 \otimes (c \cdot h)_1. \quad (2.23)$$

For $\tau \in \text{Hom}(C, H \otimes C)$ with inverse λ , we write

$$\tau(c) = c_{-1} \otimes c_0, \quad \lambda(c) = c_{(-1)} \otimes c_{(0)}. \quad (2.24)$$

We then have

$$c_{-1}c_{0,(-1)} \otimes c_{0,(0)} = c_{(-1)}c_{(0),-1} \otimes c_{(0),0} = 1 \otimes c. \quad (2.25)$$

For $\gamma \in \text{Hom}(C, C \otimes H)$ with inverse μ , we write

$$\gamma(c) = c_0 \otimes c_1, \quad \mu(c) = c_{(0)} \otimes c_{(1)}. \quad (2.26)$$

Let $\mathcal{T}(C)$ and $\mathcal{L}(C)$ be the sets of twistings and left-hand twistings of C , respectively, and $U(\mathcal{T}(C))$ and $U(\mathcal{L}(C))$ the sets of invertible twistings and left-hand twistings, respectively.

PROPOSITION 2.1. *Take $\tau \in U(\mathcal{T}(C))$ with inverse λ . Define $\ell(\tau) : C \rightarrow C \otimes H$ by*

$$\ell(\tau)(c) = c_{0,(0)} \cdot \bar{S}(c_{0,(-1)})\bar{S}(c_{-1})_1 \otimes \bar{S}(c_{-1})_2. \quad (2.27)$$

Take $\gamma \in U(\mathcal{L}(C))$, with inverse μ . Define $r(\gamma) : C \rightarrow H \otimes C$ by

$$r(\gamma)(c) = S(c_1)_1 \otimes c_{0,(0)} \cdot S(c_{0,(1)})S(c_1)_2. \quad (2.28)$$

Then $\ell : U(\mathcal{T}(C)) \rightarrow U(\mathcal{L}(C))$ is a bijection with inverse r . Furthermore, $\ell(\sigma) = \sigma'$ and $r(\sigma') = \sigma$.

PROOF. It is shown in [12] that $\ell(\tau) \in U(\mathcal{L}(C))$ with sn inverse given by

$$\ell(\tau)'(c) = c_{0,(0)} \cdot \bar{S}(c_{0,(-1)})_1 \bar{S}(c_{-1})_1 \otimes \bar{S}(\bar{S}(c_{-1})_3)\bar{S}(c_{0,(-1)})_2 \bar{S}(c_{-1})_2. \quad (2.29)$$

Set $g = \bar{S}(c_{0,(-1)})$, $h = \bar{S}(c_{-1})$. Then $\ell(\tau)(c) = c_{0,(0)} \cdot gh_1 \otimes h_2$, so

$$\begin{aligned}
r(\ell(\tau))(c) &= S(h_2)_1 \otimes (c_{0,(0)} \cdot gh_1)_{0,(0)} \cdot \bar{S}((c_{0,(0)} \cdot gh_1)_{0,(-1)})_1 \\
&\quad \cdot \bar{S}((c_{0,(0)} \cdot gh_1)_{-1})_1 S(\bar{S}(\bar{S}((c_{0,(0)} \cdot gh_1)_{-1})_3) \\
&\quad \cdot \bar{S}((c_{0,(0)} \cdot gh_1)_{0,(-1)})_2 \bar{S}((c_{0,(0)} \cdot gh_1)_{-1})_2 S(h_2)_2 \\
&= S(h_2)_1 \otimes (c_{0,(0)} \cdot gh_1)_{0,(0)} \bar{S}((c_{0,(0)} \cdot gh_1)_{0,(-1)})_1 \\
&\quad \cdot \bar{S}((c_{0,(0)} \cdot gh_1)_{-1})_1 S(\bar{S}((c_{0,(0)} \cdot gh_1)_{-1})_2) \\
&\quad \cdot S(\bar{S}((c_{0,(0)} \cdot gh_1)_{0,(-1)})_2) \bar{S}((c_{0,(0)} \cdot gh_1)_{-1})_3 S(h_2)_2 \\
&= S(h_2)_1 \otimes (c_{0,(0)} \cdot gh_1)_{0,(0)} \\
&\quad \cdot \varepsilon((c_{0,(0)} \cdot gh_1)_{0,(-1)}) \bar{S}((c_{0,(0)} \cdot gh_1)_{-1}) S(h_2)_2 \quad (2.30) \\
&= S(h_2)_1 \otimes (c_{0,(0)} \cdot gh_1)_0 \cdot \bar{S}((c_{0,(0)} \cdot gh_1)_{-1}) S(h_2)_2 \\
&= S(h_2)_1 \otimes c_{0,(0),0} \cdot (gh_1)_3 \bar{S}((gh_1)_2) \bar{S}(c_{0,(0),-1}) \\
&\quad \cdot \bar{S}(S(gh_1)_1) S(h_2)_2 \\
&= S(h_3) \otimes c_{0,(0),0} \cdot \bar{S}(c_{0,(0),-1}) gh_1 S(h_2) \\
&= S(h) \otimes c_{0,(0),0} \cdot \bar{S}(c_{0,(0),-1}) g \\
&= S(\bar{S}(c_{-1})) \otimes c_{0,(0),0} \cdot \bar{S}(c_{0,(-1)} c_{0,(0),-1}) \\
&= c_{-1} \otimes c_0 = \tau(c).
\end{aligned}$$

In [12], it is also shown that $r(\gamma) \in U(\mathcal{T}(C))$, and it is straightforward to verify that the $*$ -inverse of $r(\gamma)$ is given by

$$r(\gamma)'(c) = S(S(c_1)_1) S(c_{0,(1)})_1 S(c_1)_2 \otimes c_{0,(0)} \cdot S(c_{0,(1)})_2 S(c_1)_3. \quad (2.31)$$

Then a routine verification similar to the above one shows that

$$\ell(r(\gamma))(c) = \gamma(c) \quad (2.32)$$

for all $c \in C$. It is easy to show that $\ell(\sigma) = \sigma'$ and $r(\sigma') = \sigma$. \square

2.2. The crossed coproduct. We recall the following definitions from [8, 10].

DEFINITION 2.2. Let C be a coalgebra and H a Hopf algebra. It is said that H coacts weakly on C if there is a k -linear map $\rho : C \rightarrow H \otimes C$; $\rho(c) = c_{[-1]} \otimes c_{[0]}$ satisfying the following conditions for all $c \in C$:

$$\begin{aligned}
c_{[-1]} \otimes c_{[0]1} \otimes c_{[0]2} &= c_{1[-1]} c_{2[-1]} \otimes c_{1[0]} \otimes c_{2[0]}, \\
\varepsilon_C(c_{[0]}) c_{[-1]} &= \varepsilon(c) 1_H, \quad (2.33) \\
\varepsilon_H(c_{[-1]}) c_{[0]} &= c.
\end{aligned}$$

Assume that H coacts weakly on C , and let $\alpha : C \rightarrow H \otimes H$, $\alpha(c) = \alpha_1(c) \otimes \alpha_2(c)$, be a linear map. Let $C \rtimes_{\alpha} H$ be the coalgebra whose underlying vector

space is $C \otimes H$, with comultiplication and counit given by

$$\begin{aligned}\Delta_\alpha(c \rtimes h) &= (c_1 \rtimes c_{2[-1]}\alpha_1(c_3)h_1) \otimes (c_{2[0]} \rtimes \alpha_2(c_3)h_2), \\ \varepsilon_\alpha(c \rtimes h) &= \varepsilon_C(c)\varepsilon_H(h).\end{aligned}\tag{2.34}$$

It was pointed out in [10] that $\varepsilon_\alpha(c \rtimes h)$ satisfies the counit property if and only if

$$(\varepsilon_H \otimes \text{id})\alpha(c) = (\text{id} \otimes \varepsilon_H)\alpha(c) = \varepsilon_C(c)1_H\tag{2.35}$$

and that Δ_α is coassociative if and only if α satisfies

$$\begin{aligned}c_{1[-1]}\alpha_1(c_2) \otimes \alpha_1(c_{1[0]})\alpha_2(c_2)_1 \otimes \alpha_2(c_{1[0]})\alpha_2(c_2)_2 \\ = \alpha_1(c_1)\alpha_1(c_2)_1 \otimes \alpha_2(c_1)\alpha_1(c_2)_2 \otimes \alpha_2(c_2),\end{aligned}\tag{2.36}$$

$$\begin{aligned}c_{1[-1]}\alpha_1(c_2) \otimes c_{1[0][-1]}\alpha_2(c_2) \otimes c_{1[0][0]} \\ = \alpha_1(c_1)c_{2[-1]1} \otimes \alpha_2(c_1)c_{2[-1]2} \otimes c_{2[0]}.\end{aligned}\tag{2.37}$$

In [10], (2.36) is called the cocycle condition and (2.37) is called the twisted comodule condition. Following [7], we call α , satisfying (2.35), (2.36), and (2.37), a Harrison 2-cocycle.

Now, consider two weak H -coactions $\rho, \rho' : C \rightarrow H \otimes C$, and write

$$\rho(c) = c_{[-1]} \otimes c_{[0]}, \quad \rho'(c) = c_{(-1)} \otimes c_{(0)}.\tag{2.38}$$

Also consider two 2-cocycles $\alpha, \alpha' : C \rightarrow H \otimes H$ corresponding respectively to ρ and ρ' , and write

$$\alpha(c) = \alpha_1(c) \otimes \alpha_2(c), \quad \alpha'(c) = \alpha'_1(c) \otimes \alpha'_2(c).\tag{2.39}$$

Then we can consider the crossed coproducts $C \rtimes_\alpha H$ and $C \rtimes'_{\alpha'} H$. In the next lemma, we discuss when these are isomorphic.

LEMMA 2.3. *Consider a convolution invertible map $u : C \rightarrow H$ satisfying the conditions*

$$c_{(-1)} \otimes c_{(0)} = u^{-1}(c_1)c_{2[-1]}u(c_3) \otimes c_{2[0]},\tag{2.40}$$

$$\alpha'(c) = u^{-1}(c_1)c_{2[-1]}\alpha_1(c_3)u(c_4)_1 \otimes u^{-1}(c_{2[0]})\alpha_2(c_3)u(c_4)_2\tag{2.41}$$

for all $c \in C$. Then the map

$$\phi : C \rtimes'_{\alpha'} H \longrightarrow C \rtimes_\alpha H, \quad \phi(c \rtimes' h) = c_1 \rtimes u(c_2)h\tag{2.42}$$

is a left C -colinear right H -linear coalgebra isomorphism. Every left C -colinear right H -linear coalgebra isomorphism between $C \bowtie_{\alpha} H$ and $C \bowtie'_{\alpha'} H$ is of this type.

PROOF. The proof is a dual version of a similar statement for crossed products (see [13]). \square

It was shown in [12] that the crossed coproduct construction can be viewed as a special case of the twisting construction from Section 2.1. Let H be a Hopf algebra and C a right H -module coalgebra, and view $C \otimes H$ as a right H -module coalgebra, with the right H -action being induced by the multiplication by H . It was proved in [12] that there is a bijective correspondence between crossed coproduct structures on $C \otimes H$ and twistings of $C \otimes H$. We recall the description of this bijection.

Consider a weak coaction ρ and a 2-cocycle α giving rise to the crossed coproduct $C \bowtie_{\alpha} H$, and write

$$\rho(c) = c_{[-1]} \otimes c_{[0]}, \quad \alpha(c) = \alpha_1(c) \otimes \alpha_2(c). \quad (2.43)$$

The corresponding twisting $\tau : C \otimes H \rightarrow H \otimes C \otimes H$ is defined by

$$\tau(c \otimes h) = S(h_1)c_{1[-1]}\alpha_1(c_2)h_2 \otimes c_{1[0]} \otimes \alpha_2(c_2)h_3. \quad (2.44)$$

Conversely, if τ is a twisting of $C \otimes H$, then $(C \otimes H)^{\tau} = C \bowtie_{\alpha} H$, with weak coaction ρ and 2-cocycle α given by

$$\rho(c) = (\text{id} \otimes \text{id} \otimes \varepsilon_H)\tau(c \otimes 1), \quad (2.45)$$

$$\alpha(c) = (\text{id} \otimes \varepsilon_C \otimes \text{id})\tau(c \otimes 1). \quad (2.46)$$

3. Twisted 2-cocycles. Let H be a Hopf algebra with bijective antipode S and let \bar{S} be the composition inverse of S . Take an H -module coalgebra C , and let $B = C/CH^+$.

DEFINITION 3.1. A map $\alpha : C \rightarrow H \otimes H$, $\alpha(c) = \alpha_1(c) \otimes \alpha_2(c)$ is called a twisted 2-cocycle if the following conditions are satisfied for all $h \in H$ and $c \in C$:

$$(\text{id}_H \otimes \varepsilon_H)\alpha(c) = (\varepsilon_H \otimes \text{id}_H)\alpha(c) = \varepsilon_C(c)1_H, \quad (3.1)$$

$$\alpha(c \cdot h) = S(h_1)\alpha_1(c)h_2 \otimes \bar{S}(h_4)\alpha_2(c)h_3, \quad (3.2)$$

$$\begin{aligned} \alpha_1(c_1)\alpha_1(c_3)_1 \otimes c_2 \cdot \alpha_2(c_1)\alpha_1(c_3)_2 \otimes \alpha_2(c_3) \\ = \alpha_1(c_1) \otimes c_2 \cdot \alpha_1(c_3)\alpha_2(c_1)_1 \otimes \alpha_2(c_3)\alpha_2(c_1)_2. \end{aligned} \quad (3.3)$$

Our first result is the fact that the twisted 2-cocycles can be used to define twistings on C .

PROPOSITION 3.2. *With notation as above, if $\alpha : C \rightarrow H \otimes H$ is a twisted 2-cocycle, then the map*

$$\tau_\alpha : C \longrightarrow H \otimes C, \quad \tau_\alpha(c) = \alpha_1(c_1) \otimes c_2 \cdot \alpha_2(c_1), \quad (3.4)$$

is a twisting of C .

PROOF. It follows easily from (3.1) that τ_α satisfies the normalizing condition (2.9). Next, we compute

$$\begin{aligned} (c \cdot h)_{-1} \otimes (c \cdot h)_0 &= \alpha_1((c \cdot h)_1) \otimes (c \cdot h)_2 \cdot \alpha_2((c \cdot h)_1)_1 \\ &= \alpha_1(c_1 \cdot h_1) \otimes c_2 \cdot h_2 \alpha_2(c_1 \cdot h_1) \\ &= S(h_1) \alpha_1(c_1) h_2 \otimes c_2 \cdot h_5 \bar{S}(h_4) \alpha_2(c_1) h_3 \\ &= S(h_1) \alpha_1(c_1) h_2 \otimes c_2 \cdot \alpha_2(c_1) h_3 \\ &= S(h_1) c_{-1} h_2 \otimes c_0 \cdot h_3, \end{aligned} \quad (3.5)$$

and (2.13) follows easily. Finally, we compute the left- and right-hand sides of (2.14):

$$\begin{aligned} &c_{-1} \otimes c_{0,1} \cdot c_{0,2,-1} \otimes c_{0,2,0} \\ &= (1 \otimes \Delta_\tau) \tau_\alpha(c) = \alpha_1(c_1) \otimes (c_2 \cdot \alpha_2(c_1))_1 \cdot \alpha_1(((c_2 \cdot \alpha_2(c_1))_2)_1) \\ &\quad \otimes ((c_2 \cdot \alpha_2(c_1))_2)_2 \cdot \alpha_2(((c_2 \cdot \alpha_2(c_1))_2)_1) \\ &= \alpha_1(c_1) \otimes c_2 \cdot \alpha_2(c_1)_1 \alpha_1(c_3 \cdot \alpha_2(c_1)_2) \\ &\quad \otimes (c_4 \cdot \alpha_2(c_1)_3) \cdot \alpha_2(c_3 \cdot \alpha_2(c_1)_2) \\ &= \alpha_1(c_1) \otimes c_2 \cdot \alpha_2(c_1)_1 S(\alpha_2(c_1)_2) \alpha_1(c_3) \alpha_2(c_1)_3 \\ &\quad \otimes c_4 \cdot \alpha_2(c_1)_6 \bar{S}(\alpha_2(c_1)_5) \alpha_2(c_3) \alpha_2(c_1)_4 \\ &= \alpha_1(c_1) \otimes c_2 \cdot \alpha_1(c_3) \alpha_2(c_1)_1 \otimes c_4 \cdot \alpha_2(c_3) \alpha_2(c_1)_2, \\ &c_{1,-1} c_{2,-1,1} \otimes c_{1,0} \cdot c_{2,-1,2} \otimes c_{2,0} \\ &= \sum \alpha_1(c_{11}) \alpha_1(c_{21})_1 \otimes c_{12} \cdot \alpha_2(c_{11}) \cdot \alpha_1(c_{21})_2 \otimes c_{22} \cdot \alpha_2(c_{21}) \\ &= \alpha_1(c_1) \alpha_1(c_3)_1 \otimes c_2 \cdot \alpha_2(c_1) \alpha_1(c_3)_2 \otimes c_4 \cdot \alpha_2(c_3) \\ &= \alpha_1(c_1) \otimes c_2 \cdot \alpha_1(c_3) \alpha_2(c_1)_1 \otimes c_4 \cdot \alpha_2(c_3) \alpha_2(c_1)_2. \end{aligned} \quad (3.6)$$

Thus (2.14) follows, and τ_α is a twisting. \square

There is also a relation between twisted 2-cocycles and Harrison 2-cocycles. Let C be a right H -module coalgebra. Consider the trivial weak coaction $\rho(c) = 1 \otimes c$ and $\alpha : C \rightarrow H \otimes H$. Then the cocycle condition (2.36) and the twisted

comodule condition (2.37) of [Definition 2.2](#) take the following form:

$$\begin{aligned} \alpha_1(c_2) \otimes \alpha_1(c_1) \alpha_2(c_2)_1 \otimes \alpha_2(c_1) \alpha_2(c_2)_2 \\ = \alpha_1(c_1) \alpha_1(c_2)_1 \otimes \alpha_2(c_1) \alpha_1(c_2)_2 \otimes \alpha_2(c_2), \end{aligned} \quad (3.7)$$

$$\alpha_1(c_2) \otimes \alpha_2(c_2) \otimes c_1 = \alpha_1(c_1) \otimes \alpha_2(c_1) \otimes c_2. \quad (3.8)$$

The set of Harrison 2-cocyles corresponding to the trivial weak coaction is denoted by $Z_{\text{Harr}}^2(H, C)$. Thus, $Z_{\text{Harr}}^2(H, C)$ consists of maps satisfying (2.35), (3.7), and (3.8). The set of twisted 2-cocycles $\alpha^t : C \otimes H \rightarrow H \otimes H$ in the sense of [Definition 3.1](#) will be denoted by $Z_{\text{tw}}^2(H, C \otimes H)$.

PROPOSITION 3.3. *Let C be a right H -module coalgebra. There exists a biject in between $Z_{\text{Harr}}^2(H, C)$ and $Z_{\text{tw}}^2(H, C \otimes H)$.*

PROOF. Take $\alpha^t \in Z_{\text{tw}}^2(H, C \otimes H)$ and write

$$\alpha^t(c \otimes h) = \sum \alpha_1^t(c \otimes h) \otimes \alpha_2^t(c \otimes h). \quad (3.9)$$

For all $c \in C$ and $h \in H$, we have

$$\begin{aligned} \alpha_1^t(c_1 \otimes h_1) \alpha_1^t(c_2 \otimes h_2)_1 \otimes \alpha_2^t(c_1 \otimes h_1) \alpha_1^t(c_2 \otimes h_2)_2 \otimes \alpha_2^t(c_2 \otimes h_2) \\ = \alpha_1^t(c_1 \otimes h_1) \otimes \alpha_1^t(c_2 \otimes h_2) \alpha_2^t(c_1 \otimes h_1)_1 \otimes \alpha_2^t(c_2 \otimes h_2) \alpha_2^t(c_1 \otimes h_1)_2. \end{aligned} \quad (3.10)$$

Now, define $\alpha : C \rightarrow H \otimes H$ by $\alpha(c) = \alpha^t(c \otimes 1)$. It is easy to see that α satisfies (2.35) and (3.8). Using (3.10), we compute

$$\begin{aligned} \alpha_1(c_2) \otimes \alpha_1(c_1) \alpha_2(c_2)_1 \otimes \alpha_2(c_1) \alpha_2(c_2)_2 \\ = \alpha_1(c_1) \otimes \alpha_1(c_2) \alpha_2(c_1)_1 \otimes \alpha_2(c_2) \alpha_2(c_1)_2 \\ = \alpha_1^t(c_1 \otimes 1) \otimes \alpha_1^t(c_2 \otimes 1) \alpha_2^t(c_1 \otimes 1)_1 \otimes \alpha_2^t(c_2 \otimes 1) \alpha_2^t(c_1 \otimes 1)_2 \\ = \alpha_1^t(c_1 \otimes 1) \alpha_1^t(c_2 \otimes 1)_1 \otimes \alpha_2^t(c_1 \otimes 1) \alpha_1^t(c_2 \otimes 1)_2 \otimes \alpha_2^t(c_2 \otimes 1) \\ = \alpha_1(c_1) \alpha_1(c_2)_1 \otimes \alpha_2(c_1) \alpha_1(c_2)_2 \otimes \alpha_2(c_2), \end{aligned} \quad (3.11)$$

and it follows that α also satisfies (3.7).

Conversely, let $\alpha \in Z_{\text{Harr}}^2(H, C)$, and define $\alpha^t : C \otimes H \rightarrow H \otimes H$ by

$$\alpha^t(c \otimes h) = S(h_1) \alpha_1(c) h_2 \otimes \bar{S}(h_4) \alpha_2(c) h_3. \quad (3.12)$$

We can easily show that α^t satisfies conditions (3.1) and (3.2) of [Definition 3.1](#). A straightforward computation shows that (3.3) is also satisfied:

$$\begin{aligned}
& \alpha_1^t(c_1 \otimes h_1) \alpha_1^t(c_3 \otimes h_3)_1 \otimes c_2 \otimes h_2 \alpha_2^t(c_1 \otimes h_1) \alpha_1^t(c_3 \otimes h_3)_2 \otimes \alpha_2^t(c_3 \otimes h_3) \\
&= S h_1 \alpha_1(c_1) h_2 S(h_7) \alpha_1(c_3)_1 h_8 \\
&\quad \otimes c_2 \otimes h_5 \bar{S}(h_4) \alpha_2(c_1) h_3 S(h_6) \alpha_1(c_3)_2 h_9 \otimes \bar{S}(h_{11}) \alpha_2(c_3) h_{10} \\
&= S(h_1) \alpha_1(c_1) \alpha_1(c_3)_1 h_2 \otimes c_2 \otimes \alpha_2(c_1) \alpha_1(c_3)_2 h_3 \otimes \bar{S}(h_5) \alpha_2(c_3) h_4 \\
&= S(h_1) \alpha_1(c_3) h_2 \otimes c_2 \otimes \alpha_1(c_1) \alpha_2(c_3)_1 h_3 \otimes \bar{S}(h_5) \alpha_2(c_1) \alpha_2(c_3)_2 h_4 \\
&= S(h_1) \alpha_1(c_1) h_2 \otimes c_2 \otimes \alpha_1(c_3) \alpha_2(c_1)_1 h_3 \otimes \bar{S}(h_5) \alpha_2(c_3) \alpha_2(c_1)_2 h_4 \\
&= S(h_1) \alpha_1(c_1) h_2 \otimes c_2 \otimes h_7 S(h_8) \alpha_1(c_3) h_9 \bar{S}(h_6) \alpha_2(c_1)_1 h_3 \\
&\quad \otimes \bar{S}(h_{11}) \alpha_2(c_3) h_{10} \bar{S}(h_5) \alpha_2(c_1)_2 h_4 \\
&= \alpha_1^t(c_1 \otimes h_1) \otimes c_2 \otimes h_2 \alpha_1^t(c_3 \otimes h_3) \alpha_2^t(c_1 \otimes h_1)_1 \otimes \alpha_2^t(c_3 \otimes h_3) \alpha_2^t(c_1 \otimes h_1)_2. \tag{3.13}
\end{aligned}$$

So it follows that α^t is a twisted 2-cocycle. We leave it to the reader to show that the maps between $Z_{\text{Harr}}^2(H, C)$ and $Z_{\text{tw}}^2(H, C \otimes H)$ defined above are the inverses of each other. \square

4. Equivalence of twistings. In this section, we will define an equivalence relation on the set of twistings of an H -module coalgebra C . If a twisting is invertible, then all other twistings in the same equivalence class are also invertible.

PROPOSITION 4.1. *Take $\tau, \lambda \in \mathcal{T}(C)$ and use notation [\(2.24\)](#). Consider $v \in \text{Hom}(C, H)$ satisfying the following identities, for all $h \in H, c \in C$:*

$$\varepsilon_H \circ v = \varepsilon_C, \quad v(c \cdot h) = S(h_1)v(c)h_2, \tag{4.1}$$

$$c_{1,(-1)}v(c_2)_1 \otimes c_{1,(0)} \cdot v(c_2)_2 = v(c_1)c_{2,-1} \otimes c_{2,0,1} \cdot v(c_{2,0,2}). \tag{4.2}$$

Then $\psi : C^\tau \rightarrow C^\lambda$, $\psi(c) = c_1 \cdot v(c_2)$, is a left B -colinear and right H -linear coalgebra map inducing the identity map on B . If $v \in \text{Reg}(C, H)$, then ψ is an isomorphism.

PROOF. Using the second identity in (4.1) and $B = C/CH^+$, we can easily prove that ψ is left B -colinear and right H -linear. Using the first identity in (4.1), we obtain that ψ induces a well-defined map $B \rightarrow B$, which is the identity. In order to prove that ψ is a coalgebra map, we need to check that

$$\psi(c_1 \cdot c_{2,-1}) \otimes \psi(c_{2,0}) = \psi(c)_1 \psi(c)_{2,(-1)} \otimes \psi(c)_{2,(0)}. \tag{4.3}$$

Again, we compute the left- and right-hand sides and see that they are equal:

$$\begin{aligned}
& \psi(c)_1 \psi(c)_{2,(-1)} \otimes \psi(c)_{2,(0)} \\
&= (c_1 \cdot v(c_2))_1 (c_1 \cdot v(c_2))_{2,(-1)} \otimes (c_1 \cdot v(c_2))_{2,(0)} \\
&= c_1 \cdot v(c_3)_1 (c_2 \cdot v(c_3)_2)_{(-1)} \otimes (c_2 \cdot v(c_3)_2)_{(0)} \\
&= c_1 \cdot v(c_3)_1 S(v(c_3)_2) c_{2,(-1)} v(c_3)_3 \otimes c_{2,(0)} \cdot v(c_3)_4 \\
&= c_1 \cdot c_{2,(-1)} v(c_3)_1 \otimes c_{2,(0)} \cdot v(c_3)_2 \\
&= c_1 \cdot v(c_2) c_{3,-1} \otimes c_{3,0,1} \cdot v(c_{3,0,2}), \\
& \psi(c_1 \cdot c_{2,-1}) \otimes \psi(c_{2,0}) \\
&= (c_1 \cdot c_{2,-1})_1 \cdot v((c_1 \cdot c_{2,-1})_2) \otimes (c_{2,0})_1 \cdot v((c_{2,0})_2) \\
&= c_1 \cdot (c_{3,-1})_1 v(c_2 \cdot (c_{3,-1})_2) \otimes (c_{3,0})_1 \cdot v((c_{3,0})_2) \\
&= c_1 \cdot c_{3,-1,1} S(c_{3,-1,2}) v(c_2) c_{3,-1,3} \otimes c_{3,0,1} \cdot v(c_{3,0,2}) \\
&= c_1 \cdot v(c_2) c_{3,-1} \otimes c_{3,0,1} \cdot v(c_{3,0,2}).
\end{aligned} \tag{4.4}$$

If $v \in \text{Reg}(C, H)$, then its inverse w also satisfies (4.1), and $\varphi : C^\lambda \rightarrow C^\tau$ defined by

$$\varphi(c) = c_1 \cdot w(c_2) \tag{4.5}$$

is the inverse of ψ . □

DEFINITION 4.2. It is said that $\tau, \lambda \in \mathcal{T}(C)$ are equivalent if there exists $v \in \text{Reg}(C, H)$ satisfying the conditions of [Proposition 4.1](#). This is denoted by $\tau \sim \lambda$.

LEMMA 4.3. *The relation \sim is an equivalence relation on $\mathcal{T}(C)$.*

PROOF. Clearly, $\tau \sim \tau$ through $v(c) = \varepsilon(c)1_H$.

Next, assume that $\tau \sim \lambda$, and take $v \in \text{Reg}(C, H)$ satisfying (4.1) and (4.2). Equation (4.2) is equivalent to

$$c_{(-1)} \otimes c_{(0)} = v(c_1) c_{2,-1} v^{-1}(c_3)_1 \otimes c_{2,0,1} \cdot v(c_{2,0,2}) v^{-1}(c_3)_2. \tag{4.6}$$

The inverse u of v satisfies (4.1). It also satisfies (4.2) since

$$\begin{aligned}
& u(c_1) c_{2,(-1)} \otimes c_{2,(0)1} \cdot u(c_{2,(0)2}) \\
&= u(c_1) v(c_2) c_{3,-1} v^{-1}(c_4)_1 \otimes c_{3,0,1} \cdot v(c_{3,0,3})_1 v^{-1}(c_4)_2 \\
&\quad \cdot S(v^{-1}(c_4)_3) S(v(c_{3,0,3})_2) u(c_{3,0,2}) v(c_{3,0,3})_3 v^{-1}(c_4)_4 \\
&= c_{1,-1} v^{-1}(c_2)_1 \otimes c_{1,0,1} \cdot u(c_{1,0,2}) v(c_{1,0,3}) v^{-1}(c_2)_2 \\
&= c_{1,-1} u(c_2)_1 \otimes c_{1,0,1} \cdot u(c_2)_2,
\end{aligned} \tag{4.7}$$

and it follows that $\lambda \sim \tau$.

Now, assume that $\tau \sim \lambda$ and $\lambda \sim \gamma$, and take the corresponding maps $v, u \in \text{Reg}(C, H)$. Set $w = u * v$, and write

$$\tau(c) = c_{-1} \otimes c_0, \quad \lambda(c) = c_{(-1)} \otimes c_{(0)}, \quad \gamma(c) = c_{[-1]} \otimes c_{[0]}. \quad (4.8)$$

It is easily shown that w satisfies (4.1), v satisfies (4.6), and u satisfies

$$c_{[-1]} \otimes c_{[0]} = u(c_1)c_{2,(-1)}u^{-1}(c_3)_1 \otimes c_{2,(0),1} \cdot u(c_{2,(0),2})u^{-1}(c_3)_2. \quad (4.9)$$

We compute

$$\begin{aligned} c_{[-1]} \otimes c_{[0]} &= u(c_1)v(c_2)c_{3,-1}v^{-1}(c_4)_1u^{-1}(c_5)_1 \otimes c_{3,0,1} \\ &\quad \cdot v(c_{3,0,3})_1v^{-1}(c_4)_2S(v^{-1}(c_4)_3)S(v(c_{3,0,3})_2) \\ &\quad \cdot u(c_{3,0,2})v(c_{3,0,3})_3v^{-1}(c_4)_4u^{-1}(c_5)_2 \\ &= u(c_1)v(c_2)c_{3,-1}v^{-1}(c_4)_1u^{-1}(c_5)_1 \\ &\quad \otimes c_{3,0,1} \cdot u(c_{3,0,2})v(c_{3,0,3})v^{-1}(c_4)_2u^{-1}(c_5)_2 \\ &= (u * v)(c_1)c_{2,-1}(u * v)^{-1}(c_3)_1 \\ &\quad \otimes c_{2,0,1} \cdot (u * v)(c_{2,0,2})(u * v)^{-1}(c_3)_2, \end{aligned} \quad (4.10)$$

and this proves that $\tau \sim \gamma$. \square

THEOREM 4.4. *Take $\tau \sim \lambda \in \mathcal{T}(C)$. If τ is invertible, then λ is also invertible.*

PROOF. Take $v \in \text{Reg}(C, H)$ satisfying the conditions in Proposition 4.1 and let $\psi : C^\tau \rightarrow C^\lambda$ be the coalgebra isomorphism given by

$$\psi(c) = c_1 \cdot v(c_2). \quad (4.11)$$

Let τ^{-1} be the inverse of τ , and write

$$\tau^{-1}(c) = c_{(-1)} \otimes c_{(0)}, \quad \tau(c) = c_{-1} \otimes c_0, \quad \lambda(c) = c_{(-1)} \otimes c_{(0)}. \quad (4.12)$$

Define $\mu : C \rightarrow H \otimes C$ by

$$\begin{aligned} \mu(c) &= c_{[-1]} \otimes c_{[0]} = \psi^{-1}(c)_{(-1)}v^{-1}(\psi^{-1}(c)_{(0)1})v(\psi^{-1}(c)_{(0)3})_1 \\ &\quad \otimes \psi^{-1}(c)_{(0)2} \cdot v(\psi^{-1}(c)_{(0)3})_2. \end{aligned} \quad (4.13)$$

Using the temporary notation $\psi^{-1}(c)_{(-1)} = a$ and $\psi^{-1}(c)_{(0)} = b$, it is not hard to prove that μ is a left inverse of λ . Indeed,

$$\begin{aligned}
(\mu * \lambda)(c) &= (m \otimes \text{id})(\text{id} \otimes \lambda)\mu(c) \\
&= a \nu^{-1}(b_1) \nu(b_3)_1 (b_2 \cdot \nu(b_3)_2)_{(-1)} \otimes (b_2 \cdot \nu(b_3))_{(0)} \\
&= a \nu^{-1}(b_1) \nu(b_3)_1 S(\nu(b_3)_2) b_{2,(-1)} \nu(b_3)_3 \otimes b_{2,(0)} \cdot \nu(b_3)_4 \\
&= a \nu^{-1}(b_1) b_{2,(-1)} \nu(b_3)_1 \otimes b_{2,(0)} \cdot \nu(b_3)_2 \\
&= a \nu^{-1}(b_1) \nu(b_2) b_{3,-1} \otimes b_{3,0,1} \cdot \nu(b_{3,0,2}) \\
&= ab_{-1} \otimes b_{0,1} \cdot \nu(b_{0,2}) = 1 \otimes \psi^{-1}(c)_1 \cdot \nu(\psi^{-1}(c)_2) \\
&= 1 \otimes \psi(\psi^{-1}(c)) = 1 \otimes c = \sigma(c).
\end{aligned} \tag{4.14}$$

The proof of the fact that μ is also a right inverse of λ is much more technical. From the fact that ν is invertible, and using (4.2), we obtain

$$\lambda(c) = c_{(-1)} \otimes c_{(0)} = \nu(c_1) c_{2,-1} \nu^{-1}(c_3)_1 \otimes c_{2,0,1} \cdot \nu(c_{2,0,2}) \nu^{-1}(c_3)_2. \tag{4.15}$$

Now, set $\psi^{-1}(c) = c_1 \cdot \nu^{-1}(c_2)$. We compute

$$\begin{aligned}
(\lambda * \mu)(c) &= (m \otimes \text{id})(\text{id} \otimes \mu)\lambda(c) \\
&= \nu(c_1) c_{2,-1} \nu^{-1}(c_3)_1 (c_{2,0,1} \cdot \nu(c_{2,0,2}) \nu^{-1}(c_3)_2)_{[-1]} \\
&\quad \otimes (c_{2,0,1} \cdot \nu(c_{2,0,2}) \nu^{-1}(c_3)_2)_{[0]} \\
&= \nu(c_1) c_{2,-1} \nu^{-1}(c_3)_1 S(\nu^{-1}(c_3)_2) S(\nu(c_{2,0,2})_1) (c_{2,0,1})_{[-1]} \\
&\quad \cdot \nu(c_{2,0,2})_2 \nu^{-1}(c_3)_3 \otimes (c_{2,0,1})_{[0]} \cdot \nu(c_{2,0,2})_3 \nu^{-1}(c_3)_4 \\
&= \nu(c_1) c_{2,-1} S(\nu(c_{2,0,2})_1) (c_{2,0,1})_{[-1]} \nu(c_{2,0,2})_2 \nu^{-1}(c_3)_1 \\
&\quad \otimes (c_{2,0,1})_{[0]} \cdot \nu(c_{2,0,2})_3 \nu^{-1}(c_3)_2 \\
&= \nu(c_1) c_{2,-1} S(\nu(c_{2,0,2})_1) \psi^{-1}(c_{2,0,1})_{(-1)} \nu^{-1}(\psi^{-1}(c_{2,0,1})_{(0),1}) \\
&\quad \cdot \nu(\psi^{-1}(c_{2,0,1})_{(0),3})_1 \nu(c_{2,0,2})_2 \nu^{-1}(c_3)_1 \otimes \psi^{-1}(c_{2,0,1})_{(0),2} \\
&\quad \cdot \nu(\psi^{-1}(c_{2,0,1})_{(0),3})_2 \nu(c_{2,0,2})_3 \nu^{-1}(c_3)_2 \\
&= \nu(c_1) c_{2,-1} S(\nu(c_{2,0,3})_1) (c_{2,0,1} \cdot \nu^{-1}(c_{2,0,2}))_{(-1)} \\
&\quad \cdot \nu^{-1}((c_{2,0,1} \cdot \nu^{-1}(c_{2,0,2}))_{(0),1}) \nu((c_{2,0,1} \cdot \nu^{-1}(c_{2,0,2}))_{(0),3})_1 \\
&\quad \cdot \nu(c_{2,0,3})_2 \nu^{-1}(c_3)_1 \otimes (c_{2,0,1} \cdot \nu^{-1}(c_{2,0,2}))_{(0),2} \\
&\quad \cdot \nu((c_{2,0,1} \cdot \nu^{-1}(c_{2,0,2}))_{(0),3})_2 \nu(c_{2,0,3})_3 \nu^{-1}(c_3)_2 \\
&= \nu(c_1) c_{2,-1} S(\nu(c_{2,0,3})_1) S(\nu^{-1}(c_{2,0,2})_1) c_{2,0,1,(-1)} \nu^{-1}(c_{2,0,2})_2 \\
&\quad \cdot \nu^{-1}((c_{2,0,1,(0)} \cdot \nu^{-1}(c_{2,0,2})_3)_1) \nu((c_{2,0,1,(0)} \cdot \nu^{-1}(c_{2,0,2})_3)_3)_1 \\
&\quad \cdot \nu(c_{2,0,3})_2 \nu^{-1}(c_3)_1 \otimes (c_{2,0,1,(0)} \cdot \nu^{-1}(c_{2,0,2})_3)_2 \\
&\quad \cdot \nu((c_{2,0,1,(0)} \cdot \nu^{-1}(c_{2,0,2})_3)_3)_2 \nu(c_{2,0,3})_3 \nu^{-1}(c_3)_2
\end{aligned}$$

$$\begin{aligned}
&= \nu(c_1)c_{2,-1}S(\nu^{-1}(c_{2,0,2})_1\nu(c_{2,0,3})_1)c_{2,0,1,(-1)}\nu^{-1}(c_{2,0,2})_2 \\
&\quad \cdot \nu^{-1}(c_{2,0,1,\langle 0 \rangle,1} \cdot \nu^{-1}(c_{2,0,2})_3)\nu(c_{2,0,1,\langle 0 \rangle,3} \cdot \nu^{-1}(c_{2,0,2})_5)_1\nu(c_{2,0,3})_2 \\
&\quad \cdot \nu^{-1}(c_3)_1 \otimes c_{2,0,1,\langle 0 \rangle,2} \cdot \nu^{-1}(c_{2,0,2})_4 \\
&\quad \cdot \nu(c_{2,0,1,\langle 0 \rangle,3} \cdot \nu^{-1}(c_{2,0,2})_5)_2\nu(c_{2,0,3})_3\nu^{-1}(c_3)_2 \\
&= \nu(c_1)c_{2,-1}S(\nu^{-1}(c_{2,0,2})_1\nu(c_{2,0,3})_1)c_{2,0,1,(-1)}\nu^{-1}(c_{2,0,2})_2 \\
&\quad \cdot S(\nu^{-1}(c_{2,0,2})_3)\nu^{-1}(c_{2,0,1,\langle 0 \rangle,1})\nu^{-1}(c_{2,0,2})_4S(\nu^{-1}(c_{2,0,2})_6)_1 \\
&\quad \cdot \nu(c_{2,0,1,\langle 0 \rangle,3})_1(\nu^{-1}(c_{2,0,2})_7)_1\nu(c_{2,0,3})_2\nu^{-1}(c_3)_1 \\
&\quad \otimes c_{2,0,1,\langle 0 \rangle,2} \cdot \nu^{-1}(c_{2,0,2})_5S(\nu^{-1}(c_{2,0,2})_6)_2 \\
&\quad \cdot \nu(c_{2,0,1,\langle 0 \rangle,3})_2(\nu^{-1}(c_{2,0,2})_7)_2\nu(c_{2,0,3})_3\nu^{-1}(c_3)_2 \\
&= \nu(c_1)c_{2,-1}S(\nu^{-1}(c_{2,0,2})_1\nu(c_{2,0,3})_1)c_{2,0,1,(-1)}\nu^{-1}(c_{2,0,2})_2S(\nu^{-1}(c_{2,0,2})_3) \\
&\quad \cdot \nu^{-1}(c_{2,0,1,\langle 0 \rangle,1})\nu^{-1}(c_{2,0,2})_4S(\nu^{-1}(c_{2,0,2})_7)\nu(c_{2,0,1,\langle 0 \rangle,3})_1 \\
&\quad \cdot \nu^{-1}(c_{2,0,2})_8\nu(c_{2,0,3})_2\nu^{-1}(c_3)_1 \otimes c_{2,0,1,\langle 0 \rangle,2} \cdot \nu^{-1}(c_{2,0,2})_5 \\
&\quad \cdot S(\nu^{-1}(c_{2,0,2})_6)\nu(c_{2,0,1,\langle 0 \rangle,3})_2\nu^{-1}(c_{2,0,2})_9\nu(c_{2,0,3})_3\nu^{-1}(c_3)_2 \\
&= \nu(c_1)c_{2,-1}S(\nu^{-1}(c_{2,0,2})_1\nu(c_{2,0,3})_1)c_{2,0,1,(-1)}\nu^{-1}(c_{2,0,1,\langle 0 \rangle,1}) \\
&\quad \cdot \nu(c_{2,0,1,\langle 0 \rangle,3})_1\nu^{-1}(c_{2,0,2})_2\nu(c_{2,0,3})_2\nu^{-1}(c_3)_1 \\
&\quad \otimes c_{2,0,1,\langle 0 \rangle,2} \cdot \nu(c_{2,0,1,\langle 0 \rangle,3})_2\nu^{-1}(c_{2,0,2})_3\nu(c_{2,0,3})_3\nu^{-1}(c_3)_2 \\
&= \nu(c_1)c_{2,-1}c_{2,0,(-1)}\nu^{-1}(c_{2,0,\langle 0 \rangle,1})\nu(c_{2,0,\langle 0 \rangle,3})_1\nu^{-1}(c_3)_1 \\
&\quad \otimes c_{2,0,\langle 0 \rangle,2} \cdot \nu(c_{2,0,\langle 0 \rangle,3})_2\nu^{-1}(c_3)_2 \\
&= \nu(c_1)\nu^{-1}(c_{2,1})\nu(c_{2,3})_1\nu^{-1}(c_3)_1 \otimes c_{2,2} \cdot \nu(c_{2,3})_2\nu^{-1}(c_3)_2 \\
&= \nu(c_1)\nu^{-1}(c_2)\nu(c_4)_1\nu^{-1}(c_5)_1 \otimes c_3 \cdot \nu(c_4)_2\nu^{-1}(c_5)_2 \\
&= 1 \otimes c = \sigma(c),
\end{aligned} \tag{4.16}$$

and it follows that λ is convolution invertible. \square

THEOREM 4.5. *Let C be a right H -comodule algebra and consider $\tau, \lambda \in \mathcal{T}(C \otimes H)$. Then τ and λ are equivalent in the sense of [Definition 4.2](#) if and only if there is a left C -colinear and right H -linear coalgebra isomorphism between the crossed coproducts $C \rtimes_{\alpha} H$, ρ and $C \rtimes'_{\alpha'} H$, ρ' corresponding to τ and λ .*

PROOF. Consider [\(2.38\)](#). If $\tau \sim \lambda$, then there exists $\nu \in \text{Reg}(C \otimes H, H)$ satisfying [\(4.1\)](#) and [\(4.2\)](#). Define

$$u : C \longrightarrow H, \quad u(c) = \nu^{-1}(c \otimes 1). \tag{4.17}$$

If we can show that u satisfies (2.40) and (2.41), then one implication is proved by Lemma 2.3. It follows from (4.2) that

$$\begin{aligned} & (c_1 \otimes 1)_{(-1)} v(c_2 \otimes 1)_1 \otimes (c_1 \otimes 1)_{(0)} \cdot v(c_2 \otimes 1)_2 \\ &= v(c_1 \otimes 1)(c_2 \otimes 1)_{-1} \otimes (c_2 \otimes 1)_{0,1} \cdot v((c_2 \otimes 1)_{0,2}). \end{aligned} \quad (4.18)$$

Applying $1 \otimes 1 \otimes \varepsilon$ to both sides, we find

$$\begin{aligned} & (c_1 \otimes 1)_{(-1)} v(c_2 \otimes 1) \otimes (1 \otimes \varepsilon)(c_1 \otimes 1)_{(0)} \\ &= v(c_1 \otimes 1)(c_2 \otimes 1)_{-1} \otimes (1 \otimes \varepsilon)(c_2 \otimes 1)_0, \end{aligned} \quad (4.19)$$

and using (2.45), we obtain

$$c_{\langle -1 \rangle} \otimes c_{\langle 0 \rangle} = u^{-1}(c_1)c_{2[-1]}u(c_3) \otimes c_{2[0]}. \quad (4.20)$$

So u satisfies (2.40).

Applying $1 \otimes \varepsilon \otimes 1$ to both sides of (4.18), we find

$$\begin{aligned} \alpha'(c) &= v(c_1 \otimes 1)(c_2 \otimes 1)_{-1} v^{-1}(c_3 \otimes 1)_1 \\ &\otimes (\varepsilon \otimes 1)((c_2 \otimes 1)_{0,1} \cdot v((c_2 \otimes 1)_{0,2}) v^{-1}(c_3 \otimes 1)_2). \end{aligned} \quad (4.21)$$

It follows from (2.44) that

$$\begin{aligned} & (c \otimes 1)_0 = c_{1[0]} \otimes \alpha_2(c_2), \\ & (\varepsilon \otimes 1)((c_2 \otimes 1)_{0,1} \cdot v((c_2 \otimes 1)_{0,2}) v^{-1}(c_3 \otimes 1)_2) \\ &= (\varepsilon \otimes 1)(c_{2,[0],1} \otimes \alpha_2(c_3)_1 v(c_{2,[0],2} \otimes \alpha_2(c_3)_2) v^{-1}(c_4 \otimes 1)_2) \\ &= \alpha_2(c_3)_1 v(c_{2,[0]} \otimes \alpha_2(c_3)_2) v^{-1}(c_4 \otimes 1)_2 \\ &= v(c_{2,[0]} \otimes 1) \alpha_2(c_3) v^{-1}(c_4 \otimes 1)_2 \\ &= v((1 \otimes \varepsilon)(c_2 \otimes 1)_0 \otimes 1)(\varepsilon \otimes 1)(c_3 \otimes 1)_0 v^{-1}(c_4 \otimes 1)_2. \end{aligned} \quad (4.22)$$

So,

$$\begin{aligned} \alpha'(c) &= v(c_1 \otimes 1)(c_2 \otimes 1)_{-1} (c_3 \otimes 1)_{-1} v^{-1}(c_4 \otimes 1)_1 \\ &\otimes v((1 \otimes \varepsilon)(c_2 \otimes 1)_0 \otimes 1)(\varepsilon \otimes 1)(c_3 \otimes 1)_0 v^{-1}(c_4 \otimes 1)_2 \\ &= u^{-1}(c_1)c_{2[-1]}\alpha_1(c_3)u(c_4)_1 \otimes u^{-1}(c_{2[0]})\alpha_2(c_3)u(c_4)_2, \end{aligned} \quad (4.23)$$

and (2.41) follows.

Conversely, assume that the two crossed coproducts are isomorphic. By [Lemma 2.3](#), there exists $u \in \text{Reg}(C, H)$ satisfying [\(2.40\)](#) and [\(2.41\)](#). Define

$$\nu : C \otimes H \longrightarrow H, \quad \nu(c \otimes h) = S(h_1)u^{-1}(c)h_2. \quad (4.24)$$

Then

$$\begin{aligned} \varepsilon_H \nu(c \otimes h) &= \varepsilon(S(h_1)u^{-1}(c)h_2) = \varepsilon(c)\varepsilon(h), \\ \nu((c \otimes h) \cdot g) &= \nu(c \otimes hg) = S(hg)_1u^{-1}(c)(hg)_2 \\ &= S(g_1)S(h_1)u^{-1}(c)h_2g_2 = S(g_1)\nu(c \otimes h)g_2. \end{aligned} \quad (4.25)$$

So,

$$\begin{aligned} \lambda(c \otimes h) &= (c \otimes h)_1 \otimes (c \otimes h)_0 \\ &= S(h_1)c_{1,(-1)}\alpha'_1(c_2)h_2 \otimes c_{1,(0)} \otimes \alpha'_2(c_2)h_3 \\ &= S(h_1)u^{-1}(c_1)c_{2,[-1]}u(c_3)u^{-1}(c_4)c_{5,[-1]}\alpha_1(c_6)u(c_7)_1h_2 \\ &\quad \otimes c_{2,[0]} \otimes u^{-1}(c_{5,[0]})\alpha_2(c_6)u(c_7)_2h_3 \\ &= S(h_1)u^{-1}(c_1)c_{2,[-1]}\alpha_1(c_3)u(c_4)_1h_2 \otimes c_{2,[0],1} \\ &\quad \otimes u^{-1}(c_{2,[0],2})\alpha_2(c_3)u(c_4)_2h_3 \\ &= S(h_1)u^{-1}(c_1)h_2S(h_3)c_{2,[-1]}\alpha_1(c_3)h_4S(h_7)u(c_4)_1h_8 \\ &\quad \otimes c_{2,[0],1} \otimes u^{-1}(c_{2,[0],2})\alpha_2(c_3)h_5S(h_6)u(c_4)_2h_9 \\ &= S(h_1)u^{-1}(c_1)h_2S(h_3)c_{2,[-1]}\alpha_1(c_3)h_4S(h_9)u(c_4)_1h_{10} \otimes c_{2,[0],1} \\ &\quad \otimes \alpha_2(c_3)_1h_5S(h_6)S(\alpha_2(c_3)_2)u^{-1}(c_{2,[0],2})\alpha_2(c_3)_3h_7S(h_8)u(c_4)_2h_{11} \\ &= \nu(c_1 \otimes h_1)(c_2 \otimes h_2)_{-1}\nu^{-1}(c_3 \otimes h_3)_1 \\ &\quad \otimes (c_2 \otimes h_2)_{0,1} \cdot \nu((c_2 \otimes h_2)_{0,2})\nu^{-1}(c_3 \otimes h_3)_2. \end{aligned} \quad (4.26)$$

This shows that $\tau \sim \lambda$. □

5. Twisting Hopf-Galois coextensions. Let H be a Hopf algebra with bijective antipode S , and C a right H -module coalgebra. As before, we use the following notation:

$$B = C/I, \quad I = \{c(h - \varepsilon(h)) \mid h \in H, c \in C\}. \quad (5.1)$$

For $\tau \in \mathcal{T}(C)$, we have that $C^\tau/I^\tau = C/I = B$.

Now, assume that C/B is an H -Galois coextension (see [\[5\]](#)). This means that the canonical map

$$\beta : C \otimes H \longrightarrow C \square_B C, \quad \beta(c \otimes h) = c_1 \otimes c_2 \cdot h, \quad (5.2)$$

is a bijection. Recall that, in this situation, $C \square_B C$ is the cotensor product

$$C \square_B C = \left\{ \sum_i C_i \otimes d_i \in C \otimes C \mid \sum_i c_{i_1} \otimes \pi(c_{i_2}) \otimes d_i = \sum_i C_i \otimes \pi(d_{i_1}) \otimes d_{i_2} \right\}, \quad (5.3)$$

where $\pi : C \rightarrow B$ is the natural epimorphism.

LEMMA 5.1. *With notation as above, consider the map*

$$\beta' : C \otimes H \longrightarrow C \square_B C, \quad \beta'(c \otimes h) = c_1 \cdot h \otimes c_2. \quad (5.4)$$

If the antipode S is bijective, then β is bijective (resp., injective, resp., surjective) if and only if β' is bijective (resp., injective, resp., surjective).

PROOF. The map

$$\phi : C \otimes H \longrightarrow C \otimes H, \quad \phi(c \otimes h) = c \cdot h_1 \otimes S(h_2) \quad (5.5)$$

is a bijection with inverse

$$\phi^{-1}(c \otimes h) = c \cdot h_2 \otimes \bar{S}h_1. \quad (5.6)$$

Then the statement follows from the fact that $\beta' = \beta \circ \phi$. \square

THEOREM 5.2. *Take $\tau \in U(\mathcal{T}(C))$. Then C^τ/B is an H -Galois coextension if and only if C/B is an H -Galois coextension.*

PROOF. Let λ be the inverse of τ . As before, we use the notation (2.24). Let β^τ be the canonical map corresponding to the coextension C^τ/B , that is,

$$\beta^\tau(c \otimes h) = c_1 \cdot c_{2,-1} h \otimes c_{2,0}. \quad (5.7)$$

Consider the following diagram:

$$\begin{array}{ccc} C \otimes H & \xrightarrow{\beta} & C \square_B C \\ f \downarrow & & \downarrow g \\ C \otimes H & \xrightarrow{\beta^\tau} & C \square_B C, \end{array} \quad (5.8)$$

where

$$f(c \otimes h) = c_0 \otimes \bar{S}(c_{-1})h, \quad g(c \otimes d) = c_0 \cdot \bar{S}(c_{-1}) \otimes d, \quad (5.9)$$

f and g are bijections, with inverses given by

$$f^{-1}(c \otimes h) = c_{(0)} \otimes \bar{S}(c_{(-1)})h, \quad g^{-1}(c \otimes d) = c_{(0)} \cdot \bar{S}(c_{(-1)}) \otimes d. \quad (5.10)$$

We can also compute

$$\begin{aligned}
\beta^\tau f(c \otimes h) &= \beta^\tau(c_0 \otimes \bar{S}(c_{-1})h) \\
&= c_{0,1} \cdot c_{0,2,-1} \bar{S}(c_{-1})h \otimes c_{0,2,0} \\
&= c_{1,0} \cdot c_{2,-1,2} \bar{S}(c_{1,-1}c_{2,-1,1})h \otimes c_{2,0} \\
&= c_{1,0} \cdot c_{2,-1,2} \bar{S}(c_{2,-1,1}) \bar{S}(c_{1,-1})h \otimes c_{2,0} \\
&= c_{1,0} \cdot \bar{S}(c_{1,-1})h \otimes c_2 \\
&= c_{1,0} \cdot h_3 \bar{S}(h_2) \bar{S}(c_{1,-1})h_1 \otimes c_2 \\
&= c_{1,0} \cdot h_3 \bar{S}(S(h_1)c_{1,-1}h_2) \otimes c_2 \\
&= (c_1 \cdot h)_0 \cdot \bar{S}((c_1 \cdot h)_{-1}) \otimes c_2 \\
&= g(c_1 \cdot h \otimes c_2) = g\beta(c \otimes h).
\end{aligned} \tag{5.11}$$

This shows that (5.8) is commutative and it follows that β is bijective if and only if β^τ is bijective. \square

THEOREM 5.3. *Let C/B be an H -Galois coextension and take $\tau, \lambda \in \mathcal{T}(C)$. Every left B -colinear and right H -linear coalgebra map*

$$\psi : C^\tau \longrightarrow C^\lambda \tag{5.12}$$

is of the form $\psi(c) = c_1 \cdot \nu(c_2)$, where $\nu \in \text{Hom}(C, H)$ satisfies conditions (4.1) and (4.2) of Proposition 4.1. If ψ is an isomorphism, then $\nu \in \text{Reg}(C, H)$.

PROOF. We use the notation (2.24). As in [5], we consider the map

$$\bar{\tau} = (\varepsilon \otimes 1)\beta^{-1} : C \square_B C \longrightarrow H. \tag{5.13}$$

We introduce the notation $c \diamond d = \bar{\tau}(c \otimes d)$. Then we have the following properties:

$$\varepsilon_H(c \diamond d) = \varepsilon_C(c)\varepsilon_C(d), \tag{5.14}$$

$$(c \diamond d)h = c \diamond (d \cdot h), \tag{5.15}$$

$$(c \cdot h) \diamond d = S(h)(c \diamond d), \tag{5.16}$$

$$c_1 \cdot (c_2 \diamond d) = \varepsilon(c)d. \tag{5.17}$$

The map

$$\nu : C \longrightarrow H, \quad \nu(c) = c_1 \diamond \psi(c_2), \tag{5.18}$$

satisfies the property

$$c_1 \cdot \nu(c_2) = c_1 \cdot (c_2 \diamond \psi(c_3)) = \psi(c). \tag{5.19}$$

Since ψ is a coalgebra,

$$\varepsilon_H v(c) = \varepsilon_H(c_1 \diamond \psi(c_2)) = \varepsilon(\psi(c)) = \varepsilon(c), \quad (5.20)$$

and it follows that $\varepsilon_H \circ v = \varepsilon_C$.

It follows from (5.16) and (5.17) that

$$\begin{aligned} v(c \cdot h) &= (c \cdot h)_1 \diamond \psi((c \cdot h)_2) = c_1 \cdot h_1 \diamond \psi(c_2 \cdot h_2) \\ &= S(h_1)(c_1 \diamond \psi(c_2))h_2 = S(h_1)v(c)h_2. \end{aligned} \quad (5.21)$$

Since ψ is a coalgebra map, we have that

$$\begin{aligned} \psi(c_1 \cdot c_{2,-1}) \otimes \psi(c_{2,0}) &= \psi(c_1)\psi(c_{2,-1}) \otimes \psi(c_{2,0}), \\ c_1 \cdot v(c_2)c_{3,-1} \otimes c_{3,0,1} \cdot v(c_{3,0,2}) \\ &= (c_1 \cdot v(c_2))_1(c_1 \cdot v(c_2))_{2,-1} \otimes (c_1 \cdot v(c_2))_{2,0} \\ &= c_1 \cdot v(c_3)_1(c_2 \cdot v(c_3)_2)_{-1} \otimes (c_2 \cdot v(c_3)_2)_{0,0} \\ &= c_1 \cdot c_{2,-1}v(c_3)_1 \otimes c_{2,0} \cdot v(c_3)_2, \end{aligned} \quad (5.22)$$

which is equivalent to

$$\begin{aligned} c_1 \otimes c_2 \cdot v(c_3)c_{4,-1} \otimes c_{4,0,1} \cdot v(c_{4,0,2}) \\ = c_1 \otimes c_2 \cdot c_{3,-1}v(c_4)_1 \otimes c_{3,0} \cdot v(c_4)_2. \end{aligned} \quad (5.23)$$

After we apply β^{-1} to both sides, we obtain

$$\begin{aligned} c_1 \otimes v(c_2)c_{3,-1} \otimes c_{3,0,1} \cdot v(c_{3,0,2}) &= c_1 \otimes c_{2,-1}v(c_3)_1 \otimes c_{2,0} \cdot v(c_3)_2, \\ v(c_1)c_{2,-1} \otimes c_{2,0,1} \cdot v(c_{2,0,2}) &= c_{1,-1}v(c_2)_1 \otimes c_{1,0} \cdot v(c_2)_2. \end{aligned} \quad (5.24)$$

If ψ is an isomorphism, then $\psi^{-1} : C^\lambda \rightarrow C^\tau$ is a left B -colinear and right H -linear coalgebra map. Then we have a map $w : C \rightarrow H$ satisfying (4.1) and (4.2) such that

$$\psi^{-1}(c) = c_1 \cdot w(c_2). \quad (5.25)$$

For all $c \in C$, we have that

$$c = c_1 \cdot v(c_2)w(c_3) = c_1 \cdot w(c_2)v(c_3). \quad (5.26)$$

Proceeding as in the proof of (5.24), we find that v is convolution invertible.

□

ACKNOWLEDGMENT. This research was supported by the Project G.0278.01 “Construction and Applications of Noncommutative Geometry: From Algebra to Physics” of FWO Vlaanderen, and by the bilateral Project BIL99/43 “New Computational, Geometric, and Algebraic Methods Applied to Quantum Groups and Differential Operators” of the Flemish and Chinese governments.

REFERENCES

- [1] M. Beattie, C.-Y. Chen, and J. J. Zhang, *Twisted Hopf comodule algebras*, Comm. Algebra **24** (1996), no. 5, 1759–1775.
- [2] M. Beattie and B. Torrecillas, *Twistings and Hopf Galois extensions*, J. Algebra **232** (2000), no. 2, 673–696.
- [3] R. J. Blattner, M. Cohen, and S. Montgomery, *Crossed products and inner actions of Hopf algebras*, Trans. Amer. Math. Soc. **298** (1986), no. 2, 671–711.
- [4] R. J. Blattner and S. Montgomery, *Crossed products and Galois extensions of Hopf algebras*, Pacific J. Math. **137** (1989), no. 1, 37–54.
- [5] T. Brzeziński and P. M. Hajac, *Coalgebra extensions and algebra coextensions of Galois type*, Comm. Algebra **27** (1999), no. 3, 1347–1367.
- [6] S. Caenepeel, *Harrison cocycles and the group of Galois coobjects*, Algèbre Non Commutative, Groupes Quantiques et Invariants (Reims, 1995), Sémin. Congr., vol. 2, Société Mathématique de France, Paris, 1997, pp. 83–101.
- [7] S. Caenepeel, S. Dăscălescu, G. Militaru, and F. Panaite, *Coalgebra deformations of bialgebras by Harrison cocycles, copairings of Hopf algebras and double crosscoproducts*, Bull. Belg. Math. Soc. Simon Stevin **4** (1997), no. 5, 647–671.
- [8] S. Dăscălescu, G. Militaru, and Ș. Raianu, *Crossed coproducts and cleft coextensions*, Comm. Algebra **24** (1996), no. 4, 1229–1243.
- [9] S. Dăscălescu, C. Năstăsescu, and Ș. Raianu, *Hopf Algebras. An Introduction*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 235, Marcel Dekker, New York, 2001.
- [10] S. Dăscălescu, Ș. Raianu, and Y. H. Zhang, *Finite Hopf-Galois coextensions, crossed coproducts, and duality*, J. Algebra **178** (1995), no. 2, 400–413.
- [11] Y. Doi and M. Takeuchi, *Cleft comodule algebras for a bialgebra*, Comm. Algebra **14** (1986), no. 5, 801–817.
- [12] T. X. Ju and C. R. Cai, *Twisted Hopf module coalgebras*, Comm. Algebra **28** (2000), no. 1, 307–320.
- [13] S. Montgomery, *Hopf Algebras and Their Actions on Rings*, CBMS Regional Conference Series in Mathematics, vol. 82, American Mathematical Society, Rhode Island, 1993.
- [14] M. E. Sweedler, *Cohomology of algebras over Hopf algebras*, Trans. Amer. Math. Soc. **133** (1968), 205–239.
- [15] ———, *Hopf Algebras*, Mathematics Lecture Note Series, W. A. Benjamin, New York, 1969.

S. Caenepeel: Faculty of Applied Sciences, Vrije Universiteit Brussel (VUB), B-1050 Brussels, Belgium

E-mail address: scaenepe@vub.ac.be

Dingguo Wang: Department of Mathematics, Qufu Normal University, Qufu, Shandong 273165, China

E-mail address: dgwang@qtnu.edu.cn

Yanxin Wang: Department of Mathematics, Tsinghua University, Beijing 100084, China

E-mail address: wyx01@mails.tsinghua.edu.cn

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk