

## GER-TYPE AND HYERS-ULAM STABILITIES FOR THE FIRST-ORDER LINEAR DIFFERENTIAL OPERATORS OF ENTIRE FUNCTIONS

TAKESHI MIURA, GO HIRASAWA, and SIN-EI TAKAHASI

Received 22 April 2003

*Dedicated to Professor Seiji Watanabe on his 60th birthday (Kanreki).*

Let  $h$  be an entire function and  $T_h$  a differential operator defined by  $T_h f = f' + h f$ . We show that  $T_h$  has the Hyers-Ulam stability if and only if  $h$  is a nonzero constant. We also consider Ger-type stability problem for  $|1 - f'/h f| \leq \varepsilon$ .

2000 Mathematics Subject Classification: 34K20, 26D10.

**1. Introduction.** The first result, which we now call the Hyers-Ulam stability (HUS), is due to Hyers [4] who gave an answer to a question posed by Ulam (cf. [11, Chapter VI] and [12]) in 1940 concerning the stability of homomorphisms: for what metric groups  $G$  is it true that an  $\varepsilon$ -automorphism of  $G$  is necessarily near to a strict automorphism?

An answer to the above problem has been given as follows. Suppose  $E_1$  and  $E_2$  are two real Banach spaces and  $f : E_1 \rightarrow E_2$  is a mapping such that  $f(tx)$  is continuous in  $t \in \mathbb{R}$ , the set of all real numbers, for each fixed  $x \in E_1$ . If there exist  $\theta \geq 0$  and  $p \in \mathbb{R} \setminus \{1\}$  such that

$$\|f(x+y) - f(x) - f(y)\| \leq \theta(\|x\|^p + \|y\|^p) \quad (1.1)$$

for all  $x, y \in E_1$ , then there is a unique linear mapping  $T : E_1 \rightarrow E_2$  such that  $\|f(x) - T(x)\| \leq 2\theta\|x\|^p / |2 - 2^p|$  for every  $x \in E_1$ . Hyers [4] obtained the result for  $p = 0$ . Then Rassias [7] generalized the above result of Hyers to the case where  $0 \leq p < 1$ , while the proof given in [7] also works for  $p < 0$ . Gajda [2] solved the problem for  $1 < p$  and also gave an example that a similar result does not hold for  $p = 1$  (cf. [8]).

In connection with the stability of exponential functions, Alsina and Ger [1] remarked that the differential equation  $y' = y$  has the HUS. More explicitly, suppose  $I$  is an open interval,  $\varepsilon > 0$ , and  $f : I \rightarrow \mathbb{R}$  is a differentiable function such that  $|f'(t) - f(t)| \leq \varepsilon$  for all  $t \in I$ . Then, there is a differentiable function  $g : I \rightarrow \mathbb{R}$  such that  $g' = g$  and  $|f(t) - g(t)| \leq 3\varepsilon$  for all  $t \in I$ . The third and first authors of this paper along with Miyajima [10] considered the Banach-space-valued differential equation  $y' = \lambda y$ , where  $\lambda$  is a complex constant. Then they proved the HUS of  $y' = \lambda y$  under the condition that  $\operatorname{Re} \lambda \neq 0$ . Though, they treated the result as the stability of the operator  $D - I_d$ , where  $D$  denotes the ordinary differential operator and  $I_d$  the identity. Some stability results of other differential equations (or operators) are also known (cf. [5, 6, 9]).

Taking the group structure of  $\mathbb{C} \setminus \{0\}$  into account, Ger and Šemrl [3] considered the inequality

$$\left| \frac{f(x+y)}{f(x)f(y)} - 1 \right| \leq \theta \quad (x, y \in S) \quad (1.2)$$

for a mapping  $f : S \rightarrow \mathbb{C} \setminus \{0\}$ , where  $(S, +)$  is a semigroup and  $\mathbb{C}$  is the set of all complex numbers. If  $0 \leq \theta < 1$  and if  $(S, +)$  is a cancellative abelian semigroup, then they proved that there is a unique function  $g : S \rightarrow \mathbb{C} \setminus \{0\}$  such that  $g(x+y) = g(x)g(y)$  for all  $x, y \in S$  and that

$$\max \left\{ \left| \frac{f(x)}{g(x)} - 1 \right|, \left| \frac{g(x)}{f(x)} - 1 \right| \right\} \leq \sqrt{1 + \frac{1}{(1-\theta)^2}} - 2\sqrt{\frac{1+\theta}{1-\theta}} \quad (1.3)$$

for all  $x \in S$ . The stability phenomena of this kind is called Ger-type stability.

Throughout this paper,  $H(\mathbb{C})$  stands for the set of all entire functions. Let  $h \in H(\mathbb{C})$  and  $T_h : H(\mathbb{C}) \rightarrow H(\mathbb{C})$  be a linear differential operator defined by

$$T_h f(z) = f'(z) + h(z)f(z) \quad (f \in H(\mathbb{C}), z \in \mathbb{C}). \quad (1.4)$$

**DEFINITION 1.1.** The operator  $T_h$  is said to have the HUS if and only if there exists a constant  $K \geq 0$  with the following property: to each  $\varepsilon \geq 0$  and  $f, g \in H(\mathbb{C})$  satisfying  $\sup_{z \in \mathbb{C}} |T_h f(z) - g(z)| \leq \varepsilon$ , there exists an  $f_0 \in H(\mathbb{C})$  such that  $T_h f_0 = g$  and  $\sup_{z \in \mathbb{C}} |f(z) - f_0(z)| \leq K\varepsilon$ . Such  $K$  is called an HUS constant for  $T_h$ . If, in addition, the minimum of all such  $K$ 's exists, then it is called *the* HUS constant for  $T_h$ .

In this paper, we first consider the HUS of the differential operator  $T_h$ . Then we show that  $T_h$  has the HUS if and only if  $h \in H(\mathbb{C})$  is a nonzero constant function. Moreover, we give the HUS constant for  $T_h$ . Finally, we consider the Ger-type stability problem of the differential equation  $y' = \lambda y$ . To be more explicit, suppose  $\varepsilon \geq 0$  and  $f \in H(\mathbb{C})$  satisfies

$$\sup_{z \in \mathbb{C}} \left| \frac{f'(z)}{\lambda f(z)} - 1 \right| \leq \varepsilon. \quad (1.5)$$

Does there exist  $K \geq 0$  such that

$$\sup_{z \in \mathbb{C}} \left| \frac{f(z)}{ce^{\lambda t}} - 1 \right| \leq K\varepsilon \quad \text{or} \quad \sup_{z \in \mathbb{C}} \left| \frac{ce^{\lambda t}}{f(z)} - 1 \right| \leq K\varepsilon \quad (1.6)$$

holds for some  $c \in \mathbb{C} \setminus \{0\}$ ? To this problem, we give a negative answer: the Ger-type stability does not hold in general. Moreover, we show that the solution  $f \in H(\mathbb{C})$  to the differential equation  $y' = \lambda y$  is only the function which satisfies both (1.5) and (1.6).

**2. The HUS for  $T_h$ .** For simplicity, we write  $\int_0^z f(\zeta) d\zeta$  for  $\int_0^1 f(zt) z dt$ , where  $z \in \mathbb{C}$  and  $f \in H(\mathbb{C})$ . We associate to each  $h \in H(\mathbb{C})$  a function  $\tilde{h}$  defined by

$$\tilde{h}(z) = \exp \int_0^z h(\zeta) d\zeta \quad (z \in \mathbb{C}). \quad (2.1)$$

Let  $h \in H(\mathbb{C})$ . Throughout this section,  $T_h : H(\mathbb{C}) \rightarrow H(\mathbb{C})$  denotes a linear differential operator defined by (1.4). Suppose  $f, g \in H(\mathbb{C})$ . Then note that  $T_h f = g$  if and only if  $f$  is of the form

$$f(z) = \frac{1}{\tilde{h}(z)} \left\{ f(0) + \int_0^z g(\zeta) \tilde{h}(\zeta) d\zeta \right\} \quad (z \in \mathbb{C}). \quad (2.2)$$

**LEMMA 2.1.** *Suppose  $h \in H(\mathbb{C})$  is not a constant function,  $f \in H(\mathbb{C})$ , and*

$$0 < \sup_{z \in \mathbb{C}} |T_h f(z)| < \infty. \quad (2.3)$$

*Then*

$$\sup_{z \in \mathbb{C}} \left| f(z) - \frac{c}{\tilde{h}(z)} \right| = \infty \quad (2.4)$$

*for every  $c \in \mathbb{C}$ .*

**PROOF.** By hypothesis,  $T_h f$  is a bounded entire function, and so  $T_h f$  must be constant, say  $c_0 \in \mathbb{C} \setminus \{0\}$  by Liouville's theorem. Hence, by (2.2),  $f$  is of the form

$$f(z) = \frac{1}{\tilde{h}(z)} \left\{ f(0) + c_0 \int_0^z \tilde{h}(\zeta) d\zeta \right\} \quad (z \in \mathbb{C}). \quad (2.5)$$

Suppose  $\sup_{z \in \mathbb{C}} |f(z) - c_1/\tilde{h}(z)| < \infty$  for some  $c_1 \in \mathbb{C}$ . Another application of Liouville's theorem yields the existence of a constant  $c_2 \in \mathbb{C}$  such that  $c_2 = f - c_1/\tilde{h}$ , and therefore (2.5) gives

$$c_2 \tilde{h}(z) = f(0) - c_1 + c_0 \int_0^z \tilde{h}(\zeta) d\zeta \quad (z \in \mathbb{C}). \quad (2.6)$$

By differentiating both sides of (2.6) with respect to  $z$ , we obtain

$$c_2 h \tilde{h} = c_0 \tilde{h}, \quad (2.7)$$

and hence

$$c_2 h = c_0. \quad (2.8)$$

Since  $h$  is not constant, this implies that  $c_2 = 0$ . Thus,  $f = c_1/\tilde{h}$ , and hence  $T_h f = 0$  (see (2.2)), which contradicts  $0 < \sup_{z \in \mathbb{C}} |T_h f(z)|$ .  $\square$

**THEOREM 2.2.** *If  $h \in H(\mathbb{C})$ , then each of the following statements implies the other:*

- (a)  $h$  is a nonzero constant function,
- (b)  $T_h$  has the HUS.

**PROOF.** (a)  $\Rightarrow$  (b). Suppose  $h$  is a nonzero constant function, say  $\lambda \in \mathbb{C} \setminus \{0\}$ . Then,  $\tilde{h}(z) = e^{\lambda z}$  for  $z \in \mathbb{C}$ . Suppose  $\varepsilon \geq 0$  and  $f, g \in H(\mathbb{C})$  satisfy  $\sup_{z \in \mathbb{C}} |T_h f(z) - g(z)| \leq \varepsilon$ . Then there exists a  $c_0 \in \mathbb{C}$  such that  $T_h f - g = c_0$  by Liouville's theorem. Put

$$u(z) = e^{-\lambda z} \left\{ \int_0^z g(\zeta) e^{\lambda \zeta} d\zeta \right\} \quad (z \in \mathbb{C}). \quad (2.9)$$

Then  $T_h u = g$ , and so  $T_h(f - u) = c_0$ ,  $|c_0| \leq \varepsilon$ . Hence, by (2.2),  $f$  is of the form

$$\begin{aligned} f(z) &= u(z) + \frac{1}{\tilde{h}(z)} \left\{ f(0) - u(0) + c_0 \int_0^z \tilde{h}(\zeta) d\zeta \right\} \\ &= \frac{c_0}{\lambda} + u(z) + \left( f(0) - u(0) - \frac{c_0}{\lambda} \right) e^{-\lambda z} \end{aligned} \quad (2.10)$$

for all  $z \in \mathbb{C}$ . Put

$$f_0(z) = u(z) + \left( f(0) - u(0) - \frac{c_0}{\lambda} \right) e^{-\lambda z} \quad (z \in \mathbb{C}), \quad (2.11)$$

then  $T_h f_0 = g$  and

$$|f(z) - f_0(z)| = \left| \frac{c_0}{\lambda} \right| \leq \frac{\varepsilon}{|\lambda|} \quad (2.12)$$

for every  $z \in \mathbb{C}$  so that  $T_h$  has the HUS with an HUS constant  $1/|\lambda|$ .

(b)  $\Rightarrow$  (a). Put

$$f_1(z) = \frac{1}{\tilde{h}(z)} \int_0^z \tilde{h}(\zeta) d\zeta \quad (z \in \mathbb{C}). \quad (2.13)$$

Then we obtain  $T_h f_1 = 1$ . Let  $K < \infty$  be an HUS constant for  $T_h$ . Since  $T_h$  has the HUS, there is an  $f_2 \in H(\mathbb{C})$ , such that  $T_h f_2 = 0$  and

$$\sup_{z \in \mathbb{C}} |f_1(z) - f_2(z)| \leq K. \quad (2.14)$$

Note that  $f_2$  is of the form  $f_2(z) = f_2(0)/\tilde{h}(z)$  for all  $z \in \mathbb{C}$ , since  $T_h f_2 = 0$ . **Lemma 2.1**, applied to  $f_1$ , yields that  $h$  is a constant function. If  $h$  were 0, then (2.13) would be written in the form  $f_1(z) = z$  for  $z \in \mathbb{C}$ , and hence from (2.14),  $\sup_{z \in \mathbb{C}} |z - f_2(0)| \leq K < \infty$ , which is a contradiction. Thus, we conclude that  $h$  is a nonzero constant function.  $\square$

**THEOREM 2.3.** *Suppose  $\lambda \in \mathbb{C} \setminus \{0\}$ ,  $f, g \in H(\mathbb{C})$ , and  $\sup_{z \in \mathbb{C}} |T_\lambda f(z) - g(z)| < \infty$ . Then there exists a unique  $f_0 \in H(\mathbb{C})$  such that  $T_\lambda f_0 = g$  and*

$$\sup_{z \in \mathbb{C}} |f(z) - f_0(z)| < \infty. \quad (2.15)$$

Furthermore,  $1/|\lambda|$  is the HUS constant for  $T_\lambda$ .

**PROOF.** The existence of such a function  $f_0 \in H(\mathbb{C})$  is proved by **Theorem 2.2**, and so we need to show only the uniqueness. Suppose  $f_1 \in H(\mathbb{C})$  and  $f_2 \in H(\mathbb{C})$  satisfy  $T_\lambda f_j = g$  and

$$\sup_{z \in \mathbb{C}} |f(z) - f_j(z)| < \infty \quad (2.16)$$

for  $j = 1, 2$ . Since  $T_\lambda f_j = g$ ,

$$f_j(z) = e^{-\lambda z} \left\{ f_j(0) + \int_0^z g(\zeta) e^{\lambda \zeta} d\zeta \right\} \quad (z \in \mathbb{C}) \quad (2.17)$$

for  $j = 1, 2$ , and hence

$$f_1(z) - f_2(z) = (f_1(0) - f_2(0))e^{-\lambda z} \quad \forall z \in \mathbb{C}. \quad (2.18)$$

It follows from (2.16) that  $f_1 - f_2$  is constant by Liouville's theorem. Therefore,  $f_1(0) = f_2(0)$  by (2.18), which implies that  $f_1 = f_2$ , proving the uniqueness.

We show that  $1/|\lambda|$  is the HUS constant for  $T_\lambda$ . Indeed,  $1/|\lambda|$  is an HUS constant by (2.12). Conversely, let  $K$  be an arbitrary HUS constant for  $T_\lambda$ , and put

$$f_2(z) = \frac{1}{\lambda} - \frac{1}{\lambda}e^{-\lambda z} \quad (z \in \mathbb{C}). \quad (2.19)$$

A simple calculation shows that  $f_2'(z) + \lambda f_2(z) = 1$  for all  $z \in \mathbb{C}$ , and hence  $\sup_{z \in \mathbb{C}} |T_\lambda f_2(z)| = 1$ . Then, there exists an  $f_3 \in H(\mathbb{C})$  such that  $T_\lambda f_3 = 0$  and  $\sup_{z \in \mathbb{C}} |f_2(z) - f_3(z)| \leq K$ . Since  $|f_2(z) + \lambda^{-1}e^{-\lambda z}| = 1/|\lambda|$  for  $z \in \mathbb{C}$ , the uniqueness implies that  $f_3(z) = -\lambda^{-1}e^{-\lambda z}$ , which proves  $1/|\lambda| \leq K$ . Thus,  $1/|\lambda|$  is the HUS constant for  $T_\lambda$ .  $\square$

**3. Stability for the Ger-type differential inequality.** In this section, we consider the Ger-type stability problem. First, we show that the Ger-type stability does not hold in general. Indeed, the following proposition is true.

**PROPOSITION 3.1.** *For  $\lambda \in \mathbb{C} \setminus \{0\}$  and  $\varepsilon > 0$ , there exists an  $f \in H(\mathbb{C})$  with the following properties:*

$$\begin{aligned} \sup_{z \in \mathbb{C}} \left| \frac{f'(z)}{\lambda f(z)} - 1 \right| &\leq \varepsilon, \\ \sup_{z \in \mathbb{C}} \left| \frac{f(z)}{ce^{\lambda z}} - 1 \right| &= \sup_{z \in \mathbb{C}} \left| \frac{ce^{\lambda z}}{f(z)} - 1 \right| = \infty \quad \forall c \in \mathbb{C} \setminus \{0\}. \end{aligned} \quad (3.1)$$

**PROOF.** We associate to each  $\lambda \in \mathbb{C} \setminus \{0\}$  and  $\varepsilon > 0$  a function  $f$  defined by

$$f(z) = e^{(\lambda + |\lambda|\varepsilon)z} \quad (z \in \mathbb{C}). \quad (3.2)$$

As above, we obtain

$$f'(z) = (\lambda + |\lambda|\varepsilon)f(z) \quad (z \in \mathbb{C}), \quad (3.3)$$

so that

$$\left| \frac{f'(z)}{\lambda f(z)} - 1 \right| = \varepsilon \quad \forall z \in \mathbb{C}. \quad (3.4)$$

If  $c \in \mathbb{C} \setminus \{0\}$ , then we have

$$\begin{aligned} \left| \frac{f(z)}{ce^{\lambda z}} - 1 \right| &\geq \frac{1}{|c|} |e^{|\lambda|\varepsilon z}| - 1 \rightarrow \infty \quad (\operatorname{Re} z \rightarrow \infty), \\ \left| \frac{ce^{\lambda z}}{f(z)} - 1 \right| &\geq |c| |e^{-|\lambda|\varepsilon z}| - 1 \rightarrow \infty \quad (\operatorname{Re} z \rightarrow -\infty), \end{aligned} \quad (3.5)$$

and so

$$\sup_{z \in \mathbb{C}} \left| \frac{f(z)}{ce^{\lambda z}} - 1 \right| = \sup_{z \in \mathbb{C}} \left| \frac{ce^{\lambda z}}{f(z)} - 1 \right| = \infty \quad \forall c \in \mathbb{C} \setminus \{0\}. \quad (3.6)$$

□

One might ask when the Ger-type stability is true. We give an answer to this question. If the Ger-type stability holds, then the function  $f \in H(\mathbb{C})$  must be of the form  $f(z) = f(0)e^{\lambda z}$ . That is, the only solution to the differential equation  $y' = \lambda y$  has the Ger-type stability.

**THEOREM 3.2.** *Suppose  $\lambda \in \mathbb{C} \setminus \{0\}$ ,  $\varepsilon > 0$ , and  $f \in H(\mathbb{C})$  satisfies  $f(z) \neq 0$  for all  $z \in \mathbb{C}$  and (1.5) holds. Suppose*

$$\sup_{z \in \mathbb{C}} \left| \frac{f(z)}{ce^{\lambda z}} - 1 \right| \quad \text{or} \quad \sup_{z \in \mathbb{C}} \left| \frac{ce^{\lambda z}}{f(z)} - 1 \right| \quad (3.7)$$

*is finite for some  $c \in \mathbb{C} \setminus \{0\}$ ; then  $f$  is of the form  $f(z) = f(0)e^{\lambda z}$  for all  $z \in \mathbb{C}$ .*

**PROOF.** It follows from (1.5) that  $1 - f'/\lambda f$  is constant, say  $c_0 \in \mathbb{C}$ , by Liouville's theorem. Thus,  $f' = (1 - c_0)\lambda f$ , and hence

$$f(z) = f(0)e^{(1-c_0)\lambda z} \quad (z \in \mathbb{C}). \quad (3.8)$$

Suppose that there is a  $c_1 \in \mathbb{C} \setminus \{0\}$  such that

$$\sup_{z \in \mathbb{C}} \left| \frac{f(z)}{c_1 e^{\lambda z}} - 1 \right| < \infty. \quad (3.9)$$

From (3.8), it follows that

$$\sup_{z \in \mathbb{C}} \left| \frac{f(0)}{c_1} e^{-c_0 \lambda z} - 1 \right| < \infty, \quad (3.10)$$

and hence  $c_0$  must be 0, proving  $f(z) = f(0)e^{\lambda z}$  for all  $z \in \mathbb{C}$ .

Similarly, we can treat the case where

$$\sup_{z \in \mathbb{C}} \left| \frac{c_2 e^{\lambda z}}{f(z)} - 1 \right| < \infty \quad (3.11)$$

for some  $c_2 \in \mathbb{C} \setminus \{0\}$ , and so the proof is omitted. □

Thus far, we have treated entire functions. Finally, we consider the Ger-type stability problem in the category of holomorphic functions on a bounded region.

**THEOREM 3.3.** *Let  $0 \in \Omega$  be a bounded convex region of  $\mathbb{C}$  and put  $M = \sup_{z \in \Omega} |z|$ . Suppose  $\lambda \in \mathbb{C} \setminus \{0\}$ ,  $0 \leq \varepsilon \leq 1$ , and  $f : \Omega \rightarrow \mathbb{C}$  is holomorphic such that  $f(z) \neq 0$  for all  $z \in \Omega$  and*

$$\sup_{z \in \Omega} \left| \frac{f'(z)}{\lambda f(z)} - 1 \right| \leq \varepsilon. \quad (3.12)$$

Then there are  $K_\lambda > 0$  and  $c \in \mathbb{C} \setminus \{0\}$  such that

$$\max \left\{ \sup_{z \in \Omega} \left| \frac{f(z)}{ce^{\lambda z}} - 1 \right|, \sup_{z \in \Omega} \left| \frac{ce^{\lambda z}}{f(z)} - 1 \right| \right\} \leq K_\lambda \varepsilon. \quad (3.13)$$

**PROOF.** Put  $g(z) = -1 + f'(z)/\lambda f(z)$  for  $z \in \Omega$ , and so

$$f'(z) = \lambda(1 + g(z))f(z) \quad (z \in \Omega). \quad (3.14)$$

From (3.14), it follows that

$$f(z) = f(0)e^{\lambda z} \exp \int_0^z \lambda g(\zeta) d\zeta \quad (3.15)$$

for every  $z \in \Omega$ , and hence

$$\begin{aligned} \left| \frac{f(z)}{f(0)e^{\lambda z}} - 1 \right| &= \left| \exp \int_0^z \lambda g(\zeta) d\zeta - 1 \right| \leq \sum_{n=1}^{\infty} \frac{1}{n!} \left| \int_0^z \lambda g(\zeta) d\zeta \right|^n \\ &\leq \sum_{n=1}^{\infty} \frac{|\lambda \varepsilon z|^n}{n!} \leq (e^{|\lambda| M} - 1) \varepsilon \end{aligned} \quad (3.16)$$

for all  $z \in \Omega$ . Similarly, we can show that

$$\sup_{z \in \Omega} \left| \frac{f(0)e^{\lambda z}}{f(z)} - 1 \right| \leq (e^{|\lambda| M} - 1) \varepsilon, \quad (3.17)$$

and so the proof is complete.  $\square$

**ACKNOWLEDGMENT.** The first and third authors are partially supported by the Grant-in-Aid for Scientific Research, Japan Society for the Promotion of Science.

## REFERENCES

- [1] C. Alsina and R. Ger, *On some inequalities and stability results related to the exponential function*, J. Inequal. Appl. **2** (1998), no. 4, 373–380.
- [2] Z. Gajda, *On stability of additive mappings*, Int. J. Math. Math. Sci. **14** (1991), no. 3, 431–434.
- [3] R. Ger and P. Šemrl, *The stability of the exponential equation*, Proc. Amer. Math. Soc. **124** (1996), no. 3, 779–787.
- [4] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Natl. Acad. Sci. USA **27** (1941), 222–224.
- [5] T. Miura, S. Miyajima, and S.-E. Takahasi, *A characterization of Hyers-Ulam stability of first order linear differential operators*, J. Math. Anal. Appl. **286** (2003), no. 1, 136–146.
- [6] ———, *Hyers-Ulam stability of linear differential operator with constant coefficients*, Math. Nachr. **258** (2003), 90–96.
- [7] T. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), no. 2, 297–300.
- [8] T. M. Rassias and P. Šemrl, *On the behavior of mappings which do not satisfy Hyers-Ulam stability*, Proc. Amer. Math. Soc. **114** (1992), no. 4, 989–993.
- [9] H. Takagi, T. Miura, and S.-E. Takahasi, *Essential norms and stability constants of weighted composition operators on  $C(X)$* , Bull. Korean Math. Soc. **40** (2003), no. 4, 583–591.

- [10] S.-E. Takahasi, T. Miura, and S. Miyajima, *On the Hyers-Ulam stability of the Banach space-valued differential equation  $y' = \lambda y$* , Bull. Korean Math. Soc. **39** (2002), no. 2, 309–315.
- [11] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Tracts in Pure and Applied Mathematics, no. 8, Interscience Publishers, New York, 1960.
- [12] ———, *Sets, Numbers, and Universes: Selected Works*, The MIT Press, Massachusetts, 1974.

Takeshi Miura: Department of Basic Technology, Applied Mathematics and Physics, Yamagata University, Yonezawa 992-8510, Japan

*E-mail address:* [miura@yz.yamagata-u.ac.jp](mailto:miura@yz.yamagata-u.ac.jp)

Go Hirasawa: Department of Mathematics, Nippon Institute of Technology, Miyashiro, Saitama 345-8501, Japan

*E-mail address:* [hirasawa1@muh.biglobe.ne.jp](mailto:hirasawa1@muh.biglobe.ne.jp)

Sin-Ei Takahasi: Department of Basic Technology, Applied Mathematics and Physics, Yamagata University, Yonezawa 992-8510, Japan

*E-mail address:* [sin-ei@emperor.yz.yamagata-u.ac.jp](mailto:sin-ei@emperor.yz.yamagata-u.ac.jp)

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)