

## COMPLETELY MULTIPLICATIVE FUNCTIONS ARISING FROM SIMPLE OPERATIONS

VICHIAN LAOHAKOSOL and NITTIYA PABHAPOTE

Received 13 April 2003

Given two multiplicative arithmetic functions, various conditions for their convolution, powers, and logarithms to be completely multiplicative, based on values at the primes, are derived together with their applications.

2000 Mathematics Subject Classification: 11A25.

**1. Introduction.** By an *arithmetic function* we mean a complex-valued function whose domain is the set of positive integers  $\mathbb{N}$ . We define the addition and convolution of two arithmetic functions  $f$  and  $g$ , respectively, by

$$(f + g)(n) = f(n) + g(n), \quad (f * g)(n) = \sum_{ij=n} f(i)g(j). \quad (1.1)$$

It is well known (see, e.g., [2, 6, 15, 16]) that the set  $(\mathcal{A}, +, *)$  of all arithmetic functions is a unique factorization domain with the arithmetic function

$$I(n) = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{otherwise,} \end{cases} \quad (1.2)$$

being its convolution identity.

A nonzero arithmetic function  $f \in \mathcal{A}$  is called *multiplicative* if and only if

$$f(mn) = f(m)f(n) \quad \text{whenever } (m, n) = 1. \quad (1.3)$$

It is called *completely multiplicative* if this equality holds for all  $m, n \in \mathbb{N}$ .

Denote by  $\mathcal{M}$  the set of all multiplicative functions and by  $\mathcal{C}$  the set of all completely multiplicative functions. It is easy to see that the elements of  $\mathcal{M}$  are uniquely determined via their values at all prime powers, while those of  $\mathcal{C}$  are uniquely determined only via their values at the primes. Unlike  $\mathcal{M}$ , which is a group with respect to convolution,  $\mathcal{C}$  is closed neither with respect to addition nor convolution.

A natural problem is that of characterizing  $\mathcal{M}$  and  $\mathcal{C}$  by various means and there have appeared a number of such investigations, for example, by Apostol [1, 2], Carroll [5], Laohakosol et al. [10, 11], and Rearick [12, 13].

In this paper, we propose the naive problem: given  $f, g \in \mathcal{M}$ , find (simple) necessary and sufficient condition(s) for their convolution, powers, and logarithms to be in  $\mathcal{C}$ .

Though the problem seems elementary, we have not been able to locate its complete solution. As we will see, several of the techniques used to resolve certain cases of this problem are not elementary, that is, making use of logarithmic operators (see [10, 11, 13]).

The question about sum is trivial because by looking at the values at  $n = 1$  we see immediately that a sum of two multiplicative functions is never multiplicative.

To facilitate later discussions, we recall some basic facts about logarithmic operators on  $\mathcal{A}$ . For  $f \in \mathcal{A}$ , its Rearick logarithm, denoted by  $\text{Log } f \in \mathcal{A}$ , is defined by

$$\begin{aligned} (\text{Log } f)(1) &= \log f(1) \quad (\text{assuming } f(1) > 0), \\ (\text{Log } f)(n) &= \frac{1}{\log n} \sum_{d|n} f(d) f^{-1}\left(\frac{n}{d}\right) \log d \\ &= \frac{1}{\log n} (df * f^{-1})(n) \quad (n > 1), \end{aligned} \tag{1.4}$$

where  $df(n) = f(n) \log n$  denotes the log-derivation of  $f$ .

In 1973, Carlitz and Subbarao [4] defined another logarithmic operator of  $f$ , denoted by  $\beta f \in \mathcal{A}$ , as follows:

$$\begin{aligned} (\beta f)(1) &= 1 \quad (\text{assuming } f(1) = 1), \\ (\beta f)(n) &= \sum_{r=1}^{\infty} \frac{(-1)^{r-1}}{r} \sum_{\substack{a_1 \cdots a_r = n \\ a_i \neq 1 \ (i=1, \dots, r)}} f(a_1) \cdots f(a_r) \quad (n > 1). \end{aligned} \tag{1.5}$$

As proved in [10], these two logarithms are

- (i) essentially identical, that is, both take the same values except only at  $n = 1$ ,
- (ii) both bijective,
- (iii) both transform convolution into sum.

Indeed, as shown by Carlitz and Subbarao [4], the inverse (with respect to convolution) of  $\beta$  is the operator  $\gamma: \mathcal{U}_1 \rightarrow \mathcal{U}_1$ , where  $\mathcal{U}_1$  is the set of all arithmetic functions  $f$  such that  $f(1) = 1$ , defined by

$$\begin{aligned} (\gamma g)(1) &= 1, \\ (\gamma g)(n) &= \sum_{r=1}^{\infty} \frac{1}{r!} \sum_{\substack{a_1 \cdots a_r = n \\ a_i \neq 1 \ (i=1, \dots, r)}} g(a_1) \cdots g(a_r) \quad (n > 1). \end{aligned} \tag{1.6}$$

**2. Convolution.** We first look at the case of convolution.

**THEOREM 2.1.** *Let  $f, g \in \mathcal{M}$ . Then  $f * g \in \mathcal{C} \Leftrightarrow$  either*

$$\begin{aligned} g(p^a) &= f^{-1}(p^a) + (f(p) + g(p))f^{-1}(p^{a-1}) \\ &\quad + \cdots + (f(p) + g(p))^{a-1}f^{-1}(p) + (f(p) + g(p))^a \end{aligned} \tag{2.1}$$

or

$$\begin{aligned} f(p^a) &= g^{-1}(p^a) + (f(p) + g(p))g^{-1}(p^{a-1}) \\ &\quad + \cdots + (f(p) + g(p))^{a-1}g^{-1}(p) + (f(p) + g(p))^a \end{aligned} \quad (2.2)$$

for all primes  $p$  and all  $a \in \mathbb{N}$ .

**PROOF.** Writing  $g = (f * g) * f^{-1}$  and evaluating at prime powers, we get

$$\begin{aligned} g(p^a) &= f^{-1}(p^a)(f * g)(1) + f^{-1}(p^{a-1})(f * g)(p) \\ &\quad + \cdots + f^{-1}(1)(f * g)(p^a). \end{aligned} \quad (2.3)$$

If  $f * g \in \mathcal{C}$ , then  $(f * g)(p^i) = (f * g)(p)^i = (f(p) + g(p))^i$ , and (2.1) results immediately. Further, (2.2) follows by interchanging  $f$  and  $g$  in the above arguments.

On the other hand, if (2.1) holds, then comparing (2.1) and (2.3) for successive values of  $a = 1, 2, 3, \dots$ , we conclude that

$$(f * g)(p^a) = (f * g)(p)^a, \quad (2.4)$$

and so

$$(f * g) \in \mathcal{C}. \quad (2.5)$$

The same result holds similarly for (2.2).  $\square$

Two remarkable consequences are now in order.

**COROLLARY 2.2.** *Let  $f \in \mathcal{C}$  and  $g \in \mathcal{M}$ . Then*

$$f * g \in \mathcal{C} \iff g(p^a) = g(p)(g(p) + f(p))^{a-1} \quad (2.6)$$

for all primes  $p$  and all  $a \in \mathbb{N}$ .

**PROOF.** This is immediate from (2.1) in [Theorem 2.1](#), noting that  $f^{-1}(p) = -f(p)$  and

$$f \in \mathcal{C} \iff f^{-1}(p^i) = 0 \quad \forall i \geq 2. \quad (2.7) \quad \square$$

**COROLLARY 2.3.** *Let  $f, g \in \mathcal{C}$ . Then*

$$f * g \in \mathcal{C} \iff f(p)g(p) = 0 \quad (2.8)$$

for all primes  $p$ .

**PROOF.** From [Corollary 2.2](#), we have

$$\begin{aligned} f * g \in \mathcal{C} &\implies g(p^2) = g(p)(g(p) + f(p)) \\ &\implies f(p)g(p) = 0, \end{aligned} \quad (2.9)$$

while

$$\begin{aligned} f(p)g(p) = 0 &\Rightarrow g(p^a) = g(p)(g(p) + f(p))^{a-1} \\ &\Rightarrow f * g \in \mathcal{C}. \end{aligned} \tag{2.10} \quad \square$$

**3. Powers.** We first recall some of the results in [13]. For  $h \in \mathcal{A}$ , the (Rearick) exponential  $\text{Exp} h$  is defined as the unique element

$$f \in \mathcal{A}, \quad f(1) > 0, \quad \text{such that } h = \text{Log } f. \tag{3.1}$$

For  $f \in \mathcal{A}$ ,  $f(1) > 0$ , and  $r \in \mathbb{R}$ , the  $r$ th power function  $f^r$  is defined as

$$f^r = \text{Exp}(r \text{Log } f). \tag{3.2}$$

From [12], we know that if  $f \in \mathcal{M}$ ,  $r \in \mathbb{R}$ , then  $f^r \in \mathcal{M}$ , which also implies that if  $f^r \in \mathcal{M}$ ,  $r \in \mathbb{R} \setminus \{0\}$ , is multiplicative, then  $f$  is also multiplicative.

Recall also the Hsu's generalized Möbius function (see [3, 8, 11])

$$\mu_\alpha(n) = \prod_{p|n} \binom{\alpha}{\nu_p(n)} (-1)^{\nu_p(n)}, \tag{3.3}$$

where  $\alpha \in \mathbb{R}$  and  $\nu_p(n)$  is the highest power of the prime  $p$  dividing  $n$ .

**THEOREM 3.1.** *Let  $f \in \mathcal{M}$  and  $r \in \mathbb{R} \setminus \{0\}$ . Then*

$$f^r \in \mathcal{C} \iff f(p^a) = \binom{-\frac{1}{r}}{a} (-r)^a f(p)^a \tag{3.4}$$

for all primes  $p$  and all  $a \in \mathbb{N}$ .

**PROOF.** Assume  $f^r \in \mathcal{C}$ . By Haukkanen's theorem [8] and the property of power function, we have

$$\mu_{-1/r} f^r = (f^r)^{1/r} = f, \tag{3.5}$$

so that

$$f(p^a) = \mu_{-1/r}(p^a) f^r(p^a) = \binom{-\frac{1}{r}}{a} (-1)^a f^r(p)^a. \tag{3.6}$$

By the result of Carroll [5], see also [10], for  $f^r \in \mathcal{C}$ , we have

$$\begin{aligned} f^r(p) &= \text{Log } f^r(p) = r \text{Log } f(p) \\ &= \frac{r}{\log p} [f(1)f^{-1}(p) \log 1 + f(p)f^{-1}(1) \log p] = rf(p), \end{aligned} \tag{3.7}$$

and (3.4) follows.

Conversely, assuming (3.4), to show  $f^r \in \mathcal{C}$ , we use a characterization of Apostol [1], namely,

$$f^r \in \mathcal{C} \iff f^{-r}(p^a) = 0 \quad (3.8)$$

for all primes  $p$  and all integers  $a \geq 2$ .

Writing  $g = f^{-r}$ , then  $\text{Log } g = -r \text{Log } f$ . We will show that  $g(p^a) = 0$  for all  $a \geq 0$ . Since

$$(\text{Log } f)(n) = \frac{1}{\log n} (df * f^{-1})(n) \quad (n > 1), \quad (3.9)$$

then

$$f * dg = -r(g * df). \quad (3.10)$$

Thus

$$\sum_{ij=p^a} f(i)dg(j) = -r \sum_{ij=p^a} g(i)df(j). \quad (3.11)$$

Taking  $a = 1$ , we get  $g(p) = -rf(p)$ .

Taking  $a = 2$  and using  $g(p) = -rf(p)$ , we get

$$2g(p^2) = -r[2f(p^2) - (r+1)f(p)^2] = 0. \quad (3.12)$$

By induction, using (3.4), for  $a \geq 2$ , we deduce that  $g(p^a) = 0$ .  $\square$

**REMARK 3.2.** An alternative proof of the “only if” part of Theorem 3.1 can be done as follows. Assuming (3.4), to show  $f^r \in \mathcal{C}$ , we use a characterization of Carroll [5], namely, for  $f \in \mathcal{M}$ ,  $f \in \mathcal{C}$  if and only if

$$(\text{Log } f)(n) = \begin{cases} \frac{f(p)^a}{a} & \text{if } n = p^a, p \text{ prime, } a \geq 1, \\ 0 & \text{otherwise.} \end{cases} \quad (3.13)$$

From the hypothesis and using induction on  $a$ , we easily verify that

$$f^{-1}(p^a) = \left( \frac{1}{r} \right) \left( \frac{1}{a} \right) (-r)^a f(p)^a \quad (3.14)$$

for all primes  $p$  and all  $a \in \mathbb{N}$ .

Next we want to show that  $r(\text{Log } f)(p^a) = (rf(p))^a/a$ . From the definition of Rearick logarithm  $\text{Log } f$ , we have

$$\begin{aligned} (\text{Log } f)(p^a) &= \frac{1}{a} [f(p)f^{-1}(p^{a-1}) + 2f(p^2)f^{-1}(p^{a-2}) \\ &\quad + \dots + (a-1)f(p^{a-1})f^{-1}(p) + af(p^a)], \end{aligned} \quad (3.15)$$

and so, using (3.14), we get

$$(\text{Log } f)(p^a) = \frac{r^a}{a} f(p)^a \left[ \sum_{i=1}^a i \left( -\frac{1}{r} \right) \left( \frac{1}{r} \right) (-1)^a \right]. \quad (3.16)$$

Using Riordan [14, identity (3d), page 10], we deduce that

$$(\text{Log } f)(p^a) = \frac{(rf(p))^a}{a} \cdot \frac{1}{r}, \quad (3.17)$$

and the desired result follows.

**COROLLARY 3.3.** *Let  $f \in \mathcal{C}$  and  $r \in \mathbb{R} \setminus \{0, 1\}$ . Then*

$$f^r \in \mathcal{C} \iff f = I. \quad (3.18)$$

**PROOF.** For all primes  $p$  and all  $a \in \mathbb{N}$ , by Haukkanen's theorem [8] and [Theorem 3.1](#), we have

$$\begin{aligned} f^r \in \mathcal{C} &\Rightarrow (\mu_{-r} f)(p^a) = f^r(p^a) = (f^r(p))^a \\ &\Rightarrow \binom{r+a-1}{a} f(p^a) = (rf(p))^a \\ &\Rightarrow \left[ \binom{r+a-1}{a} - r^a \right] f(p)^a = 0. \end{aligned} \quad (3.19)$$

Taking  $a = 2$ , since  $r \neq 0, 1$ , we get  $f(p) = 0$  for all primes  $p$ . Being in  $\mathcal{C}$ , this yields  $f = I$ .

If  $f = I \in \mathcal{C}$ , by Haukkanen's theorem [8], we get

$$I^r(n) = (\mu_{-r} I)(n) = \mu_{-r}(n) I(n) = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{otherwise,} \end{cases} \quad (3.20)$$

and so  $I^r \in \mathcal{C}$ . □

**REMARK 3.4.** [Corollary 3.3](#) is false if the assumption  $f \in \mathcal{C}$  is dropped as seen from the following two examples.

(1) Take  $f$  as the Möbius function  $\mu \neq I$  and  $r = -1$ . We know that (see [2])  $\mu^{-1}$  is the unit function  $u$ ,  $u(n) = 1$  ( $n \geq 1$ ), that is,  $\mu^{-1} \in \mathcal{C}$ .

(2) Take  $f = |\mu| = \mu^2 \neq I$  and  $r = -1$ . Hence,  $|\mu|^{-1}$  is the Liouville function  $\lambda \in \mathcal{C}$  (see [2]).

**4. Logarithms.** Since any two of the three known logarithms are the same except at  $n = 1$ , then we may use any of them whichever appropriate. We start by observing a trivial fact that if  $f$  is multiplicative, then its Rearick logarithm  $\text{Log } f$  is not multiplicative because  $(\text{Log } f)(1) = \log f(1) = 0$ .

A natural question to consider here is that of finding a necessary and sufficient condition for the Carlitz-Subbarao logarithm  $\beta f$  to be (completely) multiplicative for a given  $f \in \mathcal{A}$ . This proves to be a difficult task, and we have the following partial result.

**THEOREM 4.1.** Let  $f \in \mathcal{A}$  with  $f(1) = 1$ . Then its Carlitz-Subbarao logarithm  $\beta f \in \mathcal{C} \Leftrightarrow$  for distinct primes  $p_1, \dots, p_k$  and positive integers  $n_1, \dots, n_k$ , there holds

$$f(p_1^{n_1} \cdots p_k^{n_k}) = \alpha(n_1, \dots, n_k) f(p_1)^{n_1} \cdots f(p_k)^{n_k}, \quad (4.1)$$

where

$$\alpha(n_1, \dots, n_k) = \sum_{r=1}^{\infty} \frac{1}{r!} I_r(n_1, \dots, n_k), \quad (4.2)$$

$I_r(n_1, \dots, n_k) = \sum^* 1$ ;  $\sum^*$  denotes the sum taken over nonnegative integers  $n_{11}, \dots, n_{r1}; \dots; n_{1k}, \dots, n_{rk}$  such that

$$\begin{aligned} n_{11} + \cdots + n_{r1} &= n_1, \dots, n_{1k} + \cdots + n_{rk} = n_k, \\ n_{11} + \cdots + n_{1k} &> 0, \dots, n_{r1} + \cdots + n_{rk} > 0. \end{aligned} \quad (4.3)$$

**PROOF.** Put  $g = \beta f$ . Then  $g(1) = \beta f(1) = 1$ . Assuming  $g = \beta f \in \mathcal{C}$ , by [4, Theorem 4.28 and equation (1.2)] of Carlitz and Subbarao, we see that for  $m > 1$ ,

$$f(m) = \sum_{r=1}^{\infty} \frac{1}{r!} \sum_{\substack{a_1 \cdots a_r = m \\ a_i \neq 1 (i=1, \dots, r)}} g(a_1) \cdots g(a_r) = g(m) c(m), \quad (4.4)$$

where

$$c(m) = \sum_{r=1}^{\infty} \frac{1}{r!} \sum_{\substack{a_1 \cdots a_r = m \\ a_i \neq 1 (i=1, \dots, r)}} 1. \quad (4.5)$$

Writing  $m$  as a unique prime factorization,  $m = p_1^{n_1} \cdots p_k^{n_k}$ , we have

$$\sum_{\substack{a_1 \cdots a_r = m \\ a_i \neq 1 (i=1, \dots, r)}} 1 = \sum^* 1. \quad (4.6)$$

The sum  $\sum^*$  depends only on  $r, n_1, \dots, n_k$ , but is independent of  $p_1, \dots, p_k$ ; we call it  $I_r(n_1, \dots, n_k)$ . Thus the function

$$c(p_1^{n_1} \cdots p_k^{n_k}) = \sum_{r=1}^{\infty} \frac{1}{r!} I_r(n_1, \dots, n_k) \quad (4.7)$$

depends only on  $r$  and  $n_1, \dots, n_k$ , but is independent of  $p_1, \dots, p_k$ , so that we may rewrite it as  $\alpha(n_1, \dots, n_k)$ . Such a function  $c(m)$  is known as prime-independent arithmetic function (see [9, page 33]). Since  $f(p) = g(p)c(p) = g(p)$ , then

$$\begin{aligned} f(p_1^{n_1} \cdots p_k^{n_k}) &= \alpha(n_1, \dots, n_k) g(p_1)^{n_1} \cdots g(p_k)^{n_k} \\ &= \alpha(n_1, \dots, n_k) f(p_1)^{n_1} \cdots f(p_k)^{n_k}, \end{aligned} \quad (4.8)$$

as to be proved.

Conversely, from  $g = \beta f$ , we get (see [4])  $f = \gamma g$ ; that is,

$$\begin{aligned} & \sum_{r=1}^{\infty} \frac{1}{r!} \sum_{\substack{a_1 \cdots a_r = p_1^{n_1} \cdots p_k^{n_k} \\ a_i \neq 1 (i=1, \dots, r)}} g(a_1) \cdots g(a_r) \\ &= f(p_1^{n_1} \cdots p_k^{n_k}) = \alpha(n_1, \dots, n_k) f(p_1)^{n_1} \cdots f(p_k)^{n_k}. \end{aligned} \quad (4.9)$$

Specializing  $k = 1 = n_1$ , we get  $g(p) = f(p)$ .

Thus

$$\begin{aligned} & \sum_{r=1}^{\infty} \frac{1}{r!} \sum_{\substack{a_1 \cdots a_r = p_1^{n_1} \cdots p_k^{n_k} \\ a_i \neq 1 (i=1, \dots, r)}} g(a_1) \cdots g(a_r) \\ &= \alpha(n_1, \dots, n_k) g(p_1)^{n_1} \cdots g(p_k)^{n_k}. \end{aligned} \quad (4.10)$$

We now show that  $g(p^i) = g(p)^i$  for each prime  $p$  and  $i \in \mathbb{N}$ .

Taking  $k = 1$  and  $n_1 = 2$  in (4.10), observing that  $\alpha(2) = \sum_{r=1}^{\infty} (1/r!) I_r(2)$ , we have

$$\frac{1}{1!} g(p^2) + \frac{1}{2!} g(p)^2 = \alpha(2) g(p)^2 = \left( \frac{1}{1!} + \frac{1}{2!} \right) g(p)^2, \quad (4.11)$$

that is,  $g(p^2) = g(p)^2$ .

Assuming  $g(p^j) = g(p)^j$  for all  $j < i$ , and using induction in (4.10), we obtain

$$\frac{1}{1!} g(p^i) + g(p)^i \sum_{r=2}^{\infty} \frac{1}{r!} I_r(i) = \alpha(i) g(p)^i. \quad (4.12)$$

As observed earlier,  $\alpha(i)$  is just the sum of the coefficients of the  $g$ 's on the left-hand side and this yields  $g(p^i) = g(p)^i$  as desired.

Next, we show that  $g(p^i q^j) = g(p)^i g(q)^j$  for distinct primes  $p, q$  and  $i, j \in \mathbb{N}$ . Taking  $k = 2$ ,  $p_1 = p$ ,  $p_2 = q$ , and  $n_1 = n_2 = 1$  in (4.10), we get

$$\alpha(1, 1) g(p) g(q) = \frac{1}{1!} g(p g) + g(p) g(q). \quad (4.13)$$

Again since  $\alpha(1, 1)$  is the sum of the coefficients of the  $g$ 's on the right-hand side, we deduce that  $g(p) g(q) = g(pq)$ . Next, using  $g(p^i) = g(p)^i$  and induction on  $i + j$ , we deduce that  $g(p^i q^j) = g(p)^i g(q)^j$ . Finally,  $g(p_1^{i_1} \cdots p_k^{i_k}) = g(p_1)^{i_1} \cdots g(p_k)^{i_k}$  follows by induction on  $k$ .  $\square$

**5. Applications.** Our first application is related to one of our earlier results in [11] which states that for  $f \in \mathcal{M}$ ,  $\alpha \in \mathbb{R} \setminus \{0, 1\}$ , if  $f^\alpha = \mu_{-\alpha} f$  and  $f$  satisfies condition (NE), then  $f \in \mathcal{C}$ . Here condition (NE) reads: if  $\alpha$  is a negative even integer, then

$$f(p^{-\alpha-1}) = f(p)^{-\alpha-1} \quad \text{for each prime } p. \quad (5.1)$$

A similar result to the one above but without condition (NE) is the following proposition easily deduced from [11, Theorem 1.2].

**PROPOSITION 5.1.** *Let  $f \in \mathcal{M}$  and  $\alpha \in \mathbb{R} \setminus \{0\}$ . Then*

$$f \in \mathcal{C} \iff f^\alpha(p^a) = \mu_{-\alpha}(p^a) \left( \frac{1}{\alpha} f^\alpha(p) \right)^a \quad (5.2)$$

for all primes  $p$  and all  $a \in \mathbb{N}$ , where  $\mu_\alpha$  denotes the Hsu's generalized Möbius function.

Our second application is related to the notion of totients (see [7]).

Recall that  $f \in \mathcal{A}$  is a totient if it is of the form

$$f = f_t * f_v^{-1}, \quad (5.3)$$

where  $f_t$  (integral component) and  $f_v$  (inverse component) are both in  $\mathcal{C}$ . Those totients belonging to  $\mathcal{C}$  are characterized in the next proposition which is an easy consequence of Corollaries 2.2 and 2.3.

**PROPOSITION 5.2.** (i) *Let  $f = f_t * f_v^{-1}$  be a totient. Then*

$$f \in \mathcal{C} \iff f_v = I \text{ or } f_v^{-1}(p) + f_t(p) = 0 \quad (5.4)$$

for all primes  $p$ .

(ii) *Let  $f = f_t * f_v^{-1}$  and  $g = g_t * g_v^{-1}$  be two totients. Then*

$$f * g \text{ is a totient} \iff f_t(p)g_t(p) = 0 = f_v(p)g_v(p) \quad (5.5)$$

for all primes  $p$ .

Another related concept is the following:  $f \in \mathcal{A}$  is said to be a *rational arithmetical function of degree  $(r, s)$*  (see, e.g., [7]) if it is of the form

$$f = g_1 * g_2 * \dots * g_r * h_1^{-1} * h_2^{-1} * \dots * h_s^{-1}, \quad g_i, h_i \in \mathcal{C}. \quad (5.6)$$

Again by Corollaries 2.2 and 2.3, we have, for example, the following characterization which can be evidently generalized.

**PROPOSITION 5.3.** (i) *Let  $f = g_1 * g_2 * h_1^{-1}$  be a rational arithmetical function of degree  $(2, 1)$ . Then*

$$f \text{ is a totient} \iff g_1(p)g_2(p) = 0 \quad (5.7)$$

for each prime  $p$ .

(ii) *Let  $f = g_1 * h_1^{-1} * h_2^{-1}$  be a rational arithmetical function of degree  $(1, 2)$ . Then*

$$f \text{ is a totient} \iff h_1(p)h_2(p) = 0 \quad (5.8)$$

for each prime  $p$ .

Our last application is related to the concept of specially multiplicative functions. Recall that an arithmetic function  $F$  is said to be *specially multiplicative* (see [16]) if there is a completely multiplicative function  $f_A$  such that

$$F(m)F(n) = \sum_{d|(m,n)} F\left(\frac{mn}{d^2}\right) f_A(d). \quad (5.9)$$

It is also known that each specially multiplicative function is a product of two completely multiplicative functions.

A straightforward consequence of [Corollary 2.3](#) is the following result.

**PROPOSITION 5.4.** *Let  $F = f * g$  be a specially multiplicative function with  $f, g \in \mathcal{C}$ . Then*

$$F \in \mathcal{C} \iff f(p)g(p) = 0 \quad (5.10)$$

for all primes  $p$ .

**ACKNOWLEDGMENT.** This work was partially supported by The University of the Thai Chamber of Commerce.

## REFERENCES

- [1] T. M. Apostol, *Some properties of completely multiplicative arithmetical functions*, Amer. Math. Monthly **78** (1971), 266–271.
- [2] ———, *Introduction to Analytic Number Theory*, Undergraduate Texts in Mathematics, Springer-Verlag, New York, 1976.
- [3] T. C. Brown, L. C. Hsu, J. Wang, and P. J.-S. Shiue, *On a certain kind of generalized number-theoretical Möbius function*, Math. Sci. **25** (2000), no. 2, 72–77.
- [4] L. Carlitz and M. V. Subbarao, *Transformation of arithmetic functions*, Duke Math. J. **40** (1973), 949–958.
- [5] T. B. Carroll, *A characterization of completely multiplicative arithmetic functions*, Amer. Math. Monthly **81** (1974), 993–995.
- [6] E. D. Cashwell and C. J. Everett, *The ring of number-theoretic functions*, Pacific J. Math. **9** (1959), 975–985.
- [7] P. Haukkanen, *Some characterizations of totients*, Int. J. Math. Math. Sci. **19** (1996), no. 2, 209–217.
- [8] ———, *On the real powers of completely multiplicative arithmetical functions*, Nieuw Arch. Wisk. (4) **15** (1997), no. 1–2, 73–77.
- [9] J. Knopfmacher, *Abstract Analytic Number Theory*, Holland Mathematical Library, vol. 12, North-Holland Publishing, Amsterdam, 1975.
- [10] V. Laohakosol, N. Pabhapote, and N. Wechwiriyakul, *Logarithmic operators and characterizations of completely multiplicative functions*, Southeast Asian Bull. Math. **25** (2001), no. 2, 273–281.
- [11] ———, *Characterizing completely multiplicative functions by generalized Möbius functions*, Int. J. Math. Math. Sci. **29** (2002), no. 11, 633–639.
- [12] D. Rearick, *Operators on algebras of arithmetic functions*, Duke Math. J. **35** (1968), 761–766.
- [13] ———, *The trigonometry of numbers*, Duke Math. J. **35** (1968), 767–776.
- [14] J. Riordan, *Combinatorial Identities*, John Wiley & Sons, New York, 1968.
- [15] H. N. Shapiro, *Introduction to the Theory of Numbers*, Pure and Applied Mathematics, John Wiley & Sons, New York, 1983.

[16] R. Sivaramakrishnan, *Classical Theory of Arithmetic Functions*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 126, Marcel Dekker, New York, 1989.

Vichian Laohakosol: Department of Mathematics, Faculty of Science, Kasetsart University, Bangkok 10900, Thailand

*E-mail address:* [fscivil@nontri.ku.ac.th](mailto:fscivil@nontri.ku.ac.th)

Nittiya Pabhapote: Department of Mathematics, Faculty of Science, The University of the Thai Chamber of Commerce, Bangkok 10400, Thailand

*E-mail address:* [nittiya\\_pab@utcc.ac.th](mailto:nittiya_pab@utcc.ac.th)

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)