

## EXTENSION OF $\alpha$ -LABELINGS OF QUADRATIC GRAPHS

KOUSHROSH ESHGHI

Received 1 June 2003

First, a new proof for the existence of an  $\alpha$ -labeling of the quadratic graph  $Q(3,4k)$  is presented. Then the existence of  $\alpha$ -labelings of special classes of quadratic graphs with some isomorphic components is shown.

2000 Mathematics Subject Classification: 05C78.

**1. Introduction.** In this paper, all graphs are finite without loops or multiple edges, and all parameters are positive integers. The symbols  $|A|$ ,  $P_n$ , and  $C_n$  denote the cardinality of set  $A$ , a snake, and a cycle with  $n$  edges, respectively. A sequence of numbers in parentheses or square brackets indicates the labels of vertices of a graph or subgraph under consideration according to whether it is a snake or cycle, respectively.

**DEFINITION 1.1.** A *graceful labeling* (or  $\beta$ -labeling) of a graph  $G = (V, E)$ , with  $m$  vertices and  $n$  edges, is a one-to-one mapping  $\Psi$  of the vertex set  $V(G)$  into the set  $\{0, 1, 2, \dots, n\}$  with this property: if we define, for any edge  $e = \{u, v\} \in E(G)$ , the value  $\Phi(e) = |\Psi(u) - \Psi(v)|$ , then  $\Phi$  is a one-to-one mapping of the set  $E(G)$  onto the set  $\{1, 2, \dots, n\}$ .

A graph is called *graceful* if it has a graceful labeling. Not all graphs are graceful. For example,  $C_5$  and  $K_5$  are not graceful.

**DEFINITION 1.2.** An  $\alpha$ -labeling of a graph  $G = (V, E)$  is a graceful labeling of  $G$  which satisfies the following additional condition: there exists a number  $\gamma$  ( $0 \leq \gamma \leq |E(G)|$ ) such that, for any edge  $e \in E(G)$  with end vertices  $u, v \in V(G)$ ,  $\min[\Psi(u), \Psi(v)] \leq \gamma < \max[\Psi(u), \Psi(v)]$ . The values of an  $\alpha$ -labeling  $\Psi$  which are less than or equal to  $\gamma$  are referred to as “*small values*” and the remaining values of  $\Psi$  as the “*large values*” of the given  $\alpha$ -labeling.

The concepts of a graceful labeling and of an  $\alpha$ -labeling were introduced by Rosa [8]. Rosa proved that any graceful Eulerian graph  $G$  satisfies the condition  $|E(G)| \equiv 0$  or  $3 \pmod{4}$ . This implies that any Eulerian graph  $G$  with an  $\alpha$ -labeling satisfies the condition  $|E(G)| \equiv 0 \pmod{4}$  ( $G$  is bipartite). It is also known that these conditions are also sufficient if  $G$  is a cycle [8]. Abrham and Kotzig [3] proved that Rosa’s condition is also sufficient for 2-regular graphs with two components. The author [4] proved the similar result for 2-regular graphs with three components with the exception of one special case.

A detailed history of the graph labeling problems and related results appears in Gallian [6]. One of the results of Abrham and Kotzig should be mentioned here.

TABLE 1.1. Some of the results about  $\alpha$ -labelings of quadratic graphs.

Graph	Graceful labeling	$\alpha$ -labeling
$Q(1, s)$	It is graceful if and only if $s \equiv 0$ or $3 \pmod{4}$ [8]	It has an $\alpha$ -labeling if and only if $s \equiv 0 \pmod{4}$ [8]
$Q(2, s)$	It is graceful if and only if $s$ is even and $s > 2$ [7]	It has an $\alpha$ -labeling if and only if $s$ is even and $s > 2$ [7]
$Q(3, 4k)$	It is graceful for each $k \geq 1$ [7]	It has an $\alpha$ -labeling for each $k > 1$ [7]
$Q(r, 3)$	It is graceful if and only if $r = 1$ [7]	It has no $\alpha$ -labeling [7]
$Q(r, 4)$	It is graceful for each $r \geq 1$ [2]	It has an $\alpha$ -labeling for each $r \geq 1, r \neq 3$ [2]
$Q(r, 5)$	It is not graceful for any $r \geq 1$ [7]	It has no $\alpha$ -labeling [7]
$Q(5, 4k)$	It is graceful for all $k \geq 1$ [5]	It has an $\alpha$ -labeling for all $k \geq 1$ [5]

**DEFINITION 1.3.** If  $G$  is a 2-regular graph on  $n$  vertices and  $n$  edges, which has a graceful labeling  $\Psi$ , then there exists exactly one number  $x$  ( $0 < x < n$ ) such that  $\Psi(v) \neq x$  for all  $v \in V(G)$ ; this number  $x$  is referred to as the missing value of the graceful labeling [1].

In the work done here, the problem of existence of an  $\alpha$ -labeling of a special class of 2-regular graphs, called quadratic graph, is investigated.

**DEFINITION 1.4.** A quadratic graph  $Q(r, s)$  is a 2-regular graph with  $r$  components, each of which is a cycle of length  $s$ .

Some of the results about  $\alpha$ -labelings of quadratic graphs published in the literature are summarized in Table 1.1.

## 2. The existence of an $\alpha$ -labeling of $Q(3, 4k)$

**THEOREM 2.1.** A  $Q(3, 4k)$ -graph has an  $\alpha$ -labeling for each  $k > 1$ .

**PROOF.** In this case we have a graph consisting of three cycles of length  $4k$ . We know that an  $\alpha$ -labeling for this graph was constructed by Kotzig in [7]. We now present a different construction of an  $\alpha$ -labeling of  $3C_{4k}$  for  $k > 1$ ; its advantage is that it makes it possible to obtain certain results in Section 3.

The vertices of the first  $C_{4k}$  are successively labeled as follows:  $[0, 12k, 1, 12k - 1, 2, 12k - 2, \dots, k - 1, 11k + 1, k + 1, 11k, \dots, 2k - 1, 10k + 2, 2k, 10k + 1]$ . The resulting edge values of the first  $C_{4k}$  are then  $12k, 12k - 1, 12k - 2, \dots, 10k + 2, 10k, \dots, 8k + 2, 8k + 1$ , and  $10k + 1$ .

The vertices of the second  $C_{4k}$  are consecutively labeled by the numbers  $[4k, 8k, 4k + 1, 8k - 1, 4k + 2, 8k - 2, \dots, 5k - 1, 7k + 1, 5k + 1, 7k, \dots, 6k - 1, 6k + 2, 6k, 6k + 1]$ . The

TABLE 2.1. An  $\alpha$ -labeling of  $Q(3,4k)$  for  $2 \leq k \leq 5$ .




$k$	$Q(3,4k)$	An $\alpha$ -labeling of $Q(3,4k)$
5	$Q(3,20)$	$[0, 51, 10, 52, 9, 53, 8, 54, 7, 55, 6, 56, 4, 57, 3, 58, 2, 59, 1, 60], [20, 31, 30, 32, 29, 33, 28, 34, 27, 35, 26, 36, 24, 37, 23, 38, 22, 39, 21, 40, 20], [5, 44, 17, 43, 14, 42, 18, 41, 16, 50, 19, 49, 11, 48, 12, 47, 25, 46, 13, 45]$
4	$Q(3,16)$	$[0, 41, 8, 42, 7, 43, 6, 44, 5, 45, 3, 46, 2, 47, 1, 48], [16, 25, 24, 26, 23, 27, 22, 28, 21, 29, 19, 30, 18, 31, 17, 32], [4, 35, 15, 40, 11, 38, 20, 37, 9, 39, 13, 34, 10, 33, 14, 36]$
3	$Q(3,12)$	$[0, 31, 6, 32, 5, 33, 4, 34, 2, 35, 1, 36], [12, 19, 18, 20, 17, 21, 16, 22, 14, 23, 13, 24], [3, 26, 10, 25, 8, 30, 11, 29, 15, 28, 7, 27]$
2	$Q(3,8)$	$[0, 21, 4, 22, 3, 23, 1, 24], [8, 13, 12, 14, 11, 15, 9, 16], [2, 17, 5, 19, 10, 20, 7, 18]$

resulting edge values of the second  $C_{4k}$  are then  $4k, 4k - 1, 4k - 2, \dots, 2k + 2, 2k, 2k - 1, \dots, 3, 2, 1$ , and  $2k + 1$ . The missing value of the first  $C_{4k}$  is equal to  $k$  and the missing value of the second  $C_{4k}$  is equal to  $5k$ . The missing value of the main graph is equal to  $3k$  and  $\gamma = 6k$ .

Now we construct the third cycle  $C_{4k}$ . First suppose that  $k \geq 6$ . Next we construct three snakes. The vertices of the first snake are successively labeled as  $10k - 1, 2k + 1, 10k - 2, 2k + 2, \dots, 3k - 4, 9k + 3, 3k - 3$ , and  $9k + 2$ . The resulting edge values of this snake are then  $8k - 2, 8k - 3, 8k - 4, \dots, 6k + 7, 6k + 6$ , and  $6k + 5$ . The vertices of the second snake are consecutively labeled by the numbers  $9k - 5, 3k - 1, 9k - 2, 3k + 1$ , and  $9k - 1$ ; this yields the edge values  $6k - 4, 6k - 1, 6k - 3$ , and  $6k - 2$ . Finally the vertices of the third snake are labeled as  $9k - 1, k, 9k, 3k - 2, 9k + 1, 5k$ , and  $9k + 2$ ; this yields the edge values  $8k - 1, 8k, 6k + 2, 6k + 3, 4k + 1$ , and  $4k + 2$ . Now we generate the edge labels  $6k + 4, 6k + 1$ , and  $6k$  by connecting the following pairs of vertices to each other respectively:  $4k - 4$  and  $10k$ ;  $4k - 1$  and  $10k$ ;  $4k - 1$  and  $10k - 1$ . In order to generate the rest of the edge labels, we need to use a special type of transforming of vertex labels, described in the appendix as “transformation of labels procedure.” Therefore, in the next step, we apply the transformation of labels procedure to the remaining vertex labels, that is,  $(3k + 2, 3k + 3, \dots, 4k - 4, 4k - 3, 4k - 2)$  and  $(8k + 1, 8k + 2, \dots, 9k - 5, 9k - 4, 9k - 3)$  by considering the two vertices  $4k - 4$  and  $9k - 5$  as end vertices. This transformation generates the rest of the edge labels and the construction of the last  $C_{4k}$  is completed. The construction of an  $\alpha$ -labeling of  $Q(3,4k)$  with  $x = 3k$  and  $\gamma = 6k$  for  $2 \leq k \leq 5$  is illustrated in Table 2.1.  $\square$

**3. Existence of  $\alpha$ -labelings of general classes of quadratic graphs.** The following concept presented in [5] is very useful for further considerations in this section.

**DEFINITION 3.1.** The graph  $C_{4k}$  has a *standard labeling* if the values of the vertices of  $C_{4k}$  can be generated from an  $\alpha$ -labeling of  $C_{4k}$  by adding constant factor(s) to the small or large values (or both) of an  $\alpha$ -labeling of  $C_{4k}$ .

FIGURE 3.1. An  $\alpha$ -labeling of the graph  $C_8 \cup C_{12}$ .FIGURE 3.2. Transformations of an  $\alpha$ -labeling of the graph  $C_8$  to a standard labeling.FIGURE 3.3. An  $\alpha$ -labeling of  $C_{12} \cup Q(2,4)$ .

**EXAMPLE 3.2.** In Figure 3.1, an  $\alpha$ -labeling of the graph  $C_8 \cup C_{12}$  is presented. This graph consists of the disjoint union of two cycles and has 20 vertices. According to the results presented in [1], we know that in this graph the missing value is 5 and  $\gamma = 10$ .

In the above  $\alpha$ -labeling,  $C_8$  has a standard labeling because it can be generated from an  $\alpha$ -labeling of  $C_8$  only by increasing the large values of this construction by 12, see Figure 3.2.

If a graph  $C_{4k}$  has a standard labeling, it can be replaced by any  $\alpha$ -labeling of the disjoint union of cycles in the form of  $\bigcup_{i=1}^n C_{4k_i}$  by considering the constant factor(s) if there is an  $\alpha$ -labeling for  $\bigcup_{i=1}^n C_{4k_i}$  and  $k = k_1 + k_2 + \dots + k_n$ .

**EXAMPLE 3.3.** Since we know that  $Q(2,4)$  has an  $\alpha$ -labeling, the standard labeling of  $C_8$  in Figure 3.1 can be replaced by an  $\alpha$ -labeling of  $Q(2,4)$  to form an  $\alpha$ -labeling of  $C_{12} \cup Q(2,4)$  if we increase the large values of an  $\alpha$ -labeling  $Q(2,4)$  by 12, see Figure 3.3.

In the construction of an  $\alpha$ -labeling of  $Q(3,4k)$ , the first and second  $C_{4k}$  have standard labelings because the first cycle can be generated by adding  $8k$  to the large values of an  $\alpha$ -labeling of  $C_{4k}$  with  $x = k$ ,  $\gamma = 2k$ , and the second cycle can be generated by adding  $4k$  to the small and large values of an  $\alpha$ -labeling of  $C_{4k}$  with  $x = k$ ,  $\gamma = 2k$ .

In the following theorems, we use this property to extend the class of quadratic graphs with isomorphic components that have  $\alpha$ -labelings.

**THEOREM 3.4.** *The following graphs have  $\alpha$ -labelings if  $k = 3k_1$ ,  $k_i = 3k_{i+1}$ ,  $k_i > 1$ ,  $i = 1, 2, 3, \dots, n-1$ :*

- (i)  $\bigcup_{i=1}^n Q(2, 4k_i) \cup Q(2, 4k) \cup C_{4k_n}$ ,
- (ii)  $\bigcup_{i=1}^n Q(4, 4k_i) \cup Q(2, 4k_n) \cup C_{4k}$ .

**PROOF.** It is shown that in the construction of an  $\alpha$ -labeling of  $Q(3, 4k)$ , two isomorphic components  $C_{4k}$  have standard labelings. Now we apply the following transformations in order to obtain the proof of each part of the theorem.

In the construction of an  $\alpha$ -labeling of  $Q(3, 4k)$ , substitute one of the components of  $C_{4k}$  with standard labeling by  $Q(3, 4k_1)$ ,  $k = 3k_1$ . Then, since at least one component of  $Q(3, 4k_1)$  still has a standard labeling, we are able to replace it again by  $Q(3, 4k_2)$ ,  $k_1 = 3k_2$ . In the next stages, we continue to replace one component of each  $Q(3, 4k_i)$  by  $Q(3, 4k_{i+1})$ ,  $k_i = 3k_{i+1}$ , for  $i = 2, 3, \dots, n-1$ , to obtain an  $\alpha$ -labeling of the first graph of the theorem.

The proof of the second part is similar to the proof of the first part. This time we use the replacements for both components with standard labelings in an  $\alpha$ -labeling of  $Q(3, 4k_i)$ .  $\square$

**EXAMPLE 3.5.** The following classes of graphs have  $\alpha$ -labelings according to [Theorem 2.1](#), for  $k = 6$  and  $k_1 = 2$ :

$$Q(3, 8) \cup Q(2, 24), \quad Q(6, 8) \cup C_{24}. \quad (3.1)$$

**THEOREM 3.6.** *The following graphs have  $\alpha$ -labelings if  $k = \sum_{i=1}^n k_i$  and  $k_i \geq \sum_{t=i+1}^n k_t$  for  $i = 1, 2, 3, \dots, n-1$ :*

- (i)  $\bigcup_{i=1}^n C_{4k_i} \cup Q(2, 4k)$ ,
- (ii)  $\bigcup_{i=1}^n Q(2, 4k_i) \cup C_{4k}$ .

**PROOF.** In the construction of an  $\alpha$ -labeling of  $Q(3, 4k)$ , at least two cycles  $C_{4k}$  have standard  $\alpha$ -labelings. In order to obtain the different parts of [Theorem 3.4](#), apply the following replacements.

(i) Consider one of the standard labelings of  $C_{4k}$ . First we replace it by  $C_{4k_1} \cup C_{4q_1}$ , where  $q_1 \leq k_1$  and  $k = k_1 + q_1$ . Then, since  $C_{4q_1}$  still has a standard labeling [3], it can be replaced again by  $C_{4k_2} \cup C_{4q_2}$ , where  $q_2 \leq k_2$  and  $q_1 = k_2 + q_2$ . In the next stages, we continue to replace each  $C_{4q_i}$  by  $C_{4k_{i+1}} \cup C_{4q_{i+1}}$ ,  $q_{i+1} \leq k_{i+1}$ , where  $q_i = k_{i+1} + q_{i+1}$  for  $i = 2, 3, \dots, n-2$ , and  $k_n = q_{n-1}$ .

(ii) We apply the replacement procedure of the first part for both  $C_{4k}$  which have standard labelings in an  $\alpha$ -labeling of  $Q(3, 4k)$ .  $\square$

**EXAMPLE 3.7.** The following classes of graphs have  $\alpha$ -labelings, for  $r, t \geq 1$ :

$$C_{4r} \cup C_{4t} \cup Q(2, 4(r+t)), \quad C_{4(r+t)} \cup Q(2, 4r) \cup Q(2, 4t). \quad (3.2)$$

**THEOREM 3.8.** *The following graphs have  $\alpha$ -labelings if  $k = 3k_1$ ,  $k_i = 3k_{i+1}$ ,  $k_1 > 1$ ,  $i = 1, 2, 3, \dots, n-1$ :*

- (i)  $\bigcup_{i=1}^n Q(2, 4k_i) \cup Q(4, 4k) \cup C_{4k_n}$ ,
- (ii)  $\bigcup_{i=1}^n Q(4, 4k_i) \cup Q(2, 4k_n) \cup Q(3, 4k)$ ,
- (iii)  $\bigcup_{i=1}^n Q(6, 4k_i) \cup Q(3, 4k_n) \cup Q(2, 2k)$ .

**PROOF.** It has been shown that in the construction of an  $\alpha$ -labeling of  $Q(5, 4k)$ , at least three isomorphic components  $C_{4k}$  have standard labelings [5].

First consider one of the components of  $C_{4k}$  with standard labeling in the construction of an  $\alpha$ -labeling of  $Q(5, 4k)$ . Then substitute it by  $Q(3, 4k_1)$ , where  $k = 3k_1$ ,  $k_1 > 1$ . Since at least one component of  $Q(3, 4k_1)$  still has a standard labeling, it can be replaced again by  $Q(3, 4k_2)$ ,  $k_1 = 3k_2$ . In the next stages, we continue to replace one component of each  $Q(3, 4k_i)$  by  $Q(3, 4k_{i+1})$ , where  $k_i = 3k_{i+1}$ , for  $i = 2, 3, \dots, n-1$ . Finally we obtain an  $\alpha$ -labeling of the graph in the first part of the theorem.

The proof of the second (and third) part of the theorem can be easily obtained by applying the above replacements to the second (and third) isomorphic component of  $C_{4k}$  with standard labelings in an  $\alpha$ -labeling of  $Q(5, 4k)$ .  $\square$

**EXAMPLE 3.9.** The following classes of graphs have  $\alpha$ -labelings according to Theorem 3.8, for  $k = 18$ ,  $k_1 = 6$ , and  $k_2 = 2$ :

$$\begin{aligned} & Q(3, 8) \cup Q(2, 24) \cup Q(4, 72), \\ & Q(6, 8) \cup Q(4, 24) \cup Q(3, 72), \\ & Q(9, 8) \cup Q(6, 24) \cup Q(2, 36). \end{aligned} \tag{3.3}$$

**THEOREM 3.10.** *The following graphs have  $\alpha$ -labelings if  $k = 5k_1$ ,  $k_i = 5k_{i+1}$ ,  $i = 1, 2, 3, \dots, n-1$ :*

- (i)  $\bigcup_{i=1}^n Q(4, 4k_i) \cup Q(2, 4k) \cup C_{4k_n}$ ,
- (ii)  $\bigcup_{i=1}^n Q(8, 4k_i) \cup Q(2, 4k_n) \cup C_{4k}$ .

**PROOF.** In the first part of the theorem, consider one of the components of  $C_{4k}$  with standard labeling in an  $\alpha$ -labeling of  $Q(5, 4k)$ . Then substitute it by  $Q(5, 4k_1)$ ,  $k = 5k_1$ . We know that in the construction of an  $\alpha$ -labeling of  $Q(5, 4k)$ , at least three isomorphic components  $C_{4k}$  have standard labelings [5]. Then, since at least one component of  $Q(5, 4k_1)$  still has a standard labeling, we are able to replace it again by  $Q(5, 4k_2)$ ,  $k_1 = 5k_2$ . In the next stages, we continue to replace one component of each  $Q(5, 4k_i)$  by  $Q(5, 4k_{i+1})$ , where  $k_i = 5k_{i+1}$ , for  $i = 2, 3, \dots, n-1$ . Finally we obtain an  $\alpha$ -labeling of the graph in the first part of the theorem.

The proof of the second part is similar to the proof of the first part. This time we use the replacements for two isomorphic components of  $C_{4k}$  with standard labelings in an  $\alpha$ -labeling of  $Q(5, 4k)$ .  $\square$

**EXAMPLE 3.11.** The following classes of graphs have  $\alpha$ -labelings for  $r \geq 1$ :

$$Q(5, 4r) \cup Q(2, 20r), \quad Q(10, 4r) \cup C_{20r}. \tag{3.4}$$

$$0 \quad 1 \quad 2 \quad \dots \quad w = r \quad \dots \quad k-1 \quad k$$

$$2k+1 \quad 2k \quad \dots \quad z = k+1+r \quad \dots \quad k+2 \quad k+1$$

FIGURE A.1. Arrangement of vertex labels of snake  $P_{2k+1}$  according to [Lemma A.1](#).

$$n \quad n+1 \quad n+2 \quad \dots \quad n+w \quad \dots \quad n+k-1 \quad n+k$$

$$m+k \quad m+k-1 \quad \dots \quad m-(k+1)+z \quad \dots \quad m+1 \quad m$$

FIGURE A.2. Arrangement of vertex labels in the transformation.

**THEOREM 3.12.** *The following classes of graphs have  $\alpha$ -labelings for  $r, t \geq 1$ :*

- (i)  $C_{4r+2} \cup C_{4t+2} \cup Q(2, 4(r+t+1))$ ,
- (ii)  $C_{4(r+t)} \cup Q(2, 4r+2) \cup Q(2, 4t+2)$ .

**PROOF.** In the first part, we need to replace one of the standard labelings of  $C_{4k}$  in the construction of  $Q(3, 4k)$  by  $C_{4r+2} \cup C_{4t+2}$ ,  $r, t \geq 1$  and  $r+t+1 = k$ , because we know that the graph  $C_{4r+2} \cup C_{4t+2}$  has an  $\alpha$ -labeling for  $r, t \geq 1$  [\[3\]](#). In the second part of the theorem, we replace both the standard  $\alpha$ -labelings of  $C_{4k}$  and the construction of  $Q(3, 4k)$  by  $C_{4r+2} \cup C_{4t+2}$ ,  $r, t \geq 1$ , and  $r+t+1 = k$ , respectively.  $\square$

## Appendix

**Transformation of labels procedure.** The transformation presented below is used in [Theorem 2.1](#).

**LEMMA A.1** (Abrham and Kotzig [\[3\]](#)). *Let  $r$  be a nonnegative integer and let  $s$  be an odd integer,  $s = 2k+1 \geq 2r+1$ . Then  $P_s$  has an  $\alpha$ -labeling  $\Psi$  with endpoints labelled  $w$  and  $z$  that satisfy the conditions  $z-w = k+1$  and  $w=r$ . (Without loss of generality, assume that  $w < z$ .)*

Given any  $0 \leq w \leq k$ ,  $k+1 \leq z \leq 2k+1$ , and  $z-w = k+1$ , we can always construct an  $\alpha$ -labeling for a bipartite snake  $P_{2k+1}$  with edge labels 1 through  $2k+1$  and endpoints  $w$  and  $z$ , with  $y = k$ ,  $w = r$ , and  $z = k+r+1$ , see [Figure A.1](#).

Now suppose we add  $n$  to the upper half and add  $m-(k+1)$  to the lower half for any positive integers  $m$  and  $n$ , where  $m-1 > n+k$ , see [Figure A.2](#).

Then the edge labels increase by precisely  $m-(k+1)-n$ . The transformation produces the edge labels from  $[m-k-n]$  through  $[m+k-n]$  according to [Lemma A.1](#).

## REFERENCES

- [1] J. Abrham and A. Kotzig, *On the missing value in a graceful numbering of a 2-regular graph*, Congr. Numer. **65** (1988), 261–266.
- [2] ———, *All 2-regular graphs consisting of 4-cycles are graceful*, Discrete Math. **135** (1994), no. 1-3, 1–14.
- [3] ———, *Graceful valuations of 2-regular graphs with two components*, Discrete Math. **150** (1996), no. 1-3, 3–15.
- [4] K. Eshghi, *Construction of  $\alpha$ -labeling of 2-regular graphs with three components*, Ph.D. thesis, University of Toronto, Toronto, 1997.
- [5] ———,  *$\alpha$ -valuations of special classes of quadratic graphs*, Bull. Iranian Math. Soc. **28** (2002), no. 1, 29–42.
- [6] J. A. Gallian, *A dynamic survey of graph labeling*, Electron. J. Combin. **5** (1998), no. 1, 1–43.
- [7] A. Kotzig,  *$\beta$ -valuations of quadratic graphs with isomorphic components*, Utilitas Math. **7** (1975), 263–279.
- [8] A. Rosa, *On certain valuations of the vertices of a graph*, Theory of Graphs (Internat. Sympos., Rome, 1966), Gordon and Breach, New York, 1967, pp. 349–355.

Kourosh Eshghi: Industrial Engineering Department, Sharif University of Technology, Tehran 14584, Iran

E-mail address: [eshghi@sharif.edu](mailto:eshghi@sharif.edu)

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)