

GENERALIZATIONS OF PRINCIPALLY QUASI-INJECTIVE MODULES AND QUASIPRINCIPALLY INJECTIVE MODULES

ZHU ZHANMIN, XIA ZHANGSHENG, AND TAN ZHISONG

Received 22 November 2004 and in revised form 9 June 2005

Let R be a ring and M a right R -module with $S = \text{End}(M_R)$. The module M is called almost principally quasi-injective (or APQ-injective for short) if, for any $m \in M$, there exists an S -submodule X_m of M such that $l_{Mrr}(m) = Sm \oplus X_m$. The module M is called almost quasiprincipally injective (or AQP-injective for short) if, for any $s \in S$, there exists a left ideal X_s of S such that $l_S(\ker(s)) = Ss \oplus X_s$. In this paper, we give some characterizations and properties of the two classes of modules. Some results on principally quasi-injective modules and quasiprincipally injective modules are extended to these modules, respectively. Specially in the case R_R , we obtain some results on AP-injective rings as corollaries.

1. Introduction

Throughout R is a ring with identity and M is a right R -module with $S = \text{End}(M_R)$. Recall a ring R is called right principally injective [5] (or right P-injective for short) if, every homomorphism from a principally right ideal of R to R can be extended to an endomorphism of R , or equivalently, $l_r(a) = Ra$ for all $a \in R$. The notion of right P-injective rings has been generalized by many authors. For example, in [4, 8], right P-injective rings are generalized to modules in two ways, respectively. Following [4], the module M is called principally quasi-injective (or PQ-injective for short) if, each R -homomorphism from a principal submodule of M to M can be extended to an endomorphism of M . This is equivalent to saying that $l_{Mrr}(m) = Sm$ for all $m \in M$, where $l_{Mrr}(m)$ consists of all elements $z \in M$ such that $mx = 0$ implies $zx = 0$ for any $x \in R$. In [8], the module M is called quasiprincipally injective (or QP-injective for short) if, every homomorphism from an M -cyclic submodule of M to M can be extended to an endomorphism of M , or equivalently, $l_s(\ker(s)) = Ss$ for all $s \in S$. In [6], right P-injective rings are generalized to almost principally injective rings, that is, a ring R is said to be almost principally injective (or AP-injective for short) if, for any $a \in R$, there exists a left ideal X_a such that $l_r(a) = Ra \oplus X_a$. The nice structure of PQ-injective modules, QP-injective modules, and AP-injective rings draws our attention to define almost PQ-injective modules and almost QP-injective modules in similar ways to AP-injective rings, and to investigate their characterizations and properties.

2. APQ-injective modules

Definition 2.1. Let M be a right R -module and let $S = \text{End}(M_R)$. The module M is called almost principally quasi-injective (briefly, APQ-injective) if, for any $m \in M$, there exists an S -submodule X_m of M such that $l_M r_R(m) = Sm \oplus X_m$.

The concept of APQ-injective modules is explained by the following lemma.

LEMMA 2.2. Let M_R be a module and let $S = \text{End}(M_R)$, and $m \in M$.

- (1) If $l_M r_R(m) = Sm \oplus X$ for some $X \subseteq M$ as left S -modules, then $\text{Hom}_R(mR, M) = S \oplus \Gamma$ as left S -modules, where $\Gamma = \{f \in \text{Hom}_R(mR, M) \mid f(m) \in X\}$.
- (2) If $\text{Hom}_R(mR, M) = S \oplus \Gamma$ as left S -modules, then $l_M(r_R(m)) = Sm \oplus X$ as left S -modules, where $X = \{f(m) \mid f \in \Gamma\}$.
- (3) Sm is a summand of $l_M(r_R(m))$ as left S -modules if and only if S is a summand of $\text{Hom}_R(mR, M)$ as left S -modules.

Proof. The map $\theta : l_M(r_R(m)) \rightarrow \text{Hom}_R(mR, M)$ with $\theta(a) = \lambda_a$ is a left S -isomorphism, where $\lambda_a : mR \rightarrow M$ is defined by $\lambda_a(mr) = ar$, so the lemma follows. Moreover, $_S(Sm)$ is nonsmall in $l_M(r_R(m))$ if and only if S is nonsmall in $\text{Hom}_R(mR, M)$. \square

From Lemma 2.2, the following corollary follows.

COROLLARY 2.3 [4, Lemma 1.1]. Let M_R be a right R -module with $S = \text{End}(M_R)$ and $m \in M$. Then $l_M(r_R(m)) = Sm$ if and only if every R -homomorphism of mR into M extends to M .

From Corollary 2.3, we see that all PQ-injective modules are APQ-injective. Since a ring R is right P-injective (resp., AP-injective) if and only if the right R -module R_R is PQ-injective (resp., APQ-injective), and Page and Zhou [6] have given three examples of rings which are right AP-injective but not right P-injective, so in general, APQ-injective modules need not be PQ-injective.

Recall that a ring R is called right QP-injective [6, Definition 2.1], if for any $0 \neq a \in R$, there exists a left ideal X_a such that $l_R(a) = Ra + X_a$ with $a \notin X_a$. Now we extend this concept to modules.

Definition 2.4. Let M be a right R -module with $S = \text{End}(M_R)$, the module M is said to be QPQ-injective (i.e., quasiprincipally quasi-injective) if, for any nonzero element m of M , there exists an S -submodule X_m of M such that $l_M r_R(m) = Sm + X_m$ with $m \notin X_m$.

Clearly, right APQ-injective modules are QPQ-injective, but the reverse implication is not true. For example, Z -module Z_Z is QPQ-injective, but not APQ-injective.

Let M be a right R -module with $S = \text{End}(M_R)$, and $J(S)$ the Jacobson radical of S . Following [4], write $W(S) = \{w \in S \mid \ker(w) \subseteq^{\text{ess}} M\}$.

THEOREM 2.5. Let M_R be QPQ-injective with $S = \text{End}(M_R)$. Then

- (1) $J(S) \subseteq W(S)$,
- (2) $\text{Soc}(M_R) \subseteq r_M(J(S))$.

Proof. (1) Let $a \in J(S)$. If $a \notin W(S)$, then $\ker(a) \cap K = 0$ for some $0 \neq K \leq M_R$. Take $k \in K$ such that $ak \neq 0$, then $l_M(r_R(ak)) = S(ak) + X_{ak}$ with $ak \notin X_{ak}$. If $r \in r_R(ak)$, then $kr \in \ker(a) \cap K$, so $kr = 0$, and hence $r \in r_R(k)$. This shows that $r_R(ak) = r_R(k)$. Note that

$k \in l_M(r_R(k)) = l_M(r_R(ak)) = S(ak) + X_{ak}$, so we may write $k = b(ak) + x$, where $b \in S$ and $x \in X$. Then $(1 - ba)k = x$, and so $k = (1 - ba)^{-1}x$. Thus $ak = a(1 - ba)^{-1}x \in X_{ak}$, a contradiction.

(2) Let $mR \subseteq M$ be simple. Suppose $am \neq 0$ for some $a \in J(S)$. Then, since mR is simple, $r_R(am) = r_R(m)$. Since M_R is QPQ-injective, there is a left S -module X such that $am \notin X$ and $l_M r_R(am) = S(am) + X$. Note that $m \in l_M r_R(am)$, and so we may write $m = b(am) + x$, where $b \in S$ and $x \in X$. Then $(1 - ba)m = x$, so $m = (1 - ba)^{-1}x \in X$. This means that $am \in X$, a contradiction. \square

COROLLARY 2.6. *Let M_R be QPQ-injective with $S = \text{End}(M_R)$. If S is semilocal, then $\text{Soc}(M_R) \subseteq \text{Soc}({}_S M)$.*

Proof. This follows from Theorem 2.5(2) and [1, Proposition 15.17]. \square

LEMMA 2.7. *Let M_R be APQ-injective with $S = \text{End}(M_R)$. If $s \notin W(S)$, then the inclusion $\ker(s) \subset \ker(s - sts)$ is strict for some $t \in S$.*

Proof. If $s \notin W(S)$, then $\ker(s) \cap mR = 0$ for some $0 \neq m \in M$. Thus $r_R(m) = r_R(sm)$, and so $l_M r_R(m) = l_M r_R(sm) = S(sm) \oplus X_{sm}$ as left S -modules because M_R is APQ-injective. Write $m = t(sm) + x$, where $x \in X_{sm}$. Then $(s - sts)m = sx \in S(sm) \cap X_{sm}$, and hence $(s - sts)m = 0$. Therefore, the inclusion $\ker(s) \subset \ker(s - sts)$ is strict. \square

LEMMA 2.8. *Let M be a right R -module with $S = \text{End}(M_R)$. Suppose that for any sequence $\{s_1, s_2, \dots\} \subseteq S$, the chain $\ker(s_1) \subseteq \ker(s_2 s_1) \subseteq \dots$ terminates. Then*

- (1) $W(S)$ is right T -nilpotent,
- (2) $S/W(S)$ contains no infinite set of nonzero pairwise orthogonal idempotents.

Proof. This is a corollary of [2, Lemma 1.9]. \square

THEOREM 2.9. *Let M_R be APQ-injective with $S = \text{End}(M_R)$, then the following conditions are equivalent.*

- (1) S is right perfect.
- (2) For any sequence $\{s_1, s_2, \dots\} \subseteq S$, the chain $\ker(s_1) \subseteq \ker(s_2 s_1) \subseteq \dots$ terminates.

Proof. (1) \Rightarrow (2). Let $s_i \in S$, $i = 1, 2, \dots$. Since S is right perfect, S satisfies DCC on principal left ideals. So the chain $Ss_1 \supseteq Ss_2 s_1 \supseteq \dots$ terminates. Thus there exists $n > 0$ such that $S(s_n \cdots s_1) = S(s_{n+1} s_n \cdots s_1) = \dots$. It follows that $\ker(s_n \cdots s_1) = \ker(s_{n+1} s_n \cdots s_1) = \dots$.

(2) \Rightarrow (1). First we prove that $S/W(S)$ is von Neumann regular. Let $s_1 \notin W(S)$. Then $\ker(s_1)$ is not essential in M . By Lemma 2.7, there exists $t_1 \in S$ such that $\ker(s_1) \subset \ker(s_1 - s_1 t_1 s_1)$ is proper. Put $s_2 = s_1 - s_1 t_1 s_1$. If $s_2 \in W(S)$, then we have $\overline{s_1} = \overline{s_1} \cdot \overline{t_1} \cdot \overline{s_1}$ in the ring $S/W(S)$. If $s_2 \notin W(S)$, then there exists $s_3 \in S$ such that $\ker(s_2) \subset \ker(s_3)$ is proper, where $s_3 = s_2 - s_2 t_2 s_2$ for some $t_2 \in S$ by the preceding proof. Repeating the above process, we get a strictly ascending chain

$$\ker(s_1) \subset \ker(s_2) \subset \ker(s_3) \subset \dots, \quad (2.1)$$

where $s_{i+1} = s_i - s_i t_i s_i$ for some $t_i \in S$, $i = 1, 2, \dots$. Let $u_1 = s_1$, $u_2 = 1 - s_1 t_1$, $u_3 = 1 - s_2 t_2, \dots, u_{i+1} = 1 - s_i t_i, \dots$. Then $s_1 = u_1, s_2 = u_2 u_1, s_3 = u_3 u_2 u_1, \dots, s_{i+1} = u_{i+1} u_i \cdots u_2 u_1, \dots$,

whence we have the following strict ascending chain

$$\ker(u_1) \subset \ker(u_2 u_1) \subset \ker(u_3 u_2 u_1) \subset \cdots, \quad (2.2)$$

which contradicts the hypothesis. So there exists a positive integer n such that $s_{n+1} \in W(S)$. This shows that \bar{s}_n is a regular element of $S/W(S)$, and hence $\bar{s}_{n-1}, \bar{s}_{n-2}, \dots, \bar{s}_1$ are regular elements of $S/W(S)$. Thus $S/W(S)$ is regular.

Note that since M_R is APQ-injective, $J(S) \subseteq W(S)$ by Theorem 2.5(1). Since the chain $\ker(s_1) \subseteq \ker(s_2 s_1) \subseteq \cdots$ terminates, by Lemma 2.8(1), $W(S)$ is right T -nilpotent, and so it follows that $W(S) \subseteq J(S)$, and thus $S/J(S)$ is regular. By Lemma 2.8, we get that S is right perfect. \square

By Lemma 2.8 (1) and [7, Remark 2], we have the following lemma.

LEMMA 2.10. *Let M be a right R -module with $S = \text{End}(M_R)$. If M_R satisfies ACC on $\{r_M(A) \mid A \subseteq S\}$, then $W(S)$ is nilpotent.*

The next corollary follows from Theorem 2.9 and Lemma 2.10.

COROLLARY 2.11. *Let M_R be APQ-injective with $S = \text{End}(M_R)$. If M_R satisfies ACC on $\{r_M(A) \mid A \subseteq S\}$, then S is semiprimary.*

For a module M_R , a submodule X of M is called a kernel submodule if $X = \ker(f)$ for some $f \in \text{End}(M_R)$, and X is called an annihilator submodule if $X = \bigcap_{f \in A} \ker(f)$ for some $A \subseteq \text{End}(M_R)$.

COROLLARY 2.12. *Let M_R be an APQ-injective module and $S = \text{End}(M_R)$. Then*

- (1) *if M_R satisfies ACC on kernel submodules, then S is right perfect;*
- (2) *if M_R satisfies ACC on annihilator submodules, then S is semiprimary.*

3. AQP-injective modules

In this section we study a generalization of quasiprincipally injective modules.

Definition 3.1. Let M be a right R -module with $S = \text{End}(M_R)$. Then M is said to be almost quasiprincipally injective (briefly, AQP-injective) if, for any $s \in S$, there exists a left ideal X_s of S such that $l_S(\ker(s)) = Ss \bigoplus X_s$ as left S -modules.

The next result gives the relationship between the AQP-injectivity of a module and the AP-injectivity of its endomorphism ring.

THEOREM 3.2. *Let M_R be a right R -module with $S = \text{End}(M_R)$. Then*

- (1) *if S is right AP-injective, then M_R is AQP-injective,*
- (2) *if M_R is AQP-injective and M generates $\ker(s)$ for each $s \in S$, then S is right AP-injective.*

Proof. (1) Let $s \in S$. Since S is right AP-injective, there exists a left ideal I_s such that $l_Sr_S(s) = Ss \bigoplus I_s$. If $a \in l_S(\ker(s))$ and $b \in r_S(s)$, then $sb = 0$, so $bM \subseteq \ker(s)$, and hence $abM = 0$, that is, $ab = 0$. It follows that $l_S(\ker(s)) \subseteq l_Sr_S(s)$. Thus, we have $Ss \subseteq l_S(\ker(s)) \subseteq Ss \bigoplus I_s$. This shows that $l_S(\ker(s)) = Ss \bigoplus l_S(\ker(s)) \cap I_s$, and (1) is proved.

(2) Let $0 \neq s \in S$. As M_R is AQP-injective, $l_S(\ker(s)) = Ss \oplus X_s$ for some left ideal X_s of S . Assume $a \in l_S r_S(s)$. Since M generates $\ker(s)$, $\ker(s) = \sum_{t \in T} t(M)$ for some subset T of S . It is easy to see that $at = 0$ for each $t \in T$, thus $ax = 0$ for each $x \in \ker(s)$. This implies that $l_S r_S(s) \subseteq l_S(\ker(s))$, from which we have $Ss \subseteq l_S r_S(s) \subseteq Ss \oplus X_s$, and hence $l_S r_S(s) = Ss \oplus (l_S r_S(s) \cap X_s)$. Therefore, S is right AP-injective. \square

THEOREM 3.3. *Let M be a right R -module with $S = \text{End}(M_R)$. If M is an AQP-injective module which is a self-generator, then $J(S) = W(S)$.*

Proof. Let $s \in J(S)$. Then we will show that $s \in W(S)$. If not, then there exists a nonzero submodule K of M such that $\ker(s) \cap K = 0$. As M is a self-generator, $K = \sum_{t \in I} t(M)$ for some subset I of S , hence we have some $0 \neq t \in I$ such that $\ker(s) \cap t(M) = 0$. Clearly, $st \neq 0$ and $\ker(st) = \ker(t)$. Since M is AQP-injective, $l_S(\ker(st)) = S(st) \oplus X_{st}$ as left S -modules. Now $t \in l_S(\ker(t)) = l_S(\ker(st)) = S(st) \oplus X_{st}$. Write $t = u(st) + v$, where $u \in S$ and $v \in X_{st}$. Then $st - su(st) = sv \in S(st) \cap X_{st}$, hence $st - su(st) = 0$, that is, $(1 - su)st = 0$. Note that $1 - su$ is left invertible, so $st = 0$, a contradiction.

Conversely, let $s \in W(S)$. Then, for each $t \in S$, $ts \in W(S)$ and so $1 - ts \neq 0$. Since M_R is AQP-injective, $l_S(\ker(1 - ts)) = S(1 - ts) \oplus X_{1-ts}$ as left S -modules. Note that $\ker(ts) \cap \ker(1 - ts) = 0$, so we have $\ker(1 - ts) = 0$, thus $S = S(1 - ts) \oplus X_{1-ts}$, and then $1 = e + x$ for some $e \in S(1 - ts)$ and $x \in X$. It follows that $e^2 = e$ and $Se = S(1 - ts)$, and so $1 - ts = ue$ for some $u \in S$. Since $\ker(ts)$ is essential in M_R , if $e \neq 1$, there is a nonzero element $(1 - e)m \in (1 - e)M \cap \ker(ts)$. Then $(1 - ts)(1 - e)m = (1 - e)m$. But $(1 - ts)(1 - e)m = ue(1 - e)m = 0$. This is a contradiction. So $e = 1$ and hence $1 - ts$ is left invertible. The result follows. \square

Recall that a module M_R is said to satisfy the C_2 -condition if every submodule of M that is isomorphic to a direct summand of M is itself a direct summand of M . A module M is said to satisfy the C_3 -condition if whenever M_1 and M_2 are two summands of M and $M_1 \cap M_2 = 0$, then $M_1 \oplus M_2$ is a summand of M . It is well known that the C_2 -condition implies the C_3 -condition.

THEOREM 3.4. *If M_R is an AQP-injective module, then it satisfies the C_2 -condition. In particular, right AP-injective rings are right C_2 -rings.*

Proof. Let A be a direct summand of M with $A \cong B$ and $S = \text{End}(M_R)$. Let $A = eM$, let $e^2 = e \in S$, and let $\varphi : eM \rightarrow B$ be an isomorphism. Then $B = bM$ with $b = se$ for some $s \in S$, and $\ker(e) = \ker(b)$. Thus, $e \in l_S(\ker(e)) = l_S(\ker(b)) = Sb \oplus X_b$ as M_R is AQP-injective, where X_b is a left S -module. Then $e = tb + x$ with $t \in S$ and $x \in X_b$. Hence we have $b = be = btb + bx$, and thus $b = btb$. Let $f = bt$. Then $f^2 = f$ and $bM = fM$. \square

COROLLARY 3.5. *Let M be a quasiprojective right R -module and let $S = \text{End}(M_R)$. Then S is regular if and only if M_R is AQP-injective and $\text{im}(s)$ are M -projective for every $s \in S$.*

Proof. By combining Theorems 3.2, 3.4, and [9, Theorem 37.7], one can complete the proof. \square

Recall that a ring R is called right P.P. if every principally right ideal of R is projective.

COROLLARY 3.6. *A ring R is regular if and only if R is right P.P. and right AP-injective.*

Following [3], a module M is said to be weakly injective if, for any finitely generated submodule $N \subseteq E(M)$, we have $N \subseteq X \cong M$ for some $X \subseteq E(M)$.

COROLLARY 3.7. *Let M_R be an f.g. module. If M is weakly injective and AQP-injective, then M is injective. In particular, if R is a right AP-injective and a right weakly injective ring, then R is right self-injective.*

Proof. Let $x \in E(M)$. Then there exists $X \subseteq E(M)$ such that $M + xR \subseteq X \cong M$, hence X is AQP-injective, and so $M \mid X$ by Theorem 3.4. This shows that $M = X$, so $x \in M$. \square

We let $S = \text{End}(M_R)$. Following [7], an element $u \in S$ is called a right uniform element of S if $u \neq 0$ and $u(M)$ is a uniform submodule of M . In the following, we generalize some results on maximal left ideals of the endomorphism rings of quasiprincipally injective modules and on maximal right ideals of right AP-injective rings to maximal left ideals of the endomorphism rings of AQP-injective modules.

LEMMA 3.8. *Let M_R be a module with $S = \text{End}(M_R)$. Given a set $\{X_s \mid s \in S\}$ of left ideals of S , the following are equivalent.*

- (1) $l_S(\ker(s)) = Ss \bigoplus X_s$ for all $s \in S$.
- (2) $l_S(tM \cap \ker(s)) = (X_{st} : t)_l + Ss$ and $(X_{st} : t)_l \cap Ss \subseteq l_S(t)$ for all $s, t \in S$, where $(X_{st} : t)_l = \{x \in S \mid xt \in X_{st}\}$.

Proof. (1) \Rightarrow (2). Let $x \in l_S(tM \cap \ker(s))$. Then $\ker(st) \subseteq \ker(xt)$ and so $xt \in l_S(\ker(xt)) \subseteq l_S(\ker(st)) = S(st) \bigoplus X_{st}$. Write $xt = s_1(st) + y$, where $s_1 \in S$ and $y \in X_{st}$, then $(x - s_1s)t = y \in X_{st}$ and hence $x - s_1s \in (X_{st} : t)_l$. It follows that $x \in (X_{st} : t)_l + Ss$. Obviously, $Ss \subseteq l_S(tM \cap \ker(s))$. If $z \in (X_{st} : t)_l$, then $zt \in X_{st} \subseteq l_S(\ker(st))$. Let $tm \in tM \cap \ker(s)$, then $stm = 0$, hence $ztm = 0$. This shows that $z \in l_S(tM \cap \ker(s))$. Therefore, $l_S(tM \cap \ker(s)) = (X_{st} : t)_l + Ss$. If $s' \in (X_{st} : t)_l \cap Ss$, then $s'st \in X_{st} \cap S(st) = 0$, and thus $s's \in l_S(t)$.

(2) \Rightarrow (1). Let $t = 1$. \square

LEMMA 3.9. *Let M_R be an AQP-injective module with $S = \text{End}(M_R)$ and an index set $\{X_s \mid s \in S\}$ of ideals such that $X_{st} = X_{ts}$ for all $s, t \in S$. If $0 \neq u(M)$ is a uniform submodule of M , define $M_u = \{s \in S \mid \ker(s) \cap u(M) \neq 0\}$. Then M_u is the unique maximal left ideal of S which contains $\sum_{s \in S} (X_{su} : u)_l$.*

Proof. It is easy to see that M_u is a left ideal. Let $t \in (X_{su} : u)_l$, then $tu \in X_{su}$, and thus $tus \in X_{su} \cap S(us) = X_{us} \cap S(us)$, since $X_{su} = X_{us}$ is an ideal. Then $tus = 0$ and so $t \in M_u$ if $us \neq 0$. If $us = 0$, then $l_S(\ker(us)) = 0$, and so $X_{su} = X_{us} = 0$. This shows that $tu = 0$ and hence $t \in M_u$. Consequently, $(X_{su} : u)_l \subseteq M_u$ for all $s \in S$. Now if $s \notin M_u$, then $\ker(s) \cap uM = 0$, and so $S = (X_{su} : u)_l + Ss$ by Lemma 3.8, hence $S = M_u + Ss$, showing that M_u is a maximal left ideal.

Finally, let L be a left ideal of S such that $\sum_{s \in S} (X_{su} : u)_l \subseteq L \neq M_u$. Then, as above, $S = (X_{su} : u)_l + Ss$ for any $s \in L - M_u$. Therefore, $L = S$. \square

LEMMA 3.10. *Let M_R be AQP-injective with $S = \text{End}(M_R)$ and an index set $\{X_s \mid s \in S\}$ of ideals such that $X_{st} = X_{ts}$ for all $s, t \in S$ and let $W = u_1M \bigoplus u_2M \bigoplus \cdots \bigoplus u_nM$ be a direct sum of uniform submodules u_iM of M , where each $u_i \in S$. If $T \subseteq S$ is a maximal left ideal*

not of the form M_u for any $u \in S$ such that uM is uniform, then there is $t \in T$ such that $\ker(1-t) \cap W$ is essential in W .

Proof. Since $T \neq M_{u_1}$, let $\ker(a) \cap u_1M = 0$, $a \in T$, then $\ker(au_1) \subseteq \ker(u_1)$, and so $u_1 \in l_S(\ker(au_1)) = S(au_1) \oplus X_{au_1}$. Thus, there exists $s \in S$ such that $(1-sa)u_1 \in X_{au_1}$, and so $1-sa \in (X_{au_1} : u_1)_l \subseteq M_{u_1}$. Let $a_1 = sa$. If $1-a_1 \in M_{u_i}$ for all i , we are done. If, say, $1-a_1 \notin M_2$, then $(1-a_1)u_2M$ is uniform (being isomorphic to u_2M), so, as above, $(1-a') \in M_{(1-a_1)u_2}$ for some $a' \in T$. Let $a_2 = a'ta_1 - a'a_1$, then $1-a_2 \in M_{u_1} \cap M_{u_2}$, continue in this way to obtain $t \in S$, such that $\ker(1-t) \cap u_iM \neq 0$ for each i , Lemma 3.10 follows. \square

THEOREM 3.11. *Let M_R be a self-generator with finite Goldie dimension and $S = \text{End}(M_R)$. If M_R is AQP-injective with an index set $\{X_s \mid s \in S\}$ of left ideals of S such that $X_{st} = X_{ts}$ for all $s, t \in S$, then*

- (1) *if T is a maximal left ideal of S , then $T = M_u$ for some $u \in S$ such that uM is a uniform submodule of M ,*
- (2) *$S/J(S)$ is semisimple.*

Proof. Since M is a self-generator, every uniform submodule of M contains an M -cyclic submodule. Therefore, we can assume that $W = u_1M \oplus u_2M \oplus \cdots \oplus u_nM$ is essential as M_R has finite Goldie dimension. If T is not of the form A_u for some right uniform element of $u \in S$, then by Lemma 3.10, there exists some $t \in T$ such that $\ker(1-t) \cap W$ is essential in W , so $\ker(1-t)$ is essential in M . By Theorem 3.3, $1-t \in J(S) \subseteq T$, a contradiction. This proves (1). As to (2), if $s \in M_{u_1} \cap \cdots \cap M_{u_n}$, then $\ker(s) \cap u_iM \neq 0$ for each i , whence $\ker(s)$ is essential in M . Hence, $s \in J(S)$, proving (2). \square

Acknowledgment

The authors are grateful to the referee for useful suggestions and remarks.

References

- [1] F. W. Anderson and K. R. Fuller, *Rings and Categories of Modules*, Springer, New York, 1974.
- [2] N. Ding, M. F. Yousif, and Y. Zhou, *Modules with annihilator conditions*, Comm. Algebra **30** (2002), no. 5, 2309–2320.
- [3] S. K. Jain and S. R. López-Permouth, *Rings whose cyclics are essentially embeddable in projective modules*, J. Algebra **128** (1990), no. 1, 257–269.
- [4] W. K. Nicholson, J. K. Park, and M. F. Yousif, *Principally quasi-injective modules*, Comm. Algebra **27** (1999), no. 4, 1683–1693.
- [5] W. K. Nicholson and M. F. Yousif, *Principally injective rings*, J. Algebra **174** (1995), no. 1, 77–93.
- [6] S. S. Page and Y. Zhou, *Generalizations of principally injective rings*, J. Algebra **206** (1998), no. 2, 706–721.
- [7] N. V. Sanh and K. P. Shum, *Endomorphism rings of quasi-principally injective modules*, Comm. Algebra **29** (2001), no. 4, 1437–1443.
- [8] N. V. Sanh, K. P. Shum, S. Dhompongsa, and S. Wongwai, *On quasi-principally injective modules*, Algebra Colloq. **6** (1999), no. 3, 269–276.

[9] R. Wisbauer, *Foundations of Module and Ring Theory*, Algebra, Logic and Applications, vol. 3, Gordon and Breach Science, Pennsylvania, 1991.

Zhu Zhanmin: Department of Mathematics, Jiaxing University, Jiaxing, Zhejiang 314001, China
E-mail address: zhanmin_zhu@hotmail.com

Xia Zhangsheng: Department of Mathematics, Hubei Institute for Nationalities, Enshi, Hubei 445000, China
E-mail address: xzsw8577@163.com

Tan Zhisong: Department of Mathematics, Hubei Institute for Nationalities, Enshi, Hubei 445000, China
E-mail address: es_tzs@hotmail.com

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru