

ON THE PERIODIC NATURE OF SOME MAX-TYPE DIFFERENCE EQUATIONS

E. M. ELABBASY, H. EL-METWALLY, AND E. M. ELSAYED

Received 10 April 2005 and in revised form 19 June 2005

We study some qualitative behavior of solutions of some max-type difference equations with periodic coefficients. Some new results of the periodicity character of solutions of that type of difference equations will be established.

1. Introduction

Recently there has been a lot of interest in studying the global attractivity, the boundedness character, and the periodicity nature of nonlinear difference equations. In [5, 6, 8] some global convergence results were established which can be applied to nonlinear difference equations in proving that every solution of these equations converges to a periodic solution (which need not necessarily be stable). The periodic nature of nonlinear difference equations of the max type has been investigated by many authors. See for example [1, 2, 3, 4].

Our main objective in this paper is to extend the study of boundedness and periodicity to solutions of some max-type difference equations. We deal with the following difference equation:

$$x_{n+1} = \max \left\{ \frac{1}{x_n}, \frac{A_n}{x_{n-1}} \right\}, \quad n = 0, 1, \dots, \quad (1.1)$$

where $\{A_n\}_{n=0}^{\infty} = \{\dots, \alpha, \beta, \alpha, \beta, \dots\}$ is a periodic sequence of positive numbers of period two with $\beta > \alpha > 1$. The case where $\{A_n\}_{n=0}^{\infty}$ is a periodic sequence of positive numbers of period three and $A_n \in (0, 1]$ was investigated in [4].

2. Invariant interval and boundedness

In this section, we show that every solution of (1.1) is bounded and persists.

The following lemmas are quite important results in their own; however these lemmas will be used in the subsequent discussion.

LEMMA 2.1. *Every positive solution of (1.1) is bounded and persists.*

Proof. Let $\{x_n\}_{n=-1}^{\infty}$ be a solution of (1.1). It follows from (1.1) for an integer number $N \geq 0$ that

$$x_{n+1}x_n \geq 1, \quad x_{n+1}x_{n-1} \geq \alpha > 1 \quad \forall n \geq N. \quad (2.1)$$

Thus

$$\min \{x_{n+1}x_n, x_{n+1}x_{n-1}\} \geq 1 \quad (2.2)$$

or

$$x_{n+1} \min \{x_n, x_{n-1}\} \geq 1 \quad \forall n \geq N. \quad (2.3)$$

That is, there exists a positive real number m such that

$$x_n \geq m \quad \forall n \geq N. \quad (2.4)$$

Thus from (1.1), we see that

$$\begin{aligned} x_{n+1} &= \max \left\{ \frac{1}{x_n}, \frac{A_n}{x_{n-1}} \right\} \\ &\leq \max \left\{ \frac{1}{m}, \frac{A_n}{m} \right\} = M. \end{aligned} \quad (2.5)$$

Hence

$$x_n \leq M \quad \forall n \geq N. \quad (2.6)$$

Thus from inequalities (2.4) and (2.6) we get

$$0 < m \leq x_n \leq M < \infty \quad \forall n \geq N. \quad (2.7)$$

Therefore every solution of (1.1) is bounded and persists. \square

LEMMA 2.2. *Assume that $\{x_n\}_{n=-1}^{\infty}$ is a positive solution of (1.1). Suppose there exists $N \geq 0$ such that*

$$x_{N-1}, x_N \in \left[\frac{1}{\sqrt{\alpha}}, \beta\sqrt{\alpha} \right] \quad \text{for some } N \geq 0. \quad (2.8)$$

Then

$$x_n \in \left[\frac{\sqrt{\alpha}}{\beta}, \beta\sqrt{\alpha} \right] \quad \forall n \geq N. \quad (2.9)$$

Proof. Observe from (1.1) that

$$\begin{aligned} x_{N+1} &= \max \left\{ \frac{1}{x_N}, \frac{A_N}{x_{N-1}} \right\} \geq \max \left\{ \frac{1}{\beta\sqrt{\alpha}}, \frac{\alpha}{\beta\sqrt{\alpha}} \right\} = \frac{\sqrt{\alpha}}{\beta}, \\ x_{N+1} &\leq \max \{ \sqrt{\alpha}, \alpha\sqrt{\alpha} \} = \alpha\sqrt{\alpha} < \beta\sqrt{\alpha}. \end{aligned} \quad (2.10)$$

Then

$$\frac{\sqrt{\alpha}}{\beta} \leq x_{N+1} < \beta\sqrt{\alpha}. \quad (2.11)$$

Again

$$\begin{aligned} x_{N+2} &= \max \left\{ \frac{1}{x_{N+1}}, \frac{A_{N+1}}{x_N} \right\} \geq \max \left\{ \frac{1}{\beta\sqrt{\alpha}}, \frac{\beta}{\beta\sqrt{\alpha}} \right\} = \frac{1}{\sqrt{\alpha}}, \\ x_{N+2} &\leq \max \left\{ \frac{\beta}{\sqrt{\alpha}}, \beta\sqrt{\alpha} \right\} = \beta\sqrt{\alpha}. \end{aligned} \quad (2.12)$$

Then

$$\frac{1}{\sqrt{\alpha}} \leq x_{N+2} \leq \beta\sqrt{\alpha}. \quad (2.13)$$

Also we see from (1.1) that

$$\begin{aligned} x_{N+3} &= \max \left\{ \frac{1}{x_{N+2}}, \frac{A_{N+2}}{x_{N+1}} \right\} \geq \max \left\{ \frac{1}{\beta\sqrt{\alpha}}, \frac{\alpha}{\alpha\sqrt{\alpha}} \right\} = \frac{1}{\sqrt{\alpha}}, \\ x_{N+3} &\leq \max \left\{ \sqrt{\beta}, \beta\sqrt{\alpha} \right\} = \beta\sqrt{\alpha}. \end{aligned} \quad (2.14)$$

Then

$$\frac{1}{\sqrt{\alpha}} \leq x_{N+3} \leq \beta\sqrt{\alpha}. \quad (2.15)$$

Thus following the above procedure we have

$$x_n \in \left[\frac{\sqrt{\alpha}}{\beta}, \beta\sqrt{\alpha} \right] \quad \forall n \geq N. \quad (2.16)$$

The proof is complete. \square

LEMMA 2.3. *Every solution of (1.1) which is bounded below by $1/\sqrt{\alpha}$ lies in the interval $[1/\sqrt{\alpha}, \beta\sqrt{\alpha}]$.*

Proof. Let $\{x_n\}_{n=-1}^{\infty}$ be a positive solution of (1.1) and there exists $N \geq 0$ such that

$$x_{n-1} \geq \frac{1}{\sqrt{\alpha}} \quad \forall n \geq N. \quad (2.17)$$

It follows from (1.1) that

$$x_{N+1} = \max \left\{ \frac{1}{x_N}, \frac{A_N}{x_{N-1}} \right\} \leq \max \{ \sqrt{\alpha}, \sqrt{\alpha}A_N \} \leq \beta\sqrt{\alpha}. \quad (2.18)$$

Similarly, we see that

$$x_{N+2} = \max \left\{ \frac{1}{x_{N+1}}, \frac{A_{N+1}}{x_N} \right\} \leq \max \{ \sqrt{\alpha}, \sqrt{\alpha}A_{N+1} \} \leq \beta\sqrt{\alpha}. \quad (2.19)$$

The rest of the proof follows by Lemma 2.2. \square

3. The main result

In this section, we study the periodicity character of solutions of (1.1).

In the following we study the existence of periodic solutions of (1.1) with period four.

THEOREM 3.1. *Assume that $\{x_n\}_{n=-1}^{\infty}$ is a positive solution of (1.1) with*

$$\frac{1}{\sqrt{\alpha}} < x_{N-1}, \quad x_N < \sqrt{\beta}. \quad (3.1)$$

Then $\{x_n\}_{n=-1}^{\infty}$ is a four-cycle solution of (1.1).

Proof. Let $\{x_n\}_{n=-1}^{\infty}$ be a positive solution of (1.1). Suppose there exists $N \geq 0$ such that

$$\frac{1}{\sqrt{\alpha}} < x_{N-1}, \quad x_N < \sqrt{\beta}. \quad (3.2)$$

Assume that

$$x_{N-1} = p, \quad x_N = q. \quad (3.3)$$

Observe from (1.1) that

$$x_{N+1} = \max \left\{ \frac{1}{x_N}, \frac{A_N}{x_{N-1}} \right\}. \quad (3.4)$$

We consider the following two cases.

(1) $x_{N+1} = 1/x_N = 1/q$. In this case $1/x_N > \alpha/x_{N-1}$, (the case $1/x_N > \beta/x_{N-1}$ can be treated similarly) and we see that

$$\begin{aligned} x_{N+2} &= \max \left\{ \frac{1}{x_{N+1}}, \frac{A_{N+1}}{x_N} \right\} = \max \left\{ q, \frac{\beta}{q} \right\} = \frac{\beta}{q}, \\ x_{N+3} &= \max \left\{ \frac{1}{x_{N+2}}, \frac{A_{N+2}}{x_{N+1}} \right\} = \max \left\{ \frac{q}{\beta}, \alpha q \right\} = \alpha q, \\ x_{N+4} &= \max \left\{ \frac{1}{x_{N+3}}, \frac{A_{N+3}}{x_{N+2}} \right\} = \max \left\{ \frac{1}{\alpha q}, q \right\} = q, \\ x_{N+5} &= \max \left\{ \frac{1}{x_{N+4}}, \frac{A_{N+4}}{x_{N+3}} \right\} = \max \left\{ \frac{1}{q}, \frac{1}{q} \right\} = \frac{1}{q}, \\ x_{N+6} &= \max \left\{ \frac{1}{x_{N+5}}, \frac{A_{N+5}}{x_{N+4}} \right\} = \max \left\{ q, \frac{\beta}{q} \right\} = \frac{\beta}{q}. \end{aligned} \quad (3.5)$$

Then clearly the solution becomes in the form

$$\left\{ \dots, q, \frac{1}{q}, \frac{\beta}{q}, \alpha q, q, \frac{1}{q}, \frac{\beta}{q}, \alpha q, \dots \right\}. \quad (3.6)$$

(2) $x_{N+1} = \alpha/x_{N-1} = \alpha/p$. In this case we see that

$$\begin{aligned} x_{N+2} &= \max \left\{ \frac{1}{x_{N+1}}, \frac{A_{N+1}}{x_N} \right\} = \max \left\{ \frac{p}{\alpha}, \frac{\beta}{q} \right\} = \frac{\beta}{q}, \\ x_{N+3} &= \max \left\{ \frac{1}{x_{N+2}}, \frac{A_{N+2}}{x_{N+1}} \right\} = \max \left\{ \frac{q}{\beta}, p \right\} = p, \end{aligned} \quad (3.7)$$

where $x_{N-1} > 1/\sqrt{\beta} \Rightarrow \beta x_{N-1} > \sqrt{\beta} > x_N$,

$$\begin{aligned} x_{N+4} &= \max \left\{ \frac{1}{x_{N+3}}, \frac{A_{N+3}}{x_{N+2}} \right\} = \max \left\{ \frac{1}{p}, q \right\} = q, \\ x_{N+5} &= \max \left\{ \frac{1}{x_{N+4}}, \frac{A_{N+4}}{x_{N+3}} \right\} = \max \left\{ \frac{1}{q}, \frac{\alpha}{p} \right\} = \frac{\alpha}{p}, \\ x_{N+6} &= \max \left\{ \frac{1}{x_{N+5}}, \frac{A_{N+5}}{x_{N+4}} \right\} = \max \left\{ \frac{p}{\alpha}, \frac{\beta}{q} \right\} = \frac{\beta}{q}, \end{aligned} \quad (3.8)$$

and so the solution becomes in the form

$$\left\{ \dots, p, q, \frac{\alpha}{p}, \frac{\beta}{q}, p, q, \frac{\alpha}{p}, \frac{\beta}{q}, \dots \right\}. \quad (3.9)$$

□

The proof is complete.

THEOREM 3.2. *Every positive solution of (1.1) which is bounded from below by $1/\sqrt{\alpha}$ is eventually periodic with period four.*

Proof. Let $\{x_n\}_{n=-1}^{\infty}$ be a positive solution of (1.1). By Lemma 2.3, we assume

$$\frac{1}{\sqrt{\alpha}} < x_{N-1}, \quad x_N < \beta\sqrt{\alpha} \text{ for some integer } N \geq 2. \quad (3.10)$$

From (1.1), we see that

$$x_{N+1} = \max \left\{ \frac{1}{x_N}, \frac{\alpha}{x_{N-1}} \right\}. \quad (3.11)$$

We consider the following two cases.

(A₁) $x_{N+1} = 1/x_N$. In this case $1/x_N > \alpha/x_{N-1}$, and we see that

$$x_{N+2} = \max \left\{ \frac{1}{x_{N+1}}, \frac{A_{N+1}}{x_N} \right\} = \max \left\{ x_N, \frac{\beta}{x_N} \right\}. \quad (3.12)$$

We consider the following two cases.

(A₁₁) $x_{N+2} = x_N$. In this case $x_N > \beta/x_N$, and we see that

$$x_{N+3} = \max \left\{ \frac{1}{x_{N+2}}, \frac{A_{N+2}}{x_{N+1}} \right\} = \max \left\{ \frac{1}{x_N}, \alpha x_N \right\} = \alpha x_N, \quad (3.13)$$

where $x_N > 1/\sqrt{\alpha} \Rightarrow \alpha x_N > 1/x_N$,

$$\begin{aligned} x_{N+4} &= \max \left\{ \frac{1}{x_{N+3}}, \frac{A_{N+3}}{x_{N+2}} \right\} = \max \left\{ \frac{1}{\alpha x_N}, \frac{\beta}{x_N} \right\} = \frac{\beta}{x_N}, \\ x_{N+5} &= \max \left\{ \frac{1}{x_{N+4}}, \frac{A_{N+4}}{x_{N+3}} \right\} = \max \left\{ \frac{x_N}{\beta}, \frac{1}{x_N} \right\} = \frac{x_N}{\beta}, \\ x_{N+6} &= \max \left\{ \frac{1}{x_{N+5}}, \frac{A_{N+5}}{x_{N+4}} \right\} = \max \left\{ \frac{\beta}{x_N}, x_N \right\} = x_N, \\ x_{N+7} &= \max \left\{ \frac{1}{x_{N+6}}, \frac{A_{N+6}}{x_{N+5}} \right\} = \max \left\{ \frac{1}{x_N}, \frac{\alpha\beta}{x_N} \right\} = \frac{\alpha\beta}{x_N}, \\ x_{N+8} &= \max \left\{ \frac{1}{x_{N+7}}, \frac{A_{N+7}}{x_{N+6}} \right\} = \max \left\{ \frac{x_N}{\alpha\beta}, \frac{\beta}{x_N} \right\} = \frac{\beta}{x_N}, \\ x_{N+9} &= \max \left\{ \frac{1}{x_{N+8}}, \frac{A_{N+8}}{x_{N+7}} \right\} = \max \left\{ \frac{x_N}{\beta}, \frac{x_N}{\beta} \right\} = \frac{x_N}{\beta}, \\ x_{N+10} &= \max \left\{ \frac{1}{x_{N+9}}, \frac{A_{N+9}}{x_{N+8}} \right\} = \max \left\{ \frac{\beta}{x_N}, x_N \right\} = x_N, \\ x_{N+11} &= \max \left\{ \frac{1}{x_{N+10}}, \frac{A_{N+10}}{x_{N+9}} \right\} = \max \left\{ \frac{1}{x_N}, \frac{\alpha\beta}{x_N} \right\} = \frac{\alpha\beta}{x_N}, \\ x_{N+12} &= \max \left\{ \frac{1}{x_{N+11}}, \frac{A_{N+11}}{x_{N+10}} \right\} = \max \left\{ \frac{x_N}{\alpha\beta}, \frac{\beta}{x_N} \right\} = \frac{\beta}{x_N}. \end{aligned} \quad (3.14)$$

We see that the solution is in the form

$$\left\{ \dots, \frac{\beta}{x_N}, \frac{x_N}{\beta}, x_N, \frac{\alpha\beta}{x_N}, \frac{\beta}{x_N}, \frac{x_N}{\beta}, x_N, \frac{\alpha\beta}{x_N}, \dots \right\}. \quad (3.15)$$

Therefore $\{x_n\}_{n=-1}^{\infty}$ is a periodic solution with period four.

(A₁₂) $x_{N+2} = \beta/x_N$. In this case $\beta/x_N > x_N$, and we see that

$$\begin{aligned} x_{N+3} &= \max \left\{ \frac{1}{x_{N+2}}, \frac{A_{N+2}}{x_{N+1}} \right\} = \max \left\{ \frac{x_N}{\beta}, \alpha x_N \right\} = \alpha x_N, \\ x_{N+4} &= \max \left\{ \frac{1}{x_{N+3}}, \frac{A_{N+3}}{x_{N+2}} \right\} = \max \left\{ \frac{1}{\alpha x_N}, x_N \right\} = x_N, \end{aligned} \quad (3.16)$$

where $x_N > 1/\sqrt{\alpha} \Rightarrow x_N^2 > 1/\alpha \Rightarrow x_N > 1/\alpha x_N$,

$$\begin{aligned}
 x_{N+5} &= \max \left\{ \frac{1}{x_{N+4}}, \frac{A_{N+4}}{x_{N+3}} \right\} = \max \left\{ \frac{1}{x_N}, \frac{1}{x_N} \right\} = \frac{1}{x_N}, \\
 x_{N+6} &= \max \left\{ \frac{1}{x_{N+5}}, \frac{A_{N+5}}{x_{N+4}} \right\} = \max \left\{ x_N, \frac{\beta}{x_N} \right\} = \frac{\beta}{x_N}, \\
 x_{N+7} &= \max \left\{ \frac{1}{x_{N+6}}, \frac{A_{N+6}}{x_{N+5}} \right\} = \max \left\{ \frac{x_N}{\beta}, \alpha x_N \right\} = \alpha x_N, \\
 x_{N+8} &= \max \left\{ \frac{1}{x_{N+7}}, \frac{A_{N+7}}{x_{N+6}} \right\} = \max \left\{ \frac{1}{\alpha x_N}, x_N \right\} = x_N, \\
 x_{N+9} &= \max \left\{ \frac{1}{x_{N+8}}, \frac{A_{N+8}}{x_{N+7}} \right\} = \max \left\{ \frac{1}{x_N}, \frac{1}{x_N} \right\} = \frac{1}{x_N}, \\
 x_{N+10} &= \max \left\{ \frac{1}{x_{N+9}}, \frac{A_{N+9}}{x_{N+8}} \right\} = \max \left\{ x_N, \frac{\beta}{x_N} \right\} = \frac{\beta}{x_N}, \\
 x_{N+11} &= \max \left\{ \frac{1}{x_{N+10}}, \frac{A_{N+10}}{x_{N+9}} \right\} = \max \left\{ \frac{x_N}{\beta}, \alpha x_N \right\} = \alpha x_N, \\
 x_{N+12} &= \max \left\{ \frac{1}{x_{N+11}}, \frac{A_{N+11}}{x_{N+10}} \right\} = \max \left\{ \frac{1}{\alpha x_N}, x_N \right\} = x_N.
 \end{aligned} \tag{3.17}$$

Therefore $\{x_n\}_{n=-1}^{\infty}$ is a periodic solution with period four as follows:

$$\left\{ \dots, x_N, \frac{1}{x_N}, \frac{\beta}{x_N}, \alpha x_N, x_N, \frac{1}{x_N}, \frac{\beta}{x_N}, \alpha x_N, \dots \right\}. \tag{3.18}$$

(A₂) $x_{N+1} = \alpha/x_{N-1}$. In this case $\alpha/x_{N-1} > 1/x_N$, and we see that

$$x_{N+2} = \max \left\{ \frac{1}{x_{N+1}}, \frac{A_{N+1}}{x_N} \right\} = \max \left\{ \frac{x_{N-1}}{\alpha}, \frac{\beta}{x_N} \right\}. \tag{3.19}$$

We consider the following two cases.

(A₂₁) $x_{N+2} = x_{N-1}/\alpha$. In this case $x_{N-1}/\alpha > \beta/x_N$, and we see that

$$x_{N+3} = \max \left\{ \frac{1}{x_{N+2}}, \frac{A_{N+2}}{x_{N+1}} \right\} = \max \left\{ \frac{\alpha}{x_{N-1}}, x_{N-1} \right\} = x_{N-1}, \tag{3.20}$$

where $\beta\sqrt{\alpha}x_{N-1} > x_{N-1}x_N > \alpha\beta \Rightarrow x_{N-1} > \sqrt{\alpha}$,

$$\begin{aligned}
 x_{N+4} &= \max \left\{ \frac{1}{x_{N+3}}, \frac{A_{N+3}}{x_{N+2}} \right\} = \max \left\{ \frac{1}{x_{N-1}}, \frac{\alpha\beta}{x_{N-1}} \right\} = \frac{\alpha\beta}{x_{N-1}}, \\
 x_{N+5} &= \max \left\{ \frac{1}{x_{N+4}}, \frac{A_{N+4}}{x_{N+3}} \right\} = \max \left\{ \frac{x_{N-1}}{\alpha\beta}, \frac{\alpha}{x_{N-1}} \right\} = \frac{x_{N-1}}{\alpha\beta}.
 \end{aligned} \tag{3.21}$$

We consider the following two cases.

(A₂₁₁) $x_{N+5} = x_{N-1}/\alpha\beta$. In this case $x_{N-1}/\alpha\beta > \alpha/x_{N-1}$, and we see that

$$x_{N+6} = \max \left\{ \frac{1}{x_{N+5}}, \frac{A_{N+5}}{x_{N+4}} \right\} = \max \left\{ \frac{\alpha\beta}{x_{N-1}}, \frac{x_{N-1}}{\alpha} \right\} = \frac{x_{N-1}}{\alpha}, \tag{3.22}$$

where $x_{N-1}/\alpha\beta > \alpha/x_{N-1} \Rightarrow x_{N-1}/\alpha > \alpha\beta/x_{N-1}$,

$$\begin{aligned} x_{N+7} &= \max \left\{ \frac{1}{x_{N+6}}, \frac{A_{N+6}}{x_{N+5}} \right\} = \max \left\{ \frac{\alpha}{x_{N-1}}, \frac{\alpha^2\beta}{x_{N-1}} \right\} = \frac{\alpha^2\beta}{x_{N-1}}, \\ x_{N+8} &= \max \left\{ \frac{1}{x_{N+7}}, \frac{A_{N+7}}{x_{N+6}} \right\} = \max \left\{ \frac{x_{N-1}}{\alpha^2\beta}, \frac{\alpha\beta}{x_{N-1}} \right\} = \frac{\alpha\beta}{x_{N-1}}, \end{aligned} \quad (3.23)$$

where $x_{N-1} < \beta\sqrt{\alpha} \Rightarrow x_{N-1}^2 < \beta^2\alpha < \beta^2\alpha^3 \Rightarrow \alpha\beta/x_{N-1} > x_{N-1}/\alpha^2\beta$,

$$\begin{aligned} x_{N+9} &= \max \left\{ \frac{1}{x_{N+8}}, \frac{A_{N+8}}{x_{N+7}} \right\} = \max \left\{ \frac{x_{N-1}}{\alpha\beta}, \frac{x_{N-1}}{\alpha\beta} \right\} = \frac{x_{N-1}}{\alpha\beta}, \\ x_{N+10} &= \max \left\{ \frac{1}{x_{N+9}}, \frac{A_{N+9}}{x_{N+8}} \right\} = \max \left\{ \frac{\alpha\beta}{x_{N-1}}, \frac{x_{N-1}}{\alpha} \right\} = \frac{x_{N-1}}{\alpha}, \\ x_{N+11} &= \max \left\{ \frac{1}{x_{N+10}}, \frac{A_{N+10}}{x_{N+9}} \right\} = \max \left\{ \frac{\alpha}{x_{N-1}}, \frac{\alpha^2\beta}{x_{N-1}} \right\} = \frac{\alpha^2\beta}{x_{N-1}}, \\ x_{N+12} &= \max \left\{ \frac{1}{x_{N+11}}, \frac{A_{N+11}}{x_{N+10}} \right\} = \max \left\{ \frac{x_{N-1}}{\alpha^2\beta}, \frac{\alpha\beta}{x_{N-1}} \right\} = \frac{\alpha\beta}{x_{N-1}}. \end{aligned} \quad (3.24)$$

Therefore the solution can be written as

$$\left\{ \dots, \frac{\alpha\beta}{x_{N-1}}, \frac{x_{N-1}}{\alpha\beta}, \frac{x_{N-1}}{\alpha}, \frac{\alpha^2\beta}{x_{N-1}}, \frac{\alpha\beta}{x_{N-1}}, \frac{x_{N-1}}{\alpha\beta}, \frac{x_{N-1}}{\alpha}, \frac{\alpha^2\beta}{x_{N-1}}, \dots \right\}. \quad (3.25)$$

Then $\{x_n\}_{n=-1}^{\infty}$ is a periodic solution with period four.

We consider the following two cases.

(A₂₁₂) $x_{N+5} = \alpha/x_{N-1}$. In this case $\alpha/x_{N-1} > x_{N-1}/\alpha\beta$, and we see that

$$\begin{aligned} x_{N+6} &= \max \left\{ \frac{1}{x_{N+5}}, \frac{A_{N+5}}{x_{N+4}} \right\} = \max \left\{ \frac{x_{N-1}}{\alpha}, \frac{x_{N-1}}{\alpha} \right\} = \frac{x_{N-1}}{\alpha}, \\ x_{N+7} &= \max \left\{ \frac{1}{x_{N+6}}, \frac{A_{N+6}}{x_{N+5}} \right\} = \max \left\{ \frac{\alpha}{x_{N-1}}, x_{N-1} \right\} = x_{N-1}, \\ x_{N+8} &= \max \left\{ \frac{1}{x_{N+7}}, \frac{A_{N+7}}{x_{N+6}} \right\} = \max \left\{ \frac{1}{x_{N-1}}, \frac{\alpha\beta}{x_{N-1}} \right\} = \frac{\alpha\beta}{x_{N-1}}, \\ x_{N+9} &= \max \left\{ \frac{1}{x_{N+8}}, \frac{A_{N+8}}{x_{N+7}} \right\} = \max \left\{ \frac{x_{N-1}}{\alpha\beta}, \frac{\alpha}{x_{N-1}} \right\} = \frac{\alpha}{x_{N-1}}, \\ x_{N+10} &= \max \left\{ \frac{1}{x_{N+9}}, \frac{A_{N+9}}{x_{N+8}} \right\} = \max \left\{ \frac{x_{N-1}}{\alpha}, \frac{x_{N-1}}{\alpha} \right\} = \frac{x_{N-1}}{\alpha}, \\ x_{N+11} &= \max \left\{ \frac{1}{x_{N+10}}, \frac{A_{N+10}}{x_{N+9}} \right\} = \max \left\{ \frac{\alpha}{x_{N-1}}, x_{N-1} \right\} = x_{N-1}, \\ x_{N+12} &= \max \left\{ \frac{1}{x_{N+11}}, \frac{A_{N+11}}{x_{N+10}} \right\} = \max \left\{ \frac{1}{x_{N-1}}, \frac{\alpha\beta}{x_{N-1}} \right\} = \frac{\alpha\beta}{x_{N-1}}. \end{aligned} \quad (3.26)$$

It is also easy to see that the solution takes the form

$$\left\{ \dots, \frac{\alpha}{x_{N-1}}, \frac{x_{N-1}}{\alpha}, x_{N-1}, \frac{\alpha\beta}{x_{N-1}}, \frac{\alpha}{x_{N-1}}, \frac{x_{N-1}}{\alpha}, x_{N-1}, \frac{\alpha\beta}{x_{N-1}}, \dots \right\}, \quad (3.27)$$

which is periodic with period four.

(A₂₂) $x_{N+2} = \beta/x_N$. In this case $\beta/x_N > x_{N-1}/\alpha$, and we see that

$$x_{N+3} = \max \left\{ \frac{1}{x_{N+2}}, \frac{A_{N+2}}{x_{N+1}} \right\} = \max \left\{ \frac{x_N}{\beta}, x_{N-1} \right\}. \quad (3.28)$$

We consider the following two cases.

(A₂₂₁) $x_{N+3} = x_{N-1}$. In this case $x_{N-1} > x_N/\beta$, and we see that

$$\begin{aligned} x_{N+4} &= \max \left\{ \frac{1}{x_{N+3}}, \frac{A_{N+3}}{x_{N+2}} \right\} = \max \left\{ \frac{1}{x_{N-1}}, x_N \right\} = x_N, \\ x_{N+5} &= \max \left\{ \frac{1}{x_{N+4}}, \frac{A_{N+4}}{x_{N+3}} \right\} = \max \left\{ \frac{1}{x_N}, \frac{\alpha}{x_{N-1}} \right\} = \frac{\alpha}{x_{N-1}}, \\ x_{N+6} &= \max \left\{ \frac{1}{x_{N+5}}, \frac{A_{N+5}}{x_{N+4}} \right\} = \max \left\{ \frac{x_{N-1}}{\alpha}, \frac{\beta}{x_N} \right\} = \frac{\beta}{x_N}, \\ x_{N+7} &= \max \left\{ \frac{1}{x_{N+6}}, \frac{A_{N+6}}{x_{N+5}} \right\} = \max \left\{ \frac{x_N}{\beta}, x_{N-1} \right\} = x_{N-1}, \\ x_{N+8} &= \max \left\{ \frac{1}{x_{N+7}}, \frac{A_{N+7}}{x_{N+6}} \right\} = \max \left\{ \frac{1}{x_{N-1}}, x_N \right\} = x_N, \\ x_{N+9} &= \max \left\{ \frac{1}{x_{N+8}}, \frac{A_{N+8}}{x_{N+7}} \right\} = \max \left\{ \frac{1}{x_N}, \frac{\alpha}{x_{N-1}} \right\} = \frac{\alpha}{x_{N-1}}, \\ x_{N+10} &= \max \left\{ \frac{1}{x_{N+9}}, \frac{A_{N+9}}{x_{N+8}} \right\} = \max \left\{ \frac{x_{N-1}}{\alpha}, \frac{\beta}{x_N} \right\} = \frac{\beta}{x_N}, \\ x_{N+11} &= \max \left\{ \frac{1}{x_{N+10}}, \frac{A_{N+10}}{x_{N+9}} \right\} = \max \left\{ \frac{x_N}{\beta}, x_{N-1} \right\} = x_{N-1}, \\ x_{N+12} &= \max \left\{ \frac{1}{x_{N+11}}, \frac{A_{N+11}}{x_{N+10}} \right\} = \max \left\{ \frac{1}{x_{N-1}}, x_N \right\} = x_N. \end{aligned} \quad (3.29)$$

One can easily see that the solution will be in the form

$$\left\{ \dots, x_{N-1}, x_N, \frac{\alpha}{x_{N-1}}, \frac{\beta}{x_N}, x_{N-1}, x_N, \frac{\alpha}{x_{N-1}}, \frac{\beta}{x_N}, \dots \right\}, \quad (3.30)$$

and so the solution is periodic with period four.

(A_{222}) $x_{N+3} = x_N/\beta$. In this case $x_N/\beta > x_{N-1}$, and we see that

$$x_{N+4} = \max \left\{ \frac{1}{x_{N+3}}, \frac{A_{N+3}}{x_{N+2}} \right\} = \max \left\{ \frac{\beta}{x_N}, x_N \right\} = x_N, \quad (3.31)$$

where $x_N > \beta x_{N-1} > \beta/\sqrt{\alpha} > \beta/\sqrt{\beta} = \sqrt{\beta} \Rightarrow x_N^2 > \beta \Rightarrow x_N > \beta/x_N$,

$$\begin{aligned} x_{N+5} &= \max \left\{ \frac{1}{x_{N+4}}, \frac{A_{N+4}}{x_{N+3}} \right\} = \max \left\{ \frac{1}{x_N}, \frac{\alpha\beta}{x_N} \right\} = \frac{\alpha\beta}{x_N}, \\ x_{N+6} &= \max \left\{ \frac{1}{x_{N+5}}, \frac{A_{N+5}}{x_{N+4}} \right\} = \max \left\{ \frac{x_N}{\alpha\beta}, \frac{\beta}{x_N} \right\} = \frac{\beta}{x_N}, \end{aligned} \quad (3.32)$$

where $x_N < \beta\sqrt{\alpha}$,

$$\begin{aligned} x_{N+7} &= \max \left\{ \frac{1}{x_{N+6}}, \frac{A_{N+6}}{x_{N+5}} \right\} = \max \left\{ \frac{x_N}{\beta}, \frac{x_N}{\beta} \right\} = \frac{x_N}{\beta}, \\ x_{N+8} &= \max \left\{ \frac{1}{x_{N+7}}, \frac{A_{N+7}}{x_{N+6}} \right\} = \max \left\{ \frac{\beta}{x_N}, x_N \right\} = x_N, \\ x_{N+9} &= \max \left\{ \frac{1}{x_{N+8}}, \frac{A_{N+8}}{x_{N+7}} \right\} = \max \left\{ \frac{1}{x_N}, \frac{\alpha\beta}{x_N} \right\} = \frac{\alpha\beta}{x_N}, \\ x_{N+10} &= \max \left\{ \frac{1}{x_{N+9}}, \frac{A_{N+9}}{x_{N+8}} \right\} = \max \left\{ \frac{x_N}{\alpha\beta}, \frac{\beta}{x_N} \right\} = \frac{\beta}{x_N}, \\ x_{N+11} &= \max \left\{ \frac{1}{x_{N+10}}, \frac{A_{N+10}}{x_{N+9}} \right\} = \max \left\{ \frac{x_N}{\beta}, \frac{x_N}{\beta} \right\} = \frac{x_N}{\beta}, \\ x_{N+12} &= \max \left\{ \frac{1}{x_{N+11}}, \frac{A_{N+11}}{x_{N+10}} \right\} = \max \left\{ \frac{\beta}{x_N}, x_N \right\} = x_N. \end{aligned} \quad (3.33)$$

Then the solution can be written in the form

$$\left\{ \dots, \frac{\beta}{x_N}, \frac{x_N}{\beta}, x_N, \frac{\alpha\beta}{x_N}, \frac{\beta}{x_N}, \frac{x_N}{\beta}, x_N, \frac{\alpha\beta}{x_N}, \dots \right\}, \quad (3.34)$$

and so the solution is periodic with period four.

This completes the proof. The proof of Theorem 3.2 is thus completed. \square

LEMMA 3.3. *Assume $\{x_n\}_{n=-1}^{\infty}$ is a positive solution of (1.1) and suppose there exists $m \geq 2$ such that*

$$x_m < \frac{1}{\sqrt{\alpha}} < x_{m+1}. \quad (3.35)$$

Then either $\{x_n\}_{n=-1}^{\infty}$ is eventually periodic solution with period four or

$$\liminf_{n \rightarrow \infty} x_n \geq \frac{1}{\sqrt{\alpha}}. \quad (3.36)$$

Proof. Observe that $x_m < 1/\sqrt{\alpha}$ and either $x_{m+1} < \beta\sqrt{\alpha}$ or $x_{m+1} > \beta\sqrt{\alpha}$.

(i) Assume that $x_{m+1} < \beta\sqrt{\alpha}$. It follows from (1.1) that

$$x_{m+2} = \max \left\{ \frac{1}{x_{m+1}}, \frac{A_{m+1}}{x_m} \right\} = \max \left\{ \frac{1}{x_{m+1}}, \frac{\alpha}{x_m} \right\} = \frac{\alpha}{x_m}, \quad (3.37)$$

where $x_m x_{m+1} \geq 1 \Rightarrow x_{m+1} > \sqrt{\alpha} > 1$,

$$x_{m+3} = \max \left\{ \frac{1}{x_{m+2}}, \frac{A_{m+2}}{x_{m+1}} \right\} = \max \left\{ \frac{x_m}{\alpha}, \frac{\beta}{x_{m+1}} \right\} = \frac{\beta}{x_{m+1}}, \quad (3.38)$$

where $x_m x_{m+1} < \beta\sqrt{\alpha}/\sqrt{\alpha} = \beta < \alpha\beta$, and

$$x_{m+4} = \max \left\{ \frac{1}{x_{m+3}}, \frac{A_{m+3}}{x_{m+2}} \right\} = \max \left\{ \frac{x_{m+1}}{\beta}, x_m \right\}. \quad (3.39)$$

Then either

$$x_{m+4} = \frac{x_{m+1}}{\beta} \quad \text{or} \quad x_{m+4} = x_m \quad (3.40)$$

and by simple computations the solution becomes either

$$\left\{ \dots, \frac{x_{m+1}}{\beta}, x_{m+1}, \frac{\alpha\beta}{x_{m+1}}, \frac{\beta}{x_{m+1}}, \frac{x_{m+1}}{\beta}, x_{m+1}, \frac{\alpha\beta}{x_{m+1}}, \frac{\beta}{x_{m+1}}, \dots \right\}, \quad (3.41)$$

or

$$\left\{ \dots, x_m, x_{m+1}, \frac{\alpha}{x_m}, \frac{\beta}{x_{m+1}}, x_m, x_{m+1}, \frac{\alpha}{x_m}, \frac{\beta}{x_{m+1}}, \dots \right\}, \quad (3.42)$$

and so in either case $\{x_n\}_{n=-1}^{\infty}$ is a periodic solution with period four.

(ii) Assume that $x_{m+1} > \beta\sqrt{\alpha}$. In this case we see from (1.1) that

$$\begin{aligned} x_{m+2} &= \max \left\{ \frac{1}{x_{m+1}}, \frac{A_{m+1}}{x_m} \right\} = \max \left\{ \frac{1}{x_{m+1}}, \frac{\alpha}{x_m} \right\} = \frac{\alpha}{x_m}, \\ x_{m+3} &= \max \left\{ \frac{1}{x_{m+2}}, \frac{A_{m+2}}{x_{m+1}} \right\} = \max \left\{ \frac{x_m}{\alpha}, \frac{\beta}{x_{m+1}} \right\}. \end{aligned} \quad (3.43)$$

We consider the following two cases.

(B₁) $x_{m+3} = x_m/\alpha$. In this case we see that

$$\begin{aligned}
 x_{m+4} &= \max \left\{ \frac{1}{x_{m+3}}, \frac{A_{m+3}}{x_{m+2}} \right\} = \max \left\{ \frac{\alpha}{x_m}, x_m \right\} = \frac{\alpha}{x_m}, \\
 x_{m+5} &= \max \left\{ \frac{1}{x_{m+4}}, \frac{A_{m+4}}{x_{m+3}} \right\} = \max \left\{ \frac{x_m}{\alpha}, \frac{\alpha\beta}{x_m} \right\} = \frac{\alpha\beta}{x_m}, \\
 x_{m+6} &= \max \left\{ \frac{1}{x_{m+5}}, \frac{A_{m+5}}{x_{m+4}} \right\} = \max \left\{ \frac{x_m}{\alpha\beta}, x_m \right\} = x_m, \\
 x_{m+7} &= \max \left\{ \frac{1}{x_{m+6}}, \frac{A_{m+6}}{x_{m+5}} \right\} = \max \left\{ \frac{1}{x_m}, \frac{x_m}{\alpha} \right\} = \frac{1}{x_m}, \\
 x_{m+8} &= \max \left\{ \frac{1}{x_{m+7}}, \frac{A_{m+7}}{x_{m+6}} \right\} = \max \left\{ x_m, \frac{\alpha}{x_m} \right\} = \frac{\alpha}{x_m}, \\
 x_{m+9} &= \max \left\{ \frac{1}{x_{m+8}}, \frac{A_{m+8}}{x_{m+7}} \right\} = \max \left\{ \frac{x_m}{\alpha}, \beta x_m \right\} = \beta x_m, \\
 x_{m+10} &= \max \left\{ \frac{1}{x_{m+9}}, \frac{A_{m+9}}{x_{m+8}} \right\} = \max \left\{ \frac{1}{\beta x_m}, x_m \right\}.
 \end{aligned} \tag{3.44}$$

We consider the following two cases.

(B₁₁) $x_{m+10} = x_m$. In this case the solution eventually will be periodic with period four as

$$\left\{ \dots, x_m, \frac{1}{x_m}, \frac{\alpha}{x_m}, \beta x_m, x_m, \frac{1}{x_m}, \frac{\alpha}{x_m}, \beta x_m, \dots \right\}. \tag{3.45}$$

(B₁₂) $x_{m+10} = 1/\beta x_m$. In this case straightforward calculations show that the solution will be in the form

$$\left\{ \dots, \frac{x_m}{\alpha}, \frac{\alpha}{x_m}, \frac{\alpha\beta}{x_m}, x_m, \frac{1}{x_m}, \frac{\alpha}{x_m}, \beta x_m, \frac{1}{\beta x_m}, \frac{1}{x_m}, \alpha\beta x_m, \dots \right\}. \tag{3.46}$$

Thus the subsequence $\{x_{m+3i}\}_{i=0}^{\infty}$ is increasing and so

$$\lim_{i \rightarrow \infty} x_{m+3i} \geq \frac{1}{\sqrt{\alpha}}. \tag{3.47}$$

(B₂) $x_{m+3} = \beta/x_{m+1}$. This can be treated similarly to the case $x_{m+3} = x_m/\alpha$ and the solution is either periodic with period four or $\lim_{i \rightarrow \infty} x_{m+3i} \geq 1/\sqrt{\alpha}$.

The proof is complete. □

Remark 3.4. Observe by assumption that $x_m, x_{m+1} < 1/\sqrt{\alpha}$ is not possible as can be seen from (1.1).

Now, we can state the main result in this section.

THEOREM 3.5. *Every solution of (1.1) is periodic with period four.*

Proof. The proof of this theorem follows from Theorem 3.2 and Lemma 3.3. □

References

- [1] A. M. Amleh, J. Hoag, and G. Ladas, *A difference equation with eventually periodic solutions*, Comput. Math. Appl. **36** (1998), no. 10-12, 401–404.
- [2] W. J. Briden, E. A. Grove, G. Ladas, and C. M. Kent, *Eventually periodic solutions of $x_{n+1} = \max\{1/x_n, A_n/x_{n-1}\}$* , Comm. Appl. Nonlinear Anal. **6** (1999), no. 4, 31–43.
- [3] W. J. Briden, E. A. Grove, G. Ladas, and L. C. McGrath, *On the nonautonomous equation $x_{n+1} = \max\{A_n/x_n, B_n/x_{n-1}\}$* , New Developments in Difference Equations and Applications (Taipei, 1997), Gordon and Breach, Amsterdam, 1999, pp. 49–73.
- [4] E. M. Elabbasy, H. El-Metwally, and E. M. Elsayed, *On the periodicity behavior of some difference equations*, to appear.
- [5] H. El-Metwally, E. A. Grove, and G. Ladas, *A global convergence result with applications to periodic solutions*, J. Math. Anal. Appl. **245** (2000), no. 1, 161–170.
- [6] H. El-Metwally, E. A. Grove, G. Ladas, and H. D. Voulov, *On the global attractivity and the periodic character of some difference equations*, J. Differ. Equations Appl. **7** (2001), no. 6, 837–850.
- [7] V. L. Kocić and G. Ladas, *Global Behavior of Nonlinear Difference Equations of Higher Order With Applications*, Mathematics and Its Applications, vol. 256, Kluwer Academic, Dordrecht, 1993.
- [8] S. Stević, *A generalization of the Copson's theorem concerning sequences which satisfy a linear inequality*, Indian J. Math. **43** (2001), no. 3, 277–282.

E. M. Elabbasy: Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt

E-mail address: emelabbasy@mans.edu.eg

H. El-Metwally: Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt

E-mail address: helmetwally@mans.edu.eg

E. M. Elsayed: Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt

E-mail address: emelsayed@mans.edu.eg

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

| | |
|------------------------|------------------|
| Manuscript Due | December 1, 2008 |
| First Round of Reviews | March 1, 2009 |
| Publication Date | June 1, 2009 |

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru