

THE EXPECTED VARIATION OF RANDOM BOUNDED INTEGER SEQUENCES OF FINITE LENGTH


RUDOLFO ANGELES, DON RAWLINGS,
LAWRENCE SZE, AND MARK TIEFENBRUCK

Received 24 November 2004 and in revised form 15 June 2005

From the enumerative generating function of an abstract adjacency statistic, we deduce the mean and variance of the variation on random permutations, rearrangements, compositions, and bounded integer sequences of finite length.

1. Introduction

When the finite sequence of integers $w = 1, 3, 2, 2, 4, 3$ is sketched as below,



its most compelling aspect is its vertical *variation*, that is, the sum of the vertical distances between its adjacent terms. Denoted by $\text{var } w$, the vertical variation of the sequence in (1.1) is $\text{var } w = 2 + 1 + 0 + 2 + 1 = 6$. Our purpose here is to compute the mean and variance of var on four classical sets of combinatorial sequences.

To formalize matters and place our problem in the context of other work, let $[m]^n$ denote the set of sequences $w = x_1 x_2 \cdots x_n$ of length n with each $x_i \in \{1, 2, \dots, m\}$. For a real-valued function f on $[m]^2$, the f -adjacency number of $w = x_1 x_2 \cdots x_n \in [m]^n$ is defined to be

$$\text{adf } w = \sum_{k=1}^{n-1} f(x_k x_{k+1}). \quad (1.2)$$

Table 1.1

Sequences	Expected value of var	Variance of var
S_n	$\frac{n^2 - 1}{n}$	$\frac{(n-2)(n+1)(4n-7)}{90}$
$[m]^n$	$\frac{(n-1)(m^2 - 1)}{3m}$	$\frac{(m^2 - 1)(6m^2 n + 6n - 7m^2 - 2)}{90m^2}$
$R_n(\vec{i})$	$\frac{2}{n} \sum_{1 \leq x < y \leq m} (y - x) i_x i_y$	See (3.10)
$C_n(m)$	$\frac{2(n-1)}{(m-1)^{n-1}} \sum_{x=1}^{\lfloor m/2 \rfloor} (m-2x)^{\frac{n-1}{2}}$	See (3.18)

Some specializations of the f -adjacency number have been considered elsewhere. For instance, if $f(xy)$ is 1 when $x < y$ and 0 otherwise, then $\text{adf } w$ is known as the *rise number* of w [1, 3, 4]. For the selection $f(xy) = |y - x|$, $\text{adf } w = \text{var } w$. In a sorting problem of computer science, Levcopoulos and Petersson [5] introduced the related notion of oscillation ($\text{var } w - n + 1$) as a measure of the presortedness of a sequence of n distinct numbers. In [6], compositions were enumerated by their *ascent variation*, the f -adjacency statistic induced by $f(xy) = y - x$ if $x < y$ and 0 otherwise. For the case $f(xy) = h(|y - x|)$ where h is a linear, convex, or concave increasing real-valued function, Chao and Liang [2] described the arrangements of n distinct integers for which adf achieves its extreme values.

Besides considering the distribution of var on the set $[m]^n$, we also consider it on the set of rearrangements $R_n(i_1, i_2, \dots, i_m)$ consisting of sequences of length $n = i_1 + i_2 + \dots + i_m$ which contain l exactly i_l times, on the set of permutations $S_n = R_n(1, 1, \dots, 1)$ of $\{1, 2, \dots, n\}$, and on the set of compositions of m into n parts $C_n(m) = \{x_1 x_2 \dots x_n \in [m]^n : x_1 + x_2 + \dots + x_n = m\}$. For $m, n \geq 2$, Table 1.1 displays the mean and variance of var on these four sets. The k th falling factorial of n is $n^k = n(n-1) \dots (n-k+1)$, $\vec{i} = (i_1, i_2, \dots, i_m)$, and, for r a real number, $\lfloor r \rfloor$ denotes the greatest integer less than or equal to r . The results in Table 1.1 are new. David and Barton [3, Chapter 10] present the distributions of several statistics (some f -adjacency numbers, some not) primarily on permutations. We also note that Tiefenbruck [7] derived a generating function for compositions with bounded parts by a close relative of var . We leave open questions concerning the asymptotic behavior of var .

2. Enumerative factorial moments for f -adjacencies

Before working specifically with var , we discuss the enumerative generating function for adf on sequences as developed by Féodou and Rawlings [4]. Let $[m]^*$ denote the set of sequences of $1, 2, \dots, m$ of finite length (including the empty sequence of length 0). For $w = x_1 x_2 \dots x_n \in [m]^*$, we define $Z^w = z_{x_1} z_{x_2} \dots z_{x_n}$. The enumerative generating function for adf over $[m]^*$ is then defined to be $G(p) = \sum_{w \in [m]^*} p^{\text{adf } w} Z^w$.

By manipulating $G(p)$, we will obtain all of the information in Table 1.1 (and more). As a brief outline of our approach, note that the coefficient of $p^k z_1^{i_1} z_2^{i_2} \dots z_m^{i_m}$ in $G(p)$ is

just the number of rearrangements w in $R_n(\vec{i})$ with $\text{adf } w = k$. Thus, by dividing the coefficient of $z_1^{i_1} z_2^{i_2} \cdots z_m^{i_m}$ in $G'(1)$ by the cardinality of $R_n(\vec{i})$, we will obtain the mean of adf . So, in general, we compute the d th *enumerative factorial moment* $G^{(d)}(1) = \sum_{w \in [m]^*} (\text{adf } w)^d Z^w$.

From the work of Féodou and Rawlings [4], it follows that

$$G(p) = \frac{1}{D(p)}, \quad (2.1)$$

where

$$D(p) = 1 - \sum_{n \geq 1} \sum_{x_1 \cdots x_n \in [m]^n} Z^{x_1 \cdots x_n} \prod_{k=1}^{n-1} (p^{f(x_k x_{k+1})} - 1). \quad (2.2)$$

Examples are presented in [4, 6] for which D has a closed form. We do not know a closed form for D when $\text{adf} = \text{var}$ (that is, when $f(x, y) = |y - x|$). Nevertheless, (2.1) is still useful in computing the mean and variance of var .

Although the formula for taking the d -fold derivative with respect to p of a function of the form in (2.1) is known, we provide a short derivation. To avoid the quotient and chain rules, rewrite (2.1) as $GD = 1$. Differentiating the latter d times, $d \geq 1$, and dividing by $d!$ gives

$$\sum_{j=0}^d \frac{G^{(d-j)}}{(d-j)!} \frac{D^{(j)}}{j!} = 0. \quad (2.3)$$

To solve for $G^{(d)}$, consider the system

$$\begin{aligned} \frac{G^{(d)}}{d!} \frac{D^{(0)}}{0!} + \frac{G^{(d-1)}}{(d-1)!} \frac{D^{(1)}}{1!} + \frac{G^{(d-2)}}{(d-2)!} \frac{D^{(2)}}{2!} + \cdots + \frac{G^{(0)}}{0!} \frac{D^{(d)}}{d!} &= 0, \\ \frac{G^{(d-1)}}{(d-1)!} \frac{D^{(0)}}{0!} + \frac{G^{(d-2)}}{(d-2)!} \frac{D^{(1)}}{1!} + \cdots + \frac{G^{(0)}}{0!} \frac{D^{(d-1)}}{(d-1)!} &= 0, \\ &\vdots \\ \frac{G^{(1)}}{1!} \frac{D^{(0)}}{0!} + \frac{G^{(0)}}{0!} \frac{D^{(1)}}{1!} &= 0, \\ \frac{G^{(0)}}{0!} \frac{D^{(0)}}{0!} &= 1, \end{aligned} \quad (2.4)$$

where the top d equations arise from repeated application of (2.3). Cramer's rule applied

to the above system yields

$$G^{(d)} = \frac{(-1)^d}{d!} \begin{vmatrix} \frac{D^{(1)}}{1!} & \frac{D^{(2)}}{2!} & \frac{D^{(3)}}{3!} & \cdots & \frac{D^{(d)}}{d!} \\ \frac{D^{(0)}}{0!} & \frac{D^{(1)}}{1!} & \frac{D^{(2)}}{2!} & & \frac{D^{(d-1)}}{(d-1)!} \\ 0 & \frac{D^{(0)}}{0!} & \frac{D^{(1)}}{1!} & & \frac{D^{(d-2)}}{(d-2)!} \\ \vdots & & & & \vdots \\ 0 & & \cdots & 0 & \frac{D^{(0)}}{0!} \quad \frac{D^{(1)}}{1!} \end{vmatrix} \quad (2.5)$$

which, when expanded, implies that

$$G^{(d)} = \sum_{\nu=1}^d \frac{(-1)^\nu}{D^{\nu+1}} \sum_{\substack{j_1 + \cdots + j_\nu = d \\ j_k \geq 1}} \binom{d}{j_1 \cdots j_\nu} D^{(j_1)} \cdots D^{(j_\nu)}. \quad (2.6)$$

To determine the enumerative factorial moment $G^{(d)}(1)$, we see from (2.2) that

$$D^{(j)}(1) = - \sum_{r=2}^{j+1} D_r^{(j)}, \quad (2.7)$$

where

$$D_r^{(j)} = \sum_{x_1 \cdots x_r \in [m]^r} Z^{x_1 \cdots x_r} \sum_{\substack{l_1 + \cdots + l_{r-1} = j \\ l_k \geq 1}} \binom{j}{l_1 \cdots l_{r-1}} \prod_{k=1}^{r-1} f(x_k x_{k+1})^{\frac{l_k}{r}}. \quad (2.8)$$

For instance,

$$\begin{aligned} D'_2 &= \sum_{xy \in [m]^2} f(xy) z_x z_y, & D''_2 &= \sum_{xy \in [m]^2} f(xy)^2 z_x z_y, \\ D''_3 &= 2 \sum_{vxy \in [m]^3} f(vx) f(xy) z_v z_x z_y. \end{aligned} \quad (2.9)$$

Further setting $\vec{j} = (j_1, \dots, j_\nu)$, $s(\vec{j}) = j_1 + \cdots + j_\nu$,

$$\binom{d}{\vec{j}} = \binom{d}{j_1 \cdots j_\nu}, \quad \text{and} \quad D_\mu^{(\vec{j})} = \sum_{\substack{r_1 + \cdots + r_\nu = \mu \\ r_k \geq 2}} D_{r_1}^{(j_1)} \cdots D_{r_\nu}^{(j_\nu)}, \quad (2.10)$$

it follows from (2.6) and (2.7) that

$$G^{(d)}(1) = \sum_{\nu=1}^d \frac{1}{D^{\nu+1}(1)} \sum_{\substack{s(\vec{j}) = d \\ j_k \geq 1}} \binom{d}{\vec{j}} \sum_{\mu=2\nu}^{d+\nu} D_\mu^{(\vec{j})}. \quad (2.11)$$

As $D(1) = 1 - (z_1 + \dots + z_m)$, extracting the contributions made by all $w \in [m]^n$ from both sides of (2.11) gives the d th enumerative factorial moment of adf over $[m]^n$ as

$$\sum_{w \in [m]^n} (\text{adf } w)^d Z^w = \sum_{\nu=1}^d \sum_{\substack{s(\vec{j})=d \\ j_k \geq 1}} \binom{d}{\vec{j}} \sum_{\mu=2\nu}^{d+\nu} \binom{n+\nu-\mu}{\nu} \left(\sum_{i=1}^m z_i \right)^{n-\mu} D_{\mu}^{(\vec{j})} \quad (2.12)$$

valid for $d \geq 1$. When $d = 1, 2$, (2.9) and (2.12) imply that

$$\sum_{w \in [m]^n} \text{adf } w Z^w = (n-1) \left(\sum_{i=1}^m z_i \right)^{n-2} \sum_{xy \in [m]^2} f(xy) z_x z_y \quad (2.13)$$

and that

$$\begin{aligned} \sum_{w \in [m]^n} (\text{adf } w)^2 Z^w &= (n-1) \left(\sum_{i=1}^m z_i \right)^{n-2} \sum_{xy \in [m]^2} (f(xy))^2 z_x z_y \\ &\quad + 2(n-2) \left(\sum_{i=1}^m z_i \right)^{n-3} \sum_{vxy \in [m]^3} f(vx) f(xy) z_v z_x z_y \\ &\quad + (n-2)(n-3) \left(\sum_{i=1}^m z_i \right)^{n-4} \left(\sum_{xy \in [m]^2} f(xy) z_x z_y \right)^2. \end{aligned} \quad (2.14)$$

3. Discussion of Table 1.1

The entries in Table 1.1 are consequences of (2.13) and (2.14) with $f(xy) = |y - x|$ and with appropriate substitutions for Z . For the mean and variance of var on the set of bounded sequences $[m]^n$, put $z_i = 1$ for $1 \leq i \leq m$. Noting that

$$\sum_{xy \in [m]^2} |y - x| = \sum_{1 \leq x < y \leq m} 2(y - x) = 2 \binom{m+1}{3}, \quad (3.1)$$

it follows from (2.13) that the mean of var on $[m]^n$ is

$$\frac{1}{m^n} \sum_{w \in [m]^n} \text{var } w = \frac{2(n-1)m^{n-2}}{m^n} \binom{m+1}{3} = \frac{(n-1)(m^2-1)}{3m}. \quad (3.2)$$

As

$$\sum_{xy \in [m]^2} |y - x|^2 = 4 \binom{m+1}{4} \quad (3.3)$$

and as

$$\begin{aligned}
\sum_{vxy \in [m]^n} |x - v| |y - x| &= \sum_{1 \leq v < x < y \leq m} 2(x - v)(y - x) \\
&\quad + \sum_{1 \leq x < y \leq v \leq m} 4(v - x)(y - x) - \sum_{1 \leq x < y \leq m} 2(y - x)^2 \\
&= \frac{7m^2 - 8}{10} \binom{m+1}{3},
\end{aligned} \tag{3.4}$$

(2.14) implies that

$$\begin{aligned}
\frac{1}{m^n} \sum_{w \in [m]^n} (\text{var } w)^2 &= \frac{4(n-1)}{m^2} \binom{m+1}{4} + \frac{(n-2)(7m^2-8)}{5m^3} \binom{m+1}{3} \\
&\quad + \frac{4(n-2)(n-3)}{m^4} \binom{m+1}{3}^2.
\end{aligned} \tag{3.5}$$

Then, subbing the last result into

$$\frac{1}{m^n} \sum_{w \in [m]^n} (\text{var } w)^2 + \frac{(n-1)(m^2-1)}{3m} - \left(\frac{(n-1)(m^2-1)}{3m} \right)^2 \tag{3.6}$$

and simplifying gives the variance of $\text{var } w$ as recorded in Table 1.1.

For $R_n(\vec{i})$, extracting the coefficient of $z_1^{i_1} z_2^{i_2} \cdots z_m^{i_m}$ from (2.13) leads to

$$\sum_{w \in R_n(\vec{i})} \text{var } w = 2(n-1) \sum_{1 \leq x < y \leq m} (y-x) \binom{n-2}{i_1 \cdots i_x - 1 \cdots i_y - 1 \cdots i_m}. \tag{3.7}$$

As the cardinality of $R_n(\vec{i})$ is

$$\binom{n}{i_1 i_2 \cdots i_m} = \binom{n}{\vec{i}}, \tag{3.8}$$

it follows that the mean of $\text{var } w$ on $R_n(\vec{i})$ is

$$\binom{n}{\vec{i}}^{-1} \sum_{w \in R_n(\vec{i})} \text{var } w = \frac{2}{n} \sum_{1 \leq x < y \leq m} (y-x) i_x i_y. \tag{3.9}$$

Let $\vec{i}_{\setminus r} = (i_1, \dots, i_r - 1, \dots, i_n)$. For example, $(3, 2, 1, 4)_{\setminus 3 \setminus 2 \setminus 3} = (3, 1, -1, 4)$. The variance on $R_n(\vec{i})$ is then

$$\binom{n}{\vec{i}}^{-1} \sum_{w \in R_n(\vec{i})} \text{var } w^2 + \frac{2}{n} \sum_{1 \leq x < y \leq m} (y-x) i_x i_y - \left(\frac{2}{n} \sum_{1 \leq x < y \leq m} (y-x) i_x i_y \right)^2, \tag{3.10}$$

where, upon extraction of the coefficient of $z_1^{i_1} z_2^{i_2} \dots z_m^{i_m}$ from (2.14), we have

$$\begin{aligned} \sum_{w \in R_n(\vec{i})} (\text{var } w)^2 &= (n-1) \sum_{1 \leq x, y \leq m} |y-x|^2 \binom{n-2}{\vec{i}_{\setminus x \setminus y}} \\ &+ 2(n-2) \sum_{1 \leq v, x, y \leq m} |x-v| |y-x| \binom{n-3}{\vec{i}_{\setminus v \setminus x \setminus y}} \\ &+ (n-2)(n-3) \sum_{1 \leq u, v, x, y \leq m} |v-u| |y-x| \binom{n-4}{\vec{i}_{\setminus u \setminus v \setminus x \setminus y}}. \end{aligned} \quad (3.11)$$

The permutation entries in Table 1.1 follow from (3.9) and (3.10). Selecting $m = n$ and $i_k = 1$ for $1 \leq k \leq n$ in (3.9) reveals the mean of var on S_n as

$$\frac{1}{n!} \sum_{w \in S_n} \text{var } w = \frac{2}{n} \sum_{1 \leq x < y \leq n} (y-x) = \frac{2}{n} \binom{n+1}{3} = \frac{n^2-1}{3}. \quad (3.12)$$

From (3.11), with $m = n$ and $i_k = 1$ for $1 \leq k \leq n$,

$$\begin{aligned} \sum_{w \in S_n} (\text{var } w)^2 &= (n-1)! \sum_{1 \leq x, y \leq n} |y-x|^2 \\ &+ 2(n-2)! \sum_{1 \leq v, x, y \leq n} |x-v| |y-x| \\ &+ (n-2)! \sum_{\substack{1 \leq u, v, x, y \leq m \\ \{u, v\} \cap \{x, y\} = \emptyset}} |v-u| |y-x| \\ &= \left(\frac{4}{15} \right) (n-2)! (10n^2 + 14n - 27) \binom{n+1}{4}. \end{aligned} \quad (3.13)$$

So the variance of var on S_n is

$$\frac{1}{n!} \sum_{w \in S_n} \text{var } w^2 + \frac{n^2-1}{3} - \left(\frac{n^2-1}{3} \right)^2 = \frac{(n-2)(n+1)(4n-7)}{90}. \quad (3.14)$$

For $w = x_1 \dots x_n \in [m]^n$, let $\|w\| = x_1 + \dots + x_n$. For the composition results in Table 1.1, set $z_k = q^k$ for $1 \leq k \leq m$. Then (2.13) implies that

$$\sum_{w \in [m]^n} \text{var } w q^{\|w\|} = (n-1)q^{n-2} \left(\frac{1-q^m}{1-q} \right)^{n-2} \sum_{1 \leq x, y \leq m} |y-x| q^{x+y} \quad (3.15)$$

and (2.14) leads to

$$\begin{aligned}
\sum_{w \in [m]^n} (\text{var } w)^2 q^{\|w\|} &= (n-1)q^{n-2} \left(\frac{1-q^m}{1-q} \right)^{n-2} \sum_{1 \leq x, y \leq m} |y-x|^2 q^{x+y} \\
&\quad + 2(n-2)q^{n-3} \left(\frac{1-q^m}{1-q} \right)^{n-3} \sum_{1 \leq v, x, y \leq m} |x-v| |y-x| q^{v+x+y} \\
&\quad + (n-2)(n-3)q^{n-4} \left(\frac{1-q^m}{1-q} \right)^{n-4} \sum_{1 \leq u, v, x, y \leq m} |v-u| |y-x| q^{u+v+x+y}.
\end{aligned} \tag{3.16}$$

Extracting the coefficient of q^m from (3.15) to obtain

$$\begin{aligned}
\sum_{w \in C_n(m)} \text{var } w &= 2(n-1) \sum_{1 \leq x < y \leq m} (y-x) \binom{m-1-x-y}{n-3} \\
&= 2(n-1) \sum_{1 \leq x \leq \lfloor m/2 \rfloor} \binom{m-2x}{n-1}
\end{aligned} \tag{3.17}$$

and then dividing by the cardinality $\binom{m-1}{n-1}$ of $C_n(m)$ gives the mean of var as stated in Table 1.1. The variance is

$$\begin{aligned}
&\binom{m-1}{n-1}^{-1} \sum_{w \in C_n(m)} \text{var } w^2 + \frac{2(n-1)}{(m-1)^{n-1}} \sum_{1 \leq x \leq \lfloor m/2 \rfloor} (m-2x)^{n-1} \\
&\quad - \left(\frac{2(n-1)}{(m-1)^{n-1}} \sum_{1 \leq x \leq \lfloor m/2 \rfloor} (m-2x)^{n-1} \right)^2,
\end{aligned} \tag{3.18}$$

where, pulling the coefficient of q^m from (3.16), we have

$$\begin{aligned}
\sum_{w \in C_n(m)} (\text{var } w)^2 &= (n-1) \sum_{1 \leq x, y \leq m} |y-x|^2 \binom{m-1-x-y}{n-3} \\
&\quad + 2(n-2) \sum_{1 \leq v, x, y \leq m} |x-v| |y-x| \binom{m-1-v-x-y}{n-4} \\
&\quad + (n-2)(n-3) \sum_{1 \leq u, v, x, y \leq m} |v-u| |y-x| \binom{m-1-u-v-x-y}{n-5}.
\end{aligned} \tag{3.19}$$

The sums in (3.19) are marginally simplified. For instance,

$$\sum_{1 \leq x, y \leq m} |y-x|^2 \binom{m-1-x-y}{n-3} = 4 \sum_{1 \leq x \leq \lfloor m/2 \rfloor} \binom{m-2x}{n}. \tag{3.20}$$

As a part of the second sum on the right-hand side of (3.19), we note that

$$\begin{aligned} & \sum_{1 \leq v < x < y \leq m} (x - v)(y - x) \binom{m - 1 - v - x - y}{n - 4} \\ &= \sum_{2 \leq x \leq \lfloor (m+1)/2 \rfloor} \left(\binom{m - 3x + 1}{n} - \binom{m - 2x + 1}{n} + x \binom{m - 2x}{n - 1} \right). \end{aligned} \quad (3.21)$$

The four-fold sums arising in the last sum in (3.19) reduce to double sums.

Acknowledgment

This work is based on work supported by the National Science Foundation under Grant no. 0097392.

References

- [1] L. Carlitz, *Enumeration of compositions by rises, falls and levels*, Math. Nachr. **77** (1977), 361–371.
- [2] C.-C. Chao and W.-Q. Liang, *Arranging n distinct numbers on a line or a circle to reach extreme total variations*, European J. Combin. **13** (1992), no. 5, 325–334.
- [3] F. N. David and D. E. Barton, *Combinatorial Chance*, Hafner, New York, 1962.
- [4] J.-M. Fédou and D. P. Rawlings, *Adjacencies in words*, Adv. in Appl. Math. **16** (1995), no. 2, 206–218.
- [5] C. Levcopoulos and O. Petersson, *Heapsort—adapted for presorted files*, Algorithms and Data Structures (Ottawa, ON, 1989), Lecture Notes in Comput. Sci., vol. 382, Springer, Berlin, 1989, pp. 499–509.
- [6] D. P. Rawlings, *Restricted words by adjacencies*, Discrete Math. **220** (2000), no. 1-3, 183–200.
- [7] M. Tiefenbruck, *Enumerating compositions with bounded parts by variation*, REU report, California Polytechnic State University, California, 2003.

Rudolfo Angeles: Department of Mathematics, College of Science and Mathematics, California Polytechnic State University, San Luis Obispo, CA 93407, USA

Current address: Department of Statistics, Stanford University, Palo Alto, CA 94305-4065, USA

E-mail address: rangeles@stanford.edu

Don Rawlings: Department of Mathematics, College of Science and Mathematics, California Polytechnic State University, San Luis Obispo, CA 93407, USA

E-mail address: drawling@calpoly.edu

Lawrence Sze: Department of Mathematics, College of Science and Mathematics, California Polytechnic State University, San Luis Obispo, CA 93407, USA

E-mail address: lsze@calpoly.edu

Mark Tiefenbruck: Department of Mathematics, College of Science and Mathematics, California Polytechnic State University, San Luis Obispo, CA 93407, USA

Current address: 8989 Jasmine Lane Street, Cottage Grove, MN 55016-3436, USA

E-mail address: mdash@alumni.northwestern.edu

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk