

ASYMMETRIC LOCALLY CONVEX SPACES

S. COBZAŞ

Received 14 February 2005 and in revised form 6 June 2005

The aim of the present paper is to introduce the asymmetric locally convex spaces and to prove some basic properties. Among these I do mention the analogs of the Eidelheit-Tuckey separation theorems, of the Alaoglu-Bourbaki theorem on the weak compactness of the polar of a neighborhood of 0, and a Krein-Milman-type theorem. These results extend those obtained by García-Raffi et al. (2003) and Cobzaş (2004).

1. Introduction

Let X be a real vector space. An *asymmetric seminorm* on X is a positive sublinear functional $p : X \rightarrow [0; \infty)$, that is, p satisfies the conditions

- (AN1) $p(x) \geq 0$,
- (AN2) $p(tx) = tp(x)$, $t \geq 0$,
- (AN3) $p(x+y) \leq p(x) + p(y)$,

for all $x, y \in X$. The function $\bar{p} : X \rightarrow [0, \infty)$, defined by $\bar{p}(x) = p(-x)$, $x \in X$, is another positive sublinear functional on X , called the *conjugate* of p , and

$$p^s(x) = \max \{p(x), p(-x)\}, \quad x \in X, \quad (1.1)$$

is a seminorm on X . The inequalities

$$|p(x) - p(y)| \leq p^s(x - y), \quad |\bar{p}(x) - \bar{p}(y)| \leq p^s(x - y) \quad (1.2)$$

hold for all $x, y \in X$. If the seminorm p^s is a norm on X , then we say that p is an *asymmetric norm* on X . This means that, beside (AN1)–(AN3), it satisfies also the condition

- (AN4) $p(x) = 0$ and $p(-x) = 0$ imply that $x = 0$.

The pair (X, p) , where X is a linear space and p is an asymmetric seminorm on X is called a *space with asymmetric seminorm*, respectively, a *space with asymmetric norm*, if p is an asymmetric norm.

In the last years, the properties of spaces with asymmetric norms were investigated in a series of papers, emphasizing similarities as well as differences with respect to the theory of (symmetric) normed spaces, see [3, 5, 6, 7, 12, 13, 16, 17]. This study was stimulated

also by their applications in the complexity of algorithms, see [11, 14, 18]. The aim of the present paper is to develop the basic results in the theory of asymmetric locally convex spaces, a natural extension of asymmetric normed spaces and of (symmetric) locally convex spaces as well.

The function $\rho : X \times X \rightarrow [0; \infty]$ defined by $\rho(x, y) = p(y - x)$, $x, y \in X$, is an asymmetric semimetric on X . Denote by

$$B'_p(x, r) = \{x' \in X : p(x' - x) < r\}, \quad B_p(x, r) = \{x' \in X : p(x' - x) \leq r\}, \quad (1.3)$$

the open, respectively closed, ball in X of center x and radius $r > 0$. Denoting by

$$B'_p = B'_p(0, 1), \quad B_p = B_p(0, 1) \quad (1.4)$$

the corresponding unit balls, then

$$B'_p(x, r) = x + rB'_p, \quad B_p(x, r) = x + rB_p. \quad (1.5)$$

The unit balls B'_p and B_p are convex absorbing subsets of the space X and p agrees with the Minkowski functional associated to any of them.

An asymmetric seminorm p on X generates a topology τ_p on X , having as basis of neighborhoods of a point $x \in X$ the family $\{B'_p(x, r) : r > 0\}$ of open p -balls. The family $\{B_p(x, r) : r > 0\}$ of closed p -balls is also a neighborhood basis at x for τ_p .

The ball $B'_p(x, r)$ is τ_p -open but the ball $B_p(x, r)$ need not to be τ_p -closed, as can be seen from the following typical example.

Example 1.1. Consider on \mathbb{R} the asymmetric seminorm $u(\alpha) = \max\{\alpha, 0\}$, $\alpha \in \mathbb{R}$, and denote by \mathbb{R}_u the space \mathbb{R} equipped with the topology τ_u generated by u . The conjugate seminorm is $\bar{u}(\alpha) = -\min\{\alpha, 0\}$, and $u^s(\alpha) = \max\{u(\alpha), \bar{u}(\alpha)\} = |\alpha|$. The topology τ_u , called the *upper topology* of \mathbb{R} , is generated by the intervals of the form $(-\infty; a)$, $a \in \mathbb{R}$, and the family $\{(-\infty; \alpha + \epsilon) : \epsilon > 0\}$ is a neighborhood basis of a point $\alpha \in \mathbb{R}$. The set $(-\infty; 1) = B'_u(0, 1)$ is τ_u -open, and the ball $B_u(0, 1) = (-\infty; 1]$ is not τ_u -closed because $\mathbb{R} \setminus B_u(0, 1) = (1; \infty)$ is not τ_u -open.

Remark 1.2. As can be easily seen, the continuity of a mapping f from a topological space (T, τ) to (\mathbb{R}, τ_u) is equivalent to its upper semicontinuity as a mapping from (T, τ) to $(\mathbb{R}, |\cdot|)$.

The topology τ_p is translation invariant, that is, the addition $+ : X \times X \rightarrow X$ is continuous, but the multiplication by scalars $\cdot : \mathbb{R} \times X \rightarrow X$ need not be continuous, as it is shown by some examples, as, for example, that given in [5]. We will present another one in the context of Example 1.1.

Example 1.3. In the space (\mathbb{R}, u) from Example 1.1, the interval $(-\infty; 1/2)$ is a τ_u -neighborhood of $0 = (-1)0$ but for any $\alpha, \beta > 0$, the neighborhood $(-\infty; -1 + \alpha) \times (-\infty, \beta)$ of $(-1, 0)$ contains the point $(-1, -1)$ and $(-1)(-1) = 1 \notin (-\infty; 1/2)$.

The discontinuity of the multiplication by scalars, $(\alpha, x) \mapsto \alpha x$, for $\alpha = -1$ follows also from the fact that the interval $(-\infty; a)$ is τ_u -open but $-(-\infty; a) = (-a; \infty)$ is not τ_u -open.

The following proposition will be useful in the study of the continuity of linear mappings between asymmetric locally convex spaces.

PROPOSITION 1.4. *If X is a real vector space, $f, g : X \rightarrow \mathbb{R}$ are sublinear functionals, and $\alpha, \beta > 0$, then the following conditions are equivalent:*

$$\forall x \in X, \quad g(x) \leq \beta \implies f(x) \leq \alpha, \quad (1.6)$$

$$\forall x \in X, \quad u(f(x)) \leq \frac{\alpha}{\beta} u(g(x)). \quad (1.7)$$

If $g(x) \geq 0$ for all $x \in X$, then these two conditions are also equivalent to the following one:

$$\forall x \in X, \quad f(x) \leq \frac{\alpha}{\beta} g(x). \quad (1.8)$$

Proof. (1.6) \Rightarrow (1.7). Let $x \in X$. If $g(x) \leq 0$, then $g(nx) = ng(x) \leq 0 < \alpha$, $n \in \mathbb{N}$, so that $nf(x) = f(nx) \leq \beta$, $n \in \mathbb{N}$, implying that $f(x) \leq 0$ and

$$u(f(x)) = 0 = \frac{\alpha}{\beta} u(g(x)). \quad (1.9)$$

If $g(x) > 0$, then $g((\beta/g(x))x) = \beta$, so that

$$f\left(\frac{\beta}{g(x)}x\right) \leq \alpha \iff f(x) \leq \frac{\alpha}{\beta} g(x) \iff u(f(x)) \leq \frac{\alpha}{\beta} u(g(x)). \quad (1.10)$$

(1.7) \Rightarrow (1.6). Let $x \in X$. If $g(x) \leq 0 < \beta$, then $u(g(x)) = 0$, so that

$$f(x) \leq u(f(x)) \leq \frac{\alpha}{\beta} u(g(x)) = \frac{\alpha}{\beta} g(x) \leq \alpha. \quad (1.11)$$

If $g(x) > 0$, then by hypothesis,

$$f(x) \leq u(f(x)) \leq \frac{\alpha}{\beta} u(g(x)) = \frac{\alpha}{\beta} g(x) \leq \alpha. \quad (1.12)$$

Since $g(x) \geq 0$, $x \in X$, implies that $u(g(x)) = g(x)$, $x \in X$, the equivalence (1.7) \Leftrightarrow (1.8) is obvious. \square

Let now P be a family of asymmetric seminorms on a real vector space X . Denote by $\mathcal{F}(P)$ the family of all nonempty finite subsets of P , and for $F \in \mathcal{F}(P)$, $x \in X$, and $r > 0$, let

$$\begin{aligned} B_F(x, r) &= \{y \in X : p(y - x) \leq r, p \in F\} = \bigcap \{B_p(x, r) : p \in F\}, \\ B'_F(x, r) &= \{y \in X : p(y - x) < r, p \in F\} = \bigcap \{B'_p(x, r) : p \in F\} \end{aligned} \quad (1.13)$$

denote the closed, respectively, open multiball of center x and radius r . It is immediate that these multiballs are convex absorbing subsets of X .

Putting

$$p_F(x) = \max \{p(x) : p \in F\}, \quad x \in X, \quad (1.14)$$

then p_F is an asymmetric seminorm on X and

$$B_F(x, r) = B_{p_F}(x, r), \quad B'_F(x, r) = B'_{p_F}(x, r). \quad (1.15)$$

The *asymmetric locally convex* topology associated to the family P of asymmetric seminorms on a real vector space X is the topology τ_P having as basis of neighborhoods of any point $x \in X$ the family $\mathcal{N}'(x) = \{B'_F(x, r) : r > 0, F \in \mathcal{F}(P)\}$ of open multiballs. The family $\mathcal{N}(x) = \{B_F(x, r) : r > 0, F \in \mathcal{F}(P)\}$ of closed multiballs is also a neighborhood basis at x for τ_P .

It is easy to check that the family $\mathcal{N}'(x)$ fulfills the requirements of a neighborhood basis, that is,

- (BN1) $x \in B'_F(x, r)$,
- (BN2) for $B'_{F_1}(x, r_1)$ and $B'_{F_2}(x, r_2)$ in $\mathcal{N}'(x)$, there exists $B'_F(x, r) \in \mathcal{N}'(x)$ such that $B'_F(x, r) \subset B'_{F_1}(x, r_1) \cap B'_{F_2}(x, r_2)$.

For (BN2), one can take $F = F_1 \cup F_2$ and $r = \min\{r_1, r_2\}$.

Obviously, for $P = \{p\}$, we obtain the topology τ_p of an asymmetric seminormed space (X, p) considered above, that is, $\tau_{\{p\}} = \tau_p$.

The topology τ_P is derived from a quasiuniformity \mathcal{W}_P on X having as vicinities the sets

$$W_F(\epsilon) = \{(x, y) \in X \times X : p(y - x) < \epsilon, p \in F\}, \quad (1.16)$$

for $F \in \mathcal{F}(P)$ and $\epsilon > 0$. Replacing the sign $<$ by \leq in the above definition, the corresponding sets will form a basis for the same quasiuniformity \mathcal{W}_P . A good source for the properties of quasiuniform spaces is the book [10] (see also [4]). Quasiuniform structures related to asymmetric normed spaces were investigated in [1, 2, 9].

We say that the family P is *directed* if for any $p_1, p_2 \in P$, there exists $p \in P$ such that $p \geq p_i$, $i = 1, 2$, where $p \geq q$ stands for the pointwise ordering: $p(x) \geq q(x)$ for all $x \in X$. If the family P is directed, then for any τ_P -neighborhood of a point $x \in X$, there exist $p \in P$ and $r > 0$ such that $B'_p(x, r) \subset V$ (resp., $B_p(x, r) \subset V$). Indeed, if $B'_F(x, r) \subset V$, then there exists $p \in P$ such that $p \geq q$ for all $q \in F$ so that $B'_p(x, r) \subset B'_F(x, r) \subset V$. Similarly, the vicinities defined by (1.16) with $F = \{p\}$, $p \in P$, and $\epsilon > 0$ form a basis for the quasiuniformity \mathcal{W}_P .

The family

$$P_d = \{p_F : F \in \mathcal{F}(P)\}, \quad (1.17)$$

where p_F is defined by (1.14), is a directed family of asymmetric seminorms generating the same topology as P , that is, $\tau_{P_d} = \tau_P$. Therefore, without restricting the generality, we can always suppose that the family P of asymmetric seminorms is directed.

Because $B_F(x, r) = x + B_F(0, r)$ and $B'_F(x, r) = x + B'_F(0, r)$, the topology τ_p is translation invariant,

$$\mathcal{V}(x) = \{x + V : V \in \mathcal{V}(0)\}, \quad (1.18)$$

where by $\mathcal{V}(x)$ we have denoted the family of all neighborhoods with respect to τ_p of a point $x \in X$.

The addition $+ : X \times X \rightarrow X$ is continuous. Indeed, for $x, y \in X$ and the neighborhood $B'_F(x + y, r)$ of $x + y$, we have $B'_F(x, r/2) + B'_F(y, r/2) \subset B'_F(x + y, r)$.

As we have seen in Example 1.3, the multiplication by scalars need not be continuous, even in asymmetric seminormed spaces.

The topology τ_p generated by an asymmetric norm is not always Hausdorff. A necessary and sufficient condition in order that τ_p be Hausdorff is given in the following proposition.

PROPOSITION 1.5 (see [13]). *For an asymmetric seminorm p on a real vector space X , put*

$$\tilde{p}(x) = \inf \{p(x') + p(x' - x) : x' \in X\}, \quad x \in X. \quad (1.19)$$

(1) *The functional \tilde{p} is a (symmetric) seminorm on X , $\tilde{p} \leq p$, and \tilde{p} is the greatest of the seminorms on X majorized by p .*

(2) *The topology τ_p generated by p is Hausdorff if and only if $\tilde{p}(x) > 0$ for every $x \neq 0$.*

Proof. We will give a proof of the first assertion, different from that given in [13]. The second assertion will be proved in the more general context of asymmetric locally convex spaces.

First, observe that replacing x' by $x' - x$ in (1.19), we get

$$\begin{aligned} \tilde{p}(-x) &= \inf \{p(x') + p(x' + x) : x' \in X\} \\ &= \inf \{p(x' - x) + p((x' - x) + x) : x' \in X\} = \tilde{p}(x), \end{aligned} \quad (1.20)$$

so that \tilde{p} is symmetric. The positive homogeneity of \tilde{p} , $\tilde{p}(\alpha x) = \alpha \tilde{p}(x)$, $x \in X$, $\alpha \geq 0$, is obvious. For $x, y \in X$ and arbitrary $x', y' \in X$, we have

$$\tilde{p}(x + y) \leq p(x' + y') + p(x' + y' - x - y) \leq p(x') + p(x' - x) + p(y') + p(y' - y), \quad (1.21)$$

so that passing to infimum with respect to $x', y' \in X$, we obtain the subadditivity of \tilde{p} ,

$$\tilde{p}(x + y) \leq \tilde{p}(x) + \tilde{p}(y). \quad (1.22)$$

Suppose now that there exists a seminorm q on X such that $q \leq p$, that is, for all $z \in X$, $q(z) \leq p(z)$, and $\tilde{p}(x) < q(x) \leq p(x)$, for some $x \in X$. Then, by the definition of \tilde{p} , there exists $x' \in X$ such that $\tilde{p}(x) < p(x') + p(x' - x) < q(x)$, leading to the contradiction

$$q(x) \leq q(x') + q(x - x') = q(x') + q(x' - x) \leq p(x') + p(x' - x) < q(x). \quad (1.23)$$

□

The following characterization of the Hausdorff separation property for locally convex spaces is well known, see, for example, [19, Lemma VIII.1.4].

PROPOSITION 1.6. *Let (X, Q) be a locally convex space, where Q is a family of seminorms generating the topology τ_Q of X . The topology τ_Q is Hausdorff separated if and only if for every $x \in X, x \neq 0$, there exists $q \in Q$ such that $q(x) > 0$.*

In the case of asymmetric locally convex spaces, we have the following characterization.

PROPOSITION 1.7. *Let P be a family of asymmetric seminorms on a real vector space X . The asymmetric locally convex topology τ_P is Hausdorff separated if and only if for every $x \in X, x \neq 0$, there exists $p \in P$ such that $\tilde{p}(x) > 0$, where \tilde{p} is the seminorm associated to the asymmetric seminorm p through the formula (1.19).*

Proof. Suppose that P is directed and let

$$\tilde{P} = \{\tilde{p} : p \in P\}. \quad (1.24)$$

Denote by $\tau_{\tilde{P}}$ the locally convex topology on X generated by the family \tilde{P} of seminorms. The topology τ_P is finer than $\tau_{\tilde{P}}$. Indeed, $\tilde{G} \in \tau_{\tilde{P}}$ is equivalent to the fact that for every $x \in \tilde{G}$, there exist $p \in P$ and $r > 0$ such that $B'_{\tilde{p}}(x, r) \subset \tilde{G}$. Because $p(y - x) < r$ implies that $\tilde{p}(y - x) \leq p(y - x) < r$, we have $B'_p(x, r) \subset B'_{\tilde{p}}(x, r) \subset \tilde{G}$, so that $\tilde{G} \in \tau_P$. If for every $x \in X, x \neq 0$, there exists $p \in P$ such that $\tilde{p}(x) > 0$, then the locally convex topology $\tau_{\tilde{P}}$ is separated Hausdorff, and so will be the finer topology τ_P .

Conversely, suppose that the topology τ_P is Hausdorff and show that $\tilde{p}(x) = 0$ for all $p \in P$ implies that $x = 0$.

Let $x \in P$ be such that $\tilde{p}(x) = 0$ for all $p \in P$. By the definition (1.19) of the seminorm \tilde{p} , for every $p \in P$ and $n \in \mathbb{N}$, there exists an element $x_{(p,n)} \in X$ such that

$$p(x_{(p,n)}) + p(x_{(p,n)} - x) < \frac{1}{n}. \quad (1.25)$$

Define the order on $P \times \mathbb{N}$ by $(p, n) \leq (q, m)$ if and only if $p \leq q$ and $n \leq m$. Since the family P of asymmetric seminorms is directed, the set $P \times \mathbb{N}$ is also directed with respect to the order we just defined. Therefore, $\{x_{(p,n)} : (p, n) \in P \times \mathbb{N}\}$ is a net in X and by (1.25), we have

$$p(x_{(p,n)}) < \frac{1}{n}, \quad p(x_{(p,n)} - x) < \frac{1}{n}, \quad (1.26)$$

for all $(p, n) \in P \times \mathbb{N}$.

We will prove that the net $\{x_{(p,n)}\}$ converges to both 0 and x . Since the topology τ_P is Hausdorff, this will imply that $x = 0$.

To prove that the net $\{x_{(p,n)}\}$ converges to 0, we have to show that for every $p \in P$, the net $\{p(x_{(p,n)})\}$ tends to 0, that is,

$$\begin{aligned} \forall p \in P, \forall \epsilon > 0, \exists (p_0, n_0) \in P \times \mathbb{N}, \forall (q, n) \in P \times \mathbb{N}, \text{ such that} \\ (q, n) \geq (p_0, n_0) \implies p(x_{(q,n)}) < \epsilon. \end{aligned} \quad (1.27)$$

Let $p \in P$ and $\epsilon > 0$. Put $p_0 = p$ and let $n_0 \in \mathbb{N}$ be such that $1/n_0 < \epsilon$. Then for every $(q, n) \in P \times \mathbb{N}$ such that $q \geq p$ and $n \geq n_0$, we have

$$p(x_{(q,n)}) \leq q(x_{(q,n)}) < \frac{1}{n} \leq \frac{1}{n_0} < \epsilon. \quad (1.28)$$

The convergence of $\{p(x_{(p,n)} - x)\}$ to 0, which is equivalent to the τ_p -convergence of $\{x_{(p,n)}\}$ to x , can be proved similarly, using the second inequality in (1.26). \square

COROLLARY 1.8. *Let (X, P) be an asymmetric locally convex space. If the topology τ_p is Hausdorff, then for every $x \in X$, $x \neq 0$, there exists $p \in P$ such that $p(x) > 0$.*

Proof. If the topology τ_p is Hausdorff, then for every $x \in X$, $x \neq 0$, there exists $p \in P$ such that $\tilde{p}(x) > 0$. Replacing x by $-x$ and taking $x' = 0$ in the definition (1.19) of the seminorm \tilde{p} , we get

$$p(x) = p(0) + p(0 + x) \geq \tilde{p}(x) > 0. \quad (1.29)$$

\square

As in the symmetric case, asymmetric locally convex topologies can be defined through some basic families of convex absorbing sets.

A nonempty family \mathcal{C} of subsets of a real vector space X is called an *asymmetric locally convex basis* provided that

(BC1) each $C \in \mathcal{C}$ is convex and absorbing;

(BC2) for every $C_1, C_2 \in \mathcal{C}$, there exists $C \in \mathcal{C}$ such that $C \subset C_1 \cap C_2$;

(BC3) for every $C \in \mathcal{C}$ and $\alpha > 0$, there exists $D \in \mathcal{C}$ such that $D \subset \alpha C$.

Define a mapping $\mathcal{U} : X \rightarrow 2^X$ by

$$\mathcal{U}(x) = \{U \subset X : \exists C \in \mathcal{C} \text{ such that } x + C \subset U\}. \quad (1.30)$$

Recall that for an absorbing subset C of X , the Minkowski functional $p_C : X \rightarrow [0; \infty)$ is defined by

$$p_C(x) = \inf\{t > 0 : x \in tC\}. \quad (1.31)$$

If C is absorbing and convex, then p_C is a positive sublinear functional, and

$$\{x \in X : p_C(x) < 1\} \subset C \subset \{x \in X : p_C(x) \leq 1\}. \quad (1.32)$$

Conversely, if p is a positive sublinear functional on X , then $C' = \{x \in X : p(x) < 1\}$ and $C = \{x \in X : p(x) \leq 1\}$ are convex absorbing subsets of X , and $p_{C'} = p_C = p$.

Denoting by

$$P = \{p_C : C \in \mathcal{C}\} \quad (1.33)$$

the family of all Minkowski functionals associated to the sets in \mathcal{C} , then P is a family of asymmetric seminorms on X . By (BC1) and the fact that $p_C \leq p_D$ if $D \subset C$, it follows that the family P is directed.

PROPOSITION 1.9. *The family $\mathcal{U}(x)$ of subsets of X given by (1.30) satisfies the axioms of a neighborhood system, so that it defines a topology $\tau_{\mathcal{U}}$ on X . This topology agrees with the asymmetric locally convex topology generated by the family (1.33) of asymmetric seminorms.*

Proof. One can easily check that the family \mathcal{U} of subsets of X satisfies the axioms of a neighborhood system.

Since both of the topologies $\tau_{\mathcal{U}}$ and τ_P are translation invariant, in order to prove their coincidence, it suffices to show that they have the same 0-neighborhoods. Denote by \mathcal{V} the neighborhood mapping associated to τ_P . If $U \in \mathcal{U}(0)$, then there exists $C \in \mathcal{C}$ such that $C \subset U$. The inclusions

$$\{x \in X : p_C(x) < 1\} \subset C \subset U \quad (1.34)$$

show that $U \in \mathcal{V}(0)$.

Conversely, if $V \in \mathcal{V}(0)$, then there exist $C \in \mathcal{C}$ and $r > 0$ such that $\{x \in X : p_C(x) \leq r\} \subset V$. By (BC3), there exists $D \in \mathcal{C}$ such that $D \subset rC$. But then

$$D \subset rC \subset \{x \in X : p_C(x) \leq r\} \subset V, \quad (1.35)$$

so that $V \in \mathcal{U}(0)$. □

2. Bounded linear mappings between asymmetric locally convex spaces and the dual space

Let (X, P) , (Y, Q) be two asymmetric locally convex spaces with the topologies τ_P and τ_Q generated by the families P and Q of asymmetric seminorms on X and Y , respectively. In the following, when we say that (X, P) is an asymmetric locally convex space, we understand that X is a real vector space, P is a family of asymmetric seminorms on X , and τ_P is the asymmetric locally convex topology associated to P .

A linear mapping $A : X \rightarrow Y$ is called (P, Q) -bounded if for every $q \in Q$, there exist $F \in \mathcal{F}(P)$ and $L \geq 0$ such that

$$\forall x \in X, \quad q(Ax) \leq L \max \{p(x) : p \in F\}. \quad (2.1)$$

If the family P is directed, then the (P, Q) -boundedness of A is equivalent to the condition that for every $q \in Q$, there exist $p \in P$ and $L \geq 0$ such that

$$\forall x \in X, \quad q(Ax) \leq L p(x). \quad (2.2)$$

The continuity of the mapping A from (X, τ_P) to (Y, τ_Q) is called (τ_P, τ_Q) -continuity. We will use also the term (P, Q) -continuity for this property, and (P, u) -continuity in the case of (τ_P, τ_u) -continuous linear functionals.

Because both of the topologies τ_P and τ_Q are translation invariant, we have the following result. Recall that a mapping F between two quasiuniform spaces (X, \mathcal{U}) and (Y, \mathcal{W}) is called quasiuniformly continuous if for every $W \in \mathcal{W}$, there exists $U \in \mathcal{U}$ such that $(F(x), F(y)) \in W$ for every $(x, y) \in U$.

PROPOSITION 2.1. *Let (X, P) and (Y, Q) be asymmetric locally convex spaces and $A : X \rightarrow Y$ a linear mapping. The following conditions are equivalent.*

- (1) *The mapping A is (P, Q) -continuous on X .*
- (2) *The mapping A is continuous at $0 \in X$.*
- (3) *The mapping A is continuous at some point $x_0 \in X$.*

The following proposition emphasizes the equivalence of continuity and boundedness for linear mappings.

PROPOSITION 2.2. *Let (X, P) and (Y, Q) be two asymmetric locally convex spaces and $A : X \rightarrow Y$ a linear mapping. The following assertions are equivalent.*

- (1) *The mapping A is (P, Q) -continuous on X .*
- (2) *The mapping A is continuous at $0 \in X$.*
- (3) *The mapping A is (P, Q) -bounded.*
- (4) *The mapping A is quasiuniformly continuous with respect to the quasiuniformities \mathcal{W}_P and \mathcal{W}_Q .*

Proof. The equivalence (1) \Leftrightarrow (2) follows from the preceding proposition.

Suppose that the families P and Q are directed.

(2) \Leftrightarrow (3). For $q \in Q$, consider the τ_Q -neighborhood $V = B_q(0, 1)$ of $A0 = 0 \in Y$, and let U be a neighborhood of $0 \in X$ such that $A(U) \subset V$. If $p \in P$ and $r > 0$ are such that $B_p(0, r) \subset U$, then

$$\forall x \in X, \quad p(x) \leq r \implies q(Ax) \leq 1. \quad (2.3)$$

By Proposition 1.4 applied to $f(x) = q(Ax)$ and $g(x) = p(x)$, this relation implies that

$$\forall x \in X, \quad q(Ax) \leq \frac{1}{r} p(x). \quad (2.4)$$

Conversely, if A is (P, Q) -bounded, then for every τ_Q -neighborhood V of $0 \in Y$, there exist $q \in Q$ and $R > 0$ such that $B_q(0, R) \subset V$. Let $p \in P$ and $L \geq 0$ be such that the condition (2.2) is fulfilled. Taking $r := R/(L+1)$, we have

$$\forall x \in B_p(0, r), \quad q(Ax) \leq Lp(x) \leq \frac{L}{L+1}R \leq R, \quad (2.5)$$

which shows that $A(B_p(0, r)) \subset B_q(0, R) \subset V$, that is, A is continuous at $0 \in X$.

The implication (3) \Rightarrow (4) follows from the (P, Q) -boundedness of the mapping A and the definition (1.16) of the vicinities.

To prove (4) \Rightarrow (3), suppose that A is $(\mathcal{W}_P, \mathcal{W}_Q)$ -quasiuniformly continuous. For $q \in Q$, let $W = \{(y', y) \in Y \times Y : q(y - y') \leq 1\} \in \mathcal{W}_Q$, and let $U = \{(x', x) \in X \times X : p(x - x') \leq r\} \in \mathcal{W}_P$ be such that $(x', x) \in U$ implies that $(Ax', Ax) \in W$. Taking $x' = 0$, it follows that

$$\forall x \in X, \quad p(x) \leq r \implies q(Ax) \leq 1, \quad (2.6)$$

so that, by Proposition 1.4,

$$\forall x \in X, \quad q(Ax) \leq \frac{1}{r} p(x). \quad (2.7)$$

□

In the case of linear functionals on an asymmetric locally convex space, we have the following characterization of continuity, where u is as in Example 1.1.

PROPOSITION 2.3. *Let (X, P) be an asymmetric locally convex space and $\varphi : X \rightarrow \mathbb{R}$ a linear functional. The following assertions are equivalent.*

- (1) φ is (P, u) -continuous at $0 \in X$.
- (2) φ is (P, u) -continuous on X .
- (3) There exist $p \in P$ and $L \geq 0$ such that

$$\forall x \in X, \quad \varphi(x) \leq Lp(x). \quad (2.8)$$

- (4) φ is upper semicontinuous from (X, τ_P) to $(\mathbb{R}, |\cdot|)$.

Remark 2.4. If the family P is not directed, then the (P, u) -continuity of the functional φ is equivalent to the condition that there exist $F \in \mathcal{F}(P)$ and $L \geq 0$ such that

$$\forall x \in X, \quad \varphi(x) \leq L \max \{p(x) : p \in F\} = Lp_F(x). \quad (2.9)$$

The dual of an asymmetric locally convex space. For an asymmetric locally convex space (X, P) , denote by $X^b = X_P^b$ the set of all linear (P, u) -continuous functionals. If $P = \{p\}$, then we obtain the dual space X_p^b of an asymmetric normed space (X, p) considered in [13].

Let $X^\#$ be the algebraic dual space to X , that is, the space of all linear functionals on X . In contrast to the symmetric case, $X^b = X_P^b$ is not a subspace of $X^\#$, but merely a convex cone, that is,

- (i) $\varphi, \psi \in X^b \Rightarrow \varphi + \psi \in X^b$,
- (ii) $\varphi \in X^b$ and $\alpha \geq 0 \Rightarrow \alpha\varphi \in X^b$.

There are examples in the case $P = \{p\}$ of p -bounded linear functionals φ on a space with asymmetric norm (X, p) such that $-\varphi$ is not p -bounded, see [5]. A simpler example can be exhibited in the space (\mathbb{R}, u) from Example 1.1.

Example 2.5. The identity mapping $\varphi(t) = t$, $t \in \mathbb{R}$, is (τ_u, τ_u) -continuous because

$$\forall t \in \mathbb{R}, \quad \varphi(t) = t \leq \max\{t, 0\} = u(t), \quad (2.10)$$

but $-\varphi$ is not (τ_u, τ_u) -continuous, because it is impossible to find $L \geq 0$ such that $(-\varphi)(t) \leq Lu(t)$ for all $t \in \mathbb{R}$. Indeed, taking $t = -1$, we obtain the contradiction

$$1 = (-\varphi)(-1) \leq L \cdot u(-1) = 0. \quad (2.11)$$

Remark 2.6. It is easy to check that a linear functional $\varphi(t) = at$, $t \in \mathbb{R}$, is (τ_u, τ_u) -continuous if and only if $a \geq 0$. Indeed if $a \geq 0$, then $\varphi(t) = at \leq u(at) = au(t)$, $t \in \mathbb{R}$. If $a < 0$, then, reasoning as above, one concludes that φ fails to be continuous.

Suppose that the family P of asymmetric seminorms is directed, and for $p \in P$, let $p^s(x) = \max\{p(x), p(-x)\}$ be the symmetric seminorm attached to p , and let

$$P^s = \{p^s : p \in P\}. \quad (2.12)$$

Denote by $X^* = (X, P^s)^*$ the dual space of the locally convex space (X, P^s) . Since for a seminorm q and a linear functional φ we have

$$\forall x \in X, \quad \varphi(x) \leq Lq(x) \iff \forall x \in X, \quad |\varphi(x)| \leq Lq(x), \quad (2.13)$$

we have $X^b = X_P^b \subset X^* = (X, P^s)^*$. Indeed, if $\varphi \in X^b$, $p \in P$, and $L \geq 0$ are such that for all $x \in X$, $\varphi(x) \leq Lp(x)$, then, the inequality $p \leq p^s$ and the above equivalence imply that $|\varphi(x)| \leq Lp^s(x)$, $x \in X$, showing that $\varphi \in X^*$.

Let p be an asymmetric seminorm on a real vector space X and let $\varphi : X \rightarrow \mathbb{R}$ be a linear functional. Put

$$\|\varphi\|_p = \sup \varphi(B_p). \quad (2.14)$$

We say that the functional φ is p -bounded if there exists $L \geq 0$ such that

$$\forall x \in X, \quad \varphi(x) \leq Lp(x). \quad (2.15)$$

A number $L \geq 0$ satisfying (2.15) is called a p -Lipschitz constant for φ . The functional φ is p -bounded if and only if $\|\varphi\|_p < \infty$ and $\|\varphi\|_p$ is the smallest p -Lipschitz constant for φ . The p -boundedness of φ is also equivalent to its (τ_p, τ_u) -continuity. The functional $\|\cdot\|_p$ defined by (2.14) is an asymmetric norm on the asymmetric dual X_P^b of (X, p) , that is, $\|\varphi + \psi\|_p \leq \|\varphi\|_p + \|\psi\|_p$, $\|\alpha\varphi\|_p = \alpha\|\varphi\|_p$, for all $\varphi, \psi \in X_P^b$ and $\alpha \geq 0$. Also, $\|\varphi\|_p > 0$ for $\varphi \in X_P^b \setminus \{0\}$.

Similar considerations can be done with respect to the conjugate asymmetric seminorm $\bar{p}(x) = p(-x)$ of p and

$$\|\varphi\|_{\bar{p}} = \sup \varphi(B_{\bar{p}}). \quad (2.16)$$

Some properties of the norm $\|\cdot\|_p$ are collected in the following proposition.

PROPOSITION 2.7. *Let p be an asymmetric seminorm on a real vector space X and $\varphi : X \rightarrow \mathbb{R}$ a linear functional.*

(1) *The following equalities hold:*

$$\|\varphi\|_p = \sup \varphi(B'_p), \quad \|\varphi\|_{\bar{p}} = \sup \varphi(B'_{\bar{p}}). \quad (2.17)$$

Moreover, if the functional $\varphi \neq 0$ is p -bounded, then $\|\varphi\|_p > 0$ and $\varphi(x_0) = \|\varphi\|_p$, for some $x_0 \in B_p$, implies that $p(x_0) = 1$.

(2) *If $\varphi \neq 0$ is (p, \bar{p}) -bounded, then*

$$\varphi(B'_p) = (-\|\varphi\|_{\bar{p}}, \|\varphi\|_p). \quad (2.18)$$

If φ is p -bounded but not \bar{p} -bounded, then

$$\varphi(B'_p) = (-\infty; \|\varphi\|_p). \quad (2.19)$$

Proof. (1) If (x_n) is a sequence in B_p such that $\varphi(x_n) \rightarrow \|\varphi\|_p$ for $n \rightarrow \infty$, then $x'_n = (1 - 1/n)x_n \in B'_p$ and $\varphi(x'_n) \rightarrow \|\varphi\|_p$. Because

$$\sup \varphi(B'_p) \leq \sup \varphi(B_p) = \|\varphi\|_p, \quad (2.20)$$

it follows that $\sup \varphi(B'_p) = \|\varphi\|_p$.

If $\varphi \neq 0$, and $z \in X$ is such that $\varphi(z) > 0$, then the inequality $\varphi(z) \leq \|\varphi\|_p p(z)$ implies that $\|\varphi\|_p > 0$.

If $x_0 \in B_p$ is such that $\varphi(x_0) = \|\varphi\|_p$, then $\|\varphi\|_p p(x_0) \geq \varphi(x_0) = \|\varphi\|_p > 0$, so that $p(x_0) > 0$. If $p(x_0) < 1$, then $x_1 = (1/p(x_0))x_0 \in B_p$ and

$$\varphi(x_1) = \frac{\varphi(x_0)}{p(x_0)} = \frac{\|\varphi\|_p}{p(x_0)} > \|\varphi\|_p = \sup \varphi(B_p), \quad (2.21)$$

a contradiction.

(2) Suppose that φ is (p, \bar{p}) -bounded. We have

$$\begin{aligned} \|\varphi\|_{\bar{p}} &= \sup \{\varphi(x) : \bar{p}(x) < 1\} = \sup \{\varphi(-x) : p(x) < 1\} \\ &= -\inf \{\varphi(x) : p(x) < 1\}. \end{aligned} \quad (2.22)$$

Similar calculations show that

$$\|\varphi\|_{\bar{p}} = -\inf \varphi(B_p). \quad (2.23)$$

Because B'_p is convex, it follows that $\varphi(B'_p)$ is an interval in \mathbb{R} and

$$(-\|\varphi\|_{\bar{p}}; \|\varphi\|_p) \subset \varphi(B'_p) \subset [-\|\varphi\|_{\bar{p}}; \|\varphi\|_p]. \quad (2.24)$$

If $\|\varphi\|_p \in \varphi(B'_p)$, then $\|\varphi\|_p = \varphi(x_0)$, for some $x_0 \in X$ with $p(x_0) < 1$, in contradiction to the assertion (1) of the proposition. Similarly, if $-\|\varphi\|_{\bar{p}} \in \varphi(B'_p)$, then $-\|\varphi\|_{\bar{p}} = \varphi(x_1)$, for some $x_1 \in X$ with $p(x_1) < 1$. But then, for $x'_1 = (1/p(x_1))x_1 \in B_p$, we obtain the contradiction

$$\varphi(x'_1) = \frac{\varphi(x_1)}{p(x_1)} = \frac{-\|\varphi\|_{\bar{p}}}{p(x_1)} < -\|\varphi\|_{\bar{p}} = \inf \varphi(B_p). \quad (2.25)$$

If φ is p -bounded but not \bar{p} -bounded, then

$$\|\varphi\|_{\bar{p}} = \sup \varphi(B'_p) = \infty, \quad (2.26)$$

so that, by (2.22), $\inf \varphi(B'_p) = -\|\varphi\|_{\bar{p}} = -\infty$. Since $\varphi(B'_p)$ is an interval in \mathbb{R} , $\|\varphi\|_p = \sup \varphi(B'_p)$, and $\|\varphi\|_p \notin \varphi(B'_p)$, it follows that $\varphi(B'_p) = (-\infty; \|\varphi\|_p)$. \square

Extension of bounded linear functionals. As in the symmetric case, an extension result for continuous linear functionals defined on subspaces of an asymmetric locally convex space will be particularly useful in developing a duality theory for such spaces.

PROPOSITION 2.8. *Let (X, P) be an asymmetric locally convex space and Y a subspace of X .*

If $\varphi : Y \rightarrow \mathbb{R}$ is a (P, u) -continuous linear functional, then there exists a (P, u) -continuous linear functional $\Phi : X \rightarrow \mathbb{R}$ such that $\Phi|_Y = \varphi$.

Proof. Suppose that the family P is directed. By Proposition 2.3, there exist $p \in P$ and $L \geq 0$ such that

$$\forall y \in Y, \quad \varphi(y) \leq Lp(y). \quad (2.27)$$

By the Hahn-Banach dominated extension theorem, there exists a linear functional $\Phi : X \rightarrow \mathbb{R}$ such that $\Phi|_Y = \varphi$ and

$$\forall x \in X, \quad \Phi(x) \leq Lp(x), \quad (2.28)$$

which, by the same Proposition 2.3, is equivalent to the (P, u) -continuity of Φ . \square

The following existence result is well known in the symmetric case.

PROPOSITION 2.9. (1) *If p is an asymmetric norm on a real vector space X and $x_0 \in X$ is such that $p(x_0) > 0$, then there exists a p -bounded linear functional $\varphi : X \rightarrow \mathbb{R}$ such that*

- (i) $\varphi(x_0) = p(x_0)$,
- (ii) $\|\varphi\|_p = 1$.

(2) *Let (X, P) be an asymmetric locally convex space. If the topology τ_P is Hausdorff, then for every $x_0 \in X$, $x_0 \neq 0$, there exists $\psi \in X^b$ such that $\psi(x_0) = 1$.*

Proof. (1) Let $Z = \mathbb{R}x_0$ and $\varphi_0 : Z \rightarrow \mathbb{R}$ be defined by $\varphi_0(tx_0) = tp(x_0)$, $t \in \mathbb{R}$. Then φ_0 is linear and $\varphi_0(tx_0) = tp(x_0) = p(tx_0)$ for $t \geq 0$. Since $\varphi_0(tx_0) = tp(x_0) < 0 \leq p(tx_0)$ for $t < 0$, it follows that $\varphi_0(z) \leq p(z)$ for all $z \in Z$. By the Hahn-Banach extension theorem, there exists a linear functional $\varphi : X \rightarrow \mathbb{R}$ such that $\varphi|_Z = \varphi_0$ and $\varphi(x) \leq p(x)$ for all $x \in X$, implying that $\|\varphi\|_p \leq 1$. Since

$$\|\varphi\|_p = \sup \{\varphi(x) : x \in B_p\} \geq \sup \{\varphi_0(z) : z \in Z \cap B_p\} \geq \varphi_0\left(\frac{1}{p(x_0)}x_0\right) = 1, \quad (2.29)$$

it follows that $\|\varphi\|_p = 1$.

(2) If $x_0 \neq 0$ and τ_P is Hausdorff, then by Corollary 1.8, there exists $p \in P$ such that $p(x_0) > 0$. If $\varphi : X \rightarrow \mathbb{R}$ is a p -bounded linear functional satisfying the conditions (i) and (ii) of the first assertion, then we can take $\psi = (1/p(x_0))\varphi$. \square

The w^b -topology of the dual X^b . This is the analog of the weak*-topology (w^* -topology) on the dual of a locally convex space. In the case of an asymmetric normed space (X, p) , it was considered in [13].

Let (X, P) be an asymmetric locally convex space and $X^b = X_P^b$ the asymmetric dual cone. A w^b -neighborhood of an element $\varphi \in X^b$ is a subset W of X^b for which there exist

$x_1, \dots, x_n \in X$ and $\epsilon > 0$ such that

$$V_{x_1, \dots, x_n; \epsilon}(\varphi) := \{\psi \in X^b : \psi(x_i) - \varphi(x_i) < \epsilon, i = 1, \dots, n\} \subset W. \quad (2.30)$$

For $x \in X$, define $q_x : X^b \rightarrow [0; \infty)$ by $q_x(\varphi) = \varphi(x)$, $\varphi \in X^b$. Then q_x is additive and positively homogeneous on X^b and

$$V_{x_1, \dots, x_n; \epsilon}(\varphi) = \{\psi \in X^b : q_{x_i}(\psi - \varphi) < \epsilon, i = 1, \dots, n\}. \quad (2.31)$$

The w^b -convergence of a net $\{\varphi_i, i \in I\}$ to $\varphi \in X^b$ is equivalent to the fact that for every $x \in X$, the net $\{(\varphi_i - \varphi)(x), i \in I\}$ converges to 0 in (\mathbb{R}, u) , that is,

$$\forall x \in X, \forall \epsilon > 0, \exists i_0 \in I \text{ such that } \forall i \geq i_0, (\varphi_i - \varphi)(x) < \epsilon. \quad (2.32)$$

Since $X^b \subset X^*$ and

$$V_{x; \epsilon}(\varphi) \cap V_{-x; \epsilon}(\varphi) = \{\psi \in X^b : |(\psi - \varphi)(x)| < \epsilon\}, \quad (2.33)$$

it follows that the w^b -topology on X^b is induced by the w^* -topology of the space X^* .

Asymmetric polars. Let (X, P) be an asymmetric locally convex space, (X, P^s) the associated locally convex space, X^b the asymmetric dual of (X, P) , and $X^* = (X, P^s)^*$ the conjugate space of (X, P^s) .

The *polar* of a nonempty subset Y of (X, P^s) is defined by

$$Y^\circ = \{x^* \in X^* : \forall y \in Y, x^*(y) \leq 1\}. \quad (2.34)$$

Define the corresponding set in the case of the asymmetric dual X^b by

$$Y^\alpha = Y^\circ \cap X^b = \{\varphi \in X^b : \forall y \in Y, \varphi(y) \leq 1\}, \quad (2.35)$$

and call it the *asymmetric polar* of the set Y .

As it is well known, the set Y° is a convex w^* -closed subset of X^* (see, e.g., [19, page 341]). Since the w^b -topology on $X^b \subset X^*$ is induced by the w^* -topology on X^* , we have the following result.

PROPOSITION 2.10. *The asymmetric polar Y^α of a nonempty subset Y of an asymmetric locally convex space (X, P) is a convex w^b -closed subset of X^b .*

In the following proposition, we prove the asymmetric analog of the Alaoglu-Bourbaki theorem, see, for example, [8, Theorem 4.31] or [19, Satz VIII.3.11].

PROPOSITION 2.11. *The asymmetric polar of a neighborhood of the origin of an asymmetric locally convex space (X, P) is a convex w^b -compact subset of the asymmetric dual X^b .*

Proof. Suppose that P is directed. If V is a τ_P -neighborhood of $0 \in X$, then there exist $p \in P$ and $r > 0$ such that $B_p(0, r) \subset V$. Because $p^s(x) \leq r$ implies that $p(x) \leq p^s(x) \leq r$, it follows that $B_{p^s}(0, r) \subset B_p(0, r) \subset V$, so that V is a neighborhood of 0 in the locally convex space (X, P) . By the Alaoglu-Bourbaki theorem (see [19, Satz VIII.3.11]), it follows that

V° is a convex w^* -compact subset of the dual X^* . Since w^\flat -compactness of V^α is equivalent to its w^* -compactness in X^* , it is sufficient to show that the set V^α is w^* -closed in X^* .

Let $\{\varphi_i : i \in I\}$ be a net in V^α that is w^* -convergent to $f \in X^*$. This means that for every $x \in X$, the net $\{\varphi_i(x) : i \in I\}$ converges to $f(x)$ in $(\mathbb{R}, |\cdot|)$. Since for every $v \in V$, $\varphi_i(v) \leq 1$, for all $i \in I$, it follows that $f(v) \leq 1$ for all $v \in V$. Because f is linear, it is sufficient to prove its (P, u) -continuity at $0 \in X$. Consider for some $\epsilon > 0$ the τ_u -neighborhood $(-\infty; \epsilon)$ of $f(0) = 0 \in \mathbb{R}$. Then $U = (\epsilon/2)V$ is a τ_P -neighborhood of $0 \in X$, and for $v \in V$ and $u = (\epsilon/2)v \in U$, we have

$$f(u) = \frac{\epsilon}{2}f(v) \leq \frac{\epsilon}{2} < \epsilon, \quad (2.36)$$

that is, $f(U) \subset (-\infty; \epsilon)$, proving the (P, u) -continuity of f at 0.

It follows that $f \in V^\alpha$, so that V^α is w^* -closed in X^* . \square

3. The continuity of the Minkowski functional and the separation of convex sets

Proposition 2.3 can be extended to sublinear functionals.

PROPOSITION 3.1. *Let (X, P) be an asymmetric locally convex space, where P is a directed family of asymmetric seminorms on X , and let $f : X \rightarrow \mathbb{R}$ be a sublinear functional. The following assertions are equivalent.*

- (1) *The functional f is (P, u) -continuous at $0 \in X$.*
- (2) *The functional f is (P, u) -continuous on X .*
- (3) *There exist $p \in P$ and $L > 0$ such that*

$$\forall x \in X, \quad f(x) \leq Lp(x). \quad (3.1)$$

- (4) *The functional f is upper semicontinuous from (X, τ_P) to $(\mathbb{R}, |\cdot|)$.*

Proof. (1) \Rightarrow (3). Since $(-\infty; 1]$ is a τ_u -neighborhood of $f(0) = 0 \in \mathbb{R}$, there exist $p \in P$ and $r > 0$ such that $f(B_p(0, r)) \subset (-\infty; 1]$, that is,

$$\forall x \in X, \quad p(x) \leq r \implies f(x) \leq 1. \quad (3.2)$$

By Proposition 1.4, this implies that

$$\forall x \in X, \quad f(x) \leq \frac{1}{r}p(x). \quad (3.3)$$

(3) \Rightarrow (2). Let $x_0 \in X$, and for some $\epsilon > 0$, let $(-\infty; f(x_0) + \epsilon)$ be a τ_u -neighborhood of $f(x_0)$. If $p \in P$ and $L \geq 0$ are as in the assertion (3) of the proposition, then $U = x_0 + (\epsilon/(L+1))B_p$ is a τ_P -neighborhood of x_0 , and for every $z \in B_p$ and $x = x_0 + (\epsilon/(L+1))z \in U$, we have

$$f(x) \leq f(x_0) + \frac{\epsilon}{L+1}f(z) \leq f(x_0) + \epsilon \frac{L}{L+1} < f(x_0) + \epsilon, \quad (3.4)$$

proving the (P, u) -continuity of f at x_0 .

Because the implication (2) \Rightarrow (1) is trivial, it follows the equivalence of the first three assertions of the theorem.

As the equivalence (2) \Leftrightarrow (4) holds for any mapping $f : X \rightarrow \mathbb{R}$ (see Remark 1.2), it follows the equivalence of all four assertions of the proposition. \square

The above proposition has the following useful corollary.

COROLLARY 3.2. *Let f, g be sublinear functionals defined on an asymmetric locally convex space (X, P) . If $f \leq g$ and g is (P, u) -continuous, then f is (P, u) -continuous too.*

In particular, the result is true when f is linear.

Proof. By Proposition 3.1, there exist $p \in P$ and $L \geq 0$ such that for all $x \in X$, $g(x) \leq Lp(x)$. It follows that for all $x \in X$, $f(x) \leq g(x) \leq Lp(x)$, which, by the same proposition, implies the continuity of f . \square

Concerning the continuity of the Minkowski functional, we have the following result.

PROPOSITION 3.3. *Let C be a convex absorbing subset of an asymmetric locally convex space (X, P) .*

(1) *The Minkowski functional p_C is (P, u) -continuous if and only if 0 is a τ_P -interior point of C .*

(2) *If p_C is (P, u) -continuous, then*

$$\tau_P\text{-int } C = \{x \in X : p_C(x) < 1\}. \quad (3.5)$$

Proof. Suppose that the family P is directed.

(1) If 0 is a τ_P -interior point of C , then there exist $p \in P$ and $r > 0$ such that

$$B_p(0, r) \subset C \subset \{x \in X : p_C(x) \leq 1\}, \quad (3.6)$$

that is,

$$\forall x \in X, \quad p(x) \leq r \implies p_C(x) \leq 1. \quad (3.7)$$

By Proposition 1.4, we have

$$\forall x \in X, \quad p_C(x) \leq \frac{1}{r}p(x), \quad (3.8)$$

which, by Proposition 3.1, implies the (P, u) -continuity of p_C .

Conversely, suppose that p_C is (P, u) -continuous. Since the set $(-\infty; 1)$ is τ_u -open in \mathbb{R} , the set $\{x \in X : p_C(x) < 1\} = p_C^{-1}(-\infty; 1)$ is τ_P -open, contains 0, and is contained in C , implying that $0 \in \tau_P\text{-int } C$.

(2) If p_C is (P, u) -continuous, then the above inclusion shows that

$$\{x \in X : p_C(x) < 1\} \subset \tau_P\text{-int } C. \quad (3.9)$$

If $x \in \tau_P\text{-int } C$, then there exist $p_1 \in P$ and $r > 0$ such that $B'_{p_1}(x, r) \subset C$. Let $p_2 \in P$ and $L > 0$ be such that for all $x \in X$, $p_C(x) \leq Lp_2(x)$. If $p \in P$ is such that $p \geq p_1, p_2$, $i = 1, 2$, then

$B'_p(x, r) \subset B'_{p_1}(x, r) \subset C$ and

$$\forall x \in X, \quad p_C(x) \leq Lp(x). \quad (3.10)$$

If $p(x) = 0$, then, by the above inequality, $p_C(x) = 0 < 1$. If $p(x) > 0$, put $x_\alpha = (1 + \alpha)x$ for $\alpha > 0$. Since $p(x_\alpha - x) = \alpha p(x) < r$ for $0 < \alpha < r/p(x)$, it follows that $x_\alpha \in C$ for $0 < \alpha < r/p(x)$. But then, for any such α , we have

$$p_C(x) = \frac{1}{1 + \alpha} p_C(x_\alpha) \leq \frac{1}{1 + \alpha} < 1. \quad (3.11) \quad \square$$

The separation of convex sets. The separation results for convex subsets of locally convex spaces are key tools in the study of duality for these spaces as well as in optimization problems.

In the following two theorems, we prove the asymmetric analogs of the classical separation theorems of Eidelheit and Tukey (see [15, Theorems 2.2.26 and 2.2.28]).

THEOREM 3.4. *Let (X, P) be an asymmetric locally convex space. If Y_1, Y_2 are two disjoint nonempty convex subsets of X with Y_1 τ_P -open, then there exists a linear functional $\varphi \in X^b$ such that*

$$\forall y_1 \in Y_1, \forall y_2 \in Y_2 \quad \varphi(y_1) < \varphi(y_2). \quad (3.12)$$

Proof. Let $y_i^0 \in Y_i$, $i = 1, 2$, and let $x_0 = y_2^0 - y_1^0$. Since the set Y_1 is τ_P -open and the topology τ_P is translation invariant, the set

$$Y := x_0 + Y_1 - Y_2 = \cup \{x_0 - y_2 + Y_1 : y_2 \in Y_2\} \quad (3.13)$$

is τ_P -open too.

We have $0 = x_0 + y_1^0 - y_2^0 \in Y$ and $x_0 \notin Y$. Indeed, if $x_0 = x_0 + y_1 - y_2$, for some $y_1 \in Y_1$ and $y_2 \in Y_2$, then the element $y = y_1 - y_2$ would belong to the empty set $Y_1 \cap Y_2$.

By the preceding proposition, the Minkowski functional p_Y of the τ_P -open convex set Y is sublinear, (P, u) -continuous, and $p_Y(x_0) \geq 1$ since $x_0 \notin Y$. By Proposition 2.9(1), there exists a p_Y -bounded linear functional $\psi : X \rightarrow \mathbb{R}$ such that $\psi(x_0) = p_Y(x_0)$ and $\psi(x) \leq p_Y(x)$, $x \in X$. Taking $\varphi = (1/p_Y(x_0))\psi$, it follows that

$$\varphi(x_0) = 1, \quad \forall x \in X, \quad \varphi(x) = \frac{1}{p_Y(x_0)} \psi(x) \leq \frac{1}{p_Y(x_0)} p_Y(x) \leq p_Y(x). \quad (3.14)$$

By Proposition 2.3, the functional φ is (P, u) -continuous. Because Y is τ_P -open and $0 \in Y$, by Proposition 3.3, we have $Y = \{x \in X : p_Y(x) < 1\}$. Since $\varphi(x_0) = 1$, we obtain

$$\forall y_1 \in Y_1, \forall y_2 \in Y_2 \quad 1 + \varphi(y_1) - \varphi(y_2) = \varphi(x_0 + y_1 - y_2) \leq p_Y(x_0 + y_1 - y_2) < 1, \quad (3.15)$$

implying that

$$\forall y_1 \in Y_1, \forall y_2 \in Y_2, \quad \varphi(y_1) < \varphi(y_2). \quad (3.16) \quad \square$$

We prove now the asymmetric analog of Tukey's separation theorem.

THEOREM 3.5. *Let (X, P) be an asymmetric locally convex space. If Y_1, Y_2 are two nonempty disjoint convex subsets of X with Y_1 τ_P -compact and Y_2 τ_P -closed, then there exists a functional $\varphi \in X^\flat$ such that*

$$\sup \varphi(Y_1) < \inf \varphi(Y_2). \quad (3.17)$$

Proof. Suppose that P is directed.

Since $Y_1 \cap Y_2 = \emptyset$ and Y_2 is τ_P -closed, for every $y \in Y_1$, there exist $p_y \in P$ and $r_y > 0$ such that

$$(y + 2r_y B'_{p_y}) \cap Y_2 = \emptyset. \quad (3.18)$$

The τ_P -open cover $\{y + r_y B'_{p_y} : y \in Y_1\}$ of the τ_P -compact set Y_1 contains a finite subcover $\{y_k + r_k B'_{p_k} : k = 1, 2, \dots, n\}$, where $p_k = p_{y_k}$ and $r_k = r_{y_k}$ for $k = 1, \dots, n$. Take $p \in P$ such that $p \geq p_k$, $k = 1, 2, \dots, n$, put $r := \min\{r_k : k = 1, 2, \dots, n\}$ and show that

$$(Y_1 + r B'_p) \cap Y_2 = \emptyset. \quad (3.19)$$

Indeed, if $y' = y + ru$ for some $y \in Y_1$, $u \in B'_p$, and $y' \in Y_2$, then, choosing $k \in \{1, 2, \dots, n\}$ such that $y \in y_k + r_k B'_{p_k}$, we have

$$y' = y + ru \in y_k + r B'_p + r_k B'_{p_k} \subset y_k + r_k B'_{p_k} + r_k B'_{p_k} = y_k + 2r_k B'_{p_k}, \quad (3.20)$$

in contradiction to (3.18).

The set $Z := Y_1 + r B'_p$ is convex, τ_P -open, and disjoint from Y_2 . By Theorem 3.4, there exists $\varphi \in X^\flat$ such that

$$\forall y \in Y_1, \forall u \in B'_p, \forall y' \in Y_2 \quad \varphi(y) + r\varphi(u) < \varphi(y'). \quad (3.21)$$

By Proposition 2.3, there exist $q_1 \in P$ and $L > 0$ such that for all $x \in X$, $\varphi(x) \leq Lq_1(x)$. If $q \in P$ is such that $q \geq \max\{p, q_1\}$, then $\varphi(x) \leq Lq(x)$, $x \in X$, and $B'_q \subset B'_p$, so that

$$\forall y \in Y_1, \forall u \in B'_q, \forall y' \in Y_2 \quad \varphi(y) + r\varphi(u) < \varphi(y'). \quad (3.22)$$

By (3.21), $\varphi \neq 0$, so that by Proposition 2.7, $\|\varphi\|_q = \sup \varphi(B'_q) > 0$. Passing in (3.22) to supremum with respect to $u \in B'_q$, we get

$$\forall y \in Y_1, \forall y' \in Y_2 \quad \varphi(y) + r\|\varphi\|_q \leq \varphi(y'), \quad (3.23)$$

implying that

$$r\|\varphi\|_q + \sup \varphi(Y_1) \leq \inf \varphi(Y_2). \quad (3.24)$$

It follows that

$$\sup \varphi(Y_1) < \inf \varphi(Y_2). \quad (3.25)$$

□

Remark 3.6. The inequality in Theorem 3.4 cannot be reversed, in the sense that, under the same hypotheses on the sets Y_1 and Y_2 , we cannot find a (P, u) -continuous linear functional ψ on X such that

$$\forall y_2 \in Y_2, \forall y_1 \in Y_1 \quad \psi(y_2) < \psi(y_1). \quad (3.26)$$

This is due, on one side, to the fact that the functional $-\varphi$ need not be (P, u) -continuous, where φ is the linear functional given by Theorem 3.4. On the other side, analyzing the proof of Theorem 3.4, it follows that we should work with the set $Y' := x_0 + Y_2 - Y_1$ which need not be τ_P -open, because the τ_P -openness of Y_1 does not imply the τ_P -openness of $-Y_1$, see Example 1.3.

The same caution must be taken when applying Theorem 3.5.

Extreme points and the Krein-Milman theorem. We start by recalling some notions and facts. A point e of a convex subset of a vector space X is called an *extreme point* of Y provided that $(1-t)x + ty = e$, for some $x, y \in Y$ and $0 < t < 1$, implies that $x = y = e$. A nonempty convex subset Z of Y is called an *extremal subset* of Y if $(1-t)x + ty \in Z$, for some $x, y \in Y$ and $0 < t < 1$, implies that $x, y \in Z$ (in fact, $[x; y] \subset Z$, by the convexity of Z). Obviously, that a one-point set $Z = \{e\}$ is an extremal subset of Y if and only if e is an extreme point of Y . Also, if W is an extremal subset of the extremal subset Z of Y , then W is an extremal subset of Y too. In particular, if e is an extreme point of an extremal subset Z of Y , then e is an extreme point of Y . The intersection of a family of extremal subsets of Y is an extremal subset of Y provided that it is nonempty. We denote by $\text{ext } Y$ the (possibly empty) set of extreme points of the convex set Y .

The following proposition is an immediate consequence of the definitions.

PROPOSITION 3.7. *Let Y be a nonempty convex subset of a vector space X and f a linear functional on X . If the set $Z = \{z \in Y : f(z) = \sup f(Y)\}$ is nonempty, then it is an extremal subset of Y . A similar assertion holds for the set $W = \{w \in Y : f(w) = \inf f(Y)\}$.*

We can state and prove now the Krein-Milman theorem in the asymmetric case.

THEOREM 3.8. *Let (X, P) be an asymmetric locally convex space such that the topology τ_P is Hausdorff. Then any nonempty convex τ_P -compact subset Y of X coincides with the τ_P -closed convex hull of the set of its extreme points*

$$Y = \tau_P\text{-cl-co}(\text{ext } Y). \quad (3.27)$$

Proof. All the topological notions will concern the τ_P -topology of X so that we will omit “ τ_P -” in the following. By Proposition 2.9(2), for every $x \in X$, $x \neq 0$, there exists $\varphi \in X^\flat$ with $\varphi(x) = 1$.

Fact 3.9. Every nonempty convex compact subset Z of X has an extreme point.

Let

$$\mathcal{F} := \{F : F \text{ is a closed extremal subset of } Z\}, \quad (3.28)$$

and define the order in \mathcal{F} by $F_1 \leq F_2 \Leftrightarrow F_1 \subset F_2$ and show that the set \mathcal{F} is nonempty and downward inductively ordered. Because Y is τ_P -compact and the topology τ_P is Hausdorff, it follows that Y is convex and τ_P -closed, so that $Y \in \mathcal{F}$. Since a totally ordered subfamily \mathcal{G} of \mathcal{F} has the finite intersection property, by the compactness of the set Z , the set $G = \cap \mathcal{G}$ is nonempty, closed, and extremal. Therefore, $G \in \mathcal{F}$ is a lower bound for \mathcal{G} . By Zorn's lemma, the ordered set \mathcal{F} has a minimal element F_0 . If we show that F_0 is a one-point set, $F_0 = \{x_0\}$, then x_0 will be an extreme point of Z .

Suppose that F_0 contains two distinct points x_1, x_2 , and let $p \in P$ be such that $p(x_1 - x_2) > 0$. Let φ be a p -bounded linear functional such that $\varphi(x_1 - x_2) = p(x_1 - x_2) > 0$ (see Proposition 2.9(1)). It follows that $\varphi \in X^\flat$, so that φ is upper semicontinuous as a mapping from (X, τ_P) to $(\mathbb{R}, |\cdot|)$. By the compactness of the set F_0 , the set

$$F_1 = \{x \in F_0 : \varphi(x) = \sup \varphi(F_0)\} = \{x \in F_0 : \varphi(x) \geq \sup \varphi(F_0)\} \quad (3.29)$$

is nonempty and closed. By Proposition 3.7, F_1 is an extremal subset of F_0 , thus an extremal subset of Z . Therefore, $F_1 \in \mathcal{F}$, $F_1 \subset F_0$, and $x_2 \in F_0 \setminus F_1$ in contradiction to the minimality of F_0 .

Fact 3.10. $Y = \tau_P\text{-cl co}(\text{ext } Y)$.

The inclusion $\text{ext}(Y) \subset Y$ implies that $Y_1 := \tau_P\text{-cl co}(\text{ext } Y) \subset Y$. As a closed subset of a compact set, the set Y_1 is convex and compact. Supposing that there exists a point $y_0 \in Y \setminus Y_1$, then, by Theorem 3.5, there exists $\varphi \in X^\flat$ such that

$$\sup \varphi(Y_1) < \varphi(y_0). \quad (3.30)$$

Using again the upper semicontinuity of φ as a mapping from (X, τ_P) to $(\mathbb{R}, |\cdot|)$, we see that the set

$$F = \{y \in Y : \varphi(y) = \sup \varphi(Y)\} = \{y \in Y : \varphi(y) \geq \sup \varphi(Y)\} \quad (3.31)$$

is nonempty, convex, and compact, so that, by Fact 3.9, it has an extreme point e_1 . Since F is an extremal subset of Y , it follows that e_1 is an extreme point of Y , implying that $e_1 \in Y_1$. Taking into account (3.30), we obtain the contradiction

$$\sup \varphi(Y) = \varphi(e_1) \leq \sup \varphi(Y_1) < \varphi(y_0) \leq \sup \varphi(Y). \quad (3.32)$$

□

4. The asymmetric weak topology

The weak topology of a locally convex space (X, Q) is defined by the locally convex basis \mathcal{W} formed by the sets of the form

$$V'_{x_1^*, \dots, x_n^*, \epsilon} = \{x \in X : |x_i^*(x)| < \epsilon, 1 \leq i \leq n\}, \quad (4.1)$$

for $n \in \mathbb{N}$, $x_1^*, \dots, x_n^* \in X^*$ and $\epsilon > 0$. Obviously, we can suppose that $x_i^* \neq 0$, $i = 1, \dots, n$.

The duality theory for locally convex spaces is based on the following key lemma of algebraic nature.

LEMMA 4.1 (see [19, Lemma VIII.3.3]). *Let X be a vector space and $f, f_1, \dots, f_n : X \rightarrow \mathbb{R}$ linear functionals. The following assertions are equivalent.*

- (1) $f \in \text{sp}\{f_1, \dots, f_n\}$.
- (2) *There exists $L \geq 0$ such that*

$$\forall x \in X, \quad f(x) \leq L \max \{ |f_1(x)|, \dots, |f_n(x)| \}. \quad (4.2)$$

- (3) $\bigcap_{i=1}^n \ker f_i \subset \ker f$.

In our case, this lemma takes the form.

LEMMA 4.2. *Let f, f_1, \dots, f_n be real linear functionals on a vector space X , with f_1, \dots, f_n linearly independent. Then the following assertions are equivalent.*

- (1) *For all $x \in X$, $f_i(x) \leq 0$, $i = 1, \dots, n \Rightarrow f(x) \leq 0$.*
- (2) *There exists $L \geq 0$ such that for all $x \in X$, $f(x) \leq L \max\{f_i(x) : 1 \leq i \leq n\}$.*
- (3) *There exist $a_1, \dots, a_n \geq 0$, such that $f = \sum_{i=1}^n a_i f_i$.*

Proof. Since the implications (2) \Rightarrow (1) and (3) \Rightarrow (2) are obvious, it is sufficient to prove (1) \Rightarrow (3).

If $f_i(x) = 0$ for $i = 1, \dots, n$, then $f_i(-x) = -f_i(x) = 0$, $i = 1, \dots, n$, so that $f(x) \leq 0$ and $-f(x) = f(-x) \leq 0$, implying that $f(x) = 0$. Therefore Lemma 4.1(3) is fulfilled, so that there exist $a_1, \dots, a_n \in \mathbb{R}$ such that $f = \sum_{i=1}^n a_i f_i$. It remains to show that $a_j \geq 0$ for $j = 1, \dots, n$. Because f_1, \dots, f_n are linearly independent, there exist the elements $x_j \in X$ such that $f_i(x_j) = -\delta_{ij} \leq 0$, $i, j = 1, 2, \dots, n$, where δ_{ij} is the Kronecker symbol. It follows that $f(x_j) \leq 0$ and

$$-a_j = \sum_{i=1}^n a_i f_i(x_j) = f(x_j) \leq 0, \quad (4.3)$$

for $j = 1, \dots, n$. □

Define the asymmetric weak topology w^α on an asymmetric locally convex space (X, P) as the asymmetric locally convex topology generated by the asymmetric locally convex basis ${}^a\mathcal{W}'_\alpha$ formed by the sets

$$V'_{\varphi_1, \dots, \varphi_n; \epsilon} = \{x \in X : \varphi_i(x) < \epsilon, 1 \leq i \leq n\}, \quad (4.4)$$

for $n \in \mathbb{N}$, $\varphi_1, \dots, \varphi_n \in X^b$ and $\epsilon > 0$. The neighborhoods of an arbitrary point $x \in X$ are subsets of X containing a set of the form $x + V'_{\varphi_1, \dots, \varphi_n; \epsilon} = \{x' \in X : \varphi_i(x' - x) < \epsilon, 1 \leq i \leq n\}$.

The sets

$$V_{\varphi_1, \dots, \varphi_n; \epsilon} = \{x \in X : \varphi_i(x) \leq \epsilon, 1 \leq i \leq n\} \quad (4.5)$$

generate the same topology.

In the following proposition, we collect some properties of the topology w^α .

PROPOSITION 4.3. Let (X, P) be an asymmetric locally convex space and $X^\flat = (X, P)^\flat$ its asymmetric dual cone.

- (1) The topology τ_P is finer than w^α .
- (2) For $\varphi \in X^\flat$ and $\epsilon > 0$, the set $\{x \in X : \varphi(x) < \epsilon\}$ is w^α -open and $\{x \in X : \varphi(x) \geq \epsilon\}$ is w^α -closed.
- (3) A net $\{x_i : i \in I\}$ in X is w^α -convergent to $x \in X$ if and only if for every $\varphi \in X^\flat$, the net $\{\varphi(x_i)\}$ converges to $\varphi(x)$ in (\mathbb{R}, u) . This means the following:

$$\forall \varphi \in X^\flat, \forall \epsilon > 0, \exists i_0 \text{ such that } \forall i \geq i_0, \varphi(x_i - x) < \epsilon. \quad (4.6)$$

- (4) The asymmetric dual $(X, w^\alpha)^\flat$ of the asymmetric locally convex space (X, w^α) agrees with X^\flat .

Proof. Suppose that P is directed.

- (1) Let $V' = V'_{\varphi_1, \dots, \varphi_n; \epsilon}$ be an element of the locally convex basis (4.4). Because φ_i are (P, u) -continuous, there exist $p_i \in P$ and $L_i \geq 0$ such that

$$\forall x \in X, \varphi_i(x) \leq L_i p_i(x), \text{ for } i = 1, \dots, n. \quad (4.7)$$

The multiball $U' = \{x \in X : p_i(x) < \epsilon/(L+1), 1 \leq i \leq n\}$, where $L = \max L_i$, is contained in V' , showing that V' is a τ_P -neighborhood of $0 \in X$.

- (2) If $V = \{x \in X : \varphi(x) < \epsilon\}$ and $x_0 \in V$, then the w^α -neighborhood $\{x \in X : \varphi(x - x_0) < \epsilon - \varphi(x_0)\}$ of x_0 is contained in V' because

$$\varphi(x - x_0) < \epsilon - \varphi(x_0) \implies \varphi(x) = \varphi(x - x_0) + \varphi(x_0) < \epsilon. \quad (4.8)$$

The assertion (3) follows from definitions.

- (4) Because τ_P is finer than w^α , the identity map $\text{Id} : (X, \tau_P) \rightarrow (X, w^\alpha)$ is continuous, implying the (P, u) -continuity of $\varphi \circ \text{Id}$ for any $\varphi \in (X, w^\alpha)^\flat$, that is, $(X, w^\alpha)^\flat \subset (X, P)^\flat$.

Conversely, if φ is a (P, u) -continuous linear functional, then the set $V = \{x \in X : \varphi(x) < \epsilon\}$ is a w^α -neighborhood of $0 \in X$ and $\varphi(V) \subset (-\infty; \epsilon)$ for every $\epsilon > 0$, proving the (w^α, τ_u) -continuity of φ at 0, and by the linearity of φ , on the whole X . \square

As in the symmetric case, the closed convex sets are the same for the topologies τ_P and w^α .

PROPOSITION 4.4. Let (X, P) be an asymmetric locally convex space and Y a convex subset of X . Then Y is w^α -closed if and only if it is τ_P -closed.

Proof. Because τ_P is finer than w^α , it follows that any (not necessarily convex) w^α -closed subset of X is also τ_P -closed.

Suppose now that the convex set Y is τ_P but not w^α -closed. If x_0 is a point in $w^\alpha\text{-cl } Y \setminus Y$, then, applying Theorem 3.5 to the sets $\{x_0\}$ and Y , we get a functional $\varphi \in X^\flat$ such that

$$\varphi(x_0) < \inf \varphi(Y). \quad (4.9)$$

If $m := \inf \varphi(Y)$, then $V = \{x \in X : \varphi(x - x_0) < 2^{-1}(m - \varphi(x_0))\}$ is a w^α -neighborhood of x_0 . Because

$$\varphi(x) = \varphi(x - x_0) + \varphi(x_0) < \frac{m + \varphi(x_0)}{2} < m, \quad (4.10)$$

for every $x \in V$, it follows that $V \cap Y = \emptyset$, in contradiction to $x_0 \in w^\alpha\text{-cl } Y$. \square

The proposition has the following corollary.

COROLLARY 4.5. *Let (X, P) be an asymmetric locally convex space. Then for every subset Z of X , the following equality holds:*

$$w^\alpha\text{-cl co}(Y) = \tau_P\text{-cl co}(Y). \quad (4.11)$$

Proof. By the definition of the closed convex hull and the preceding proposition, we have the equalities

$$\begin{aligned} w^\alpha\text{-cl co}(Y) &= \bigcap \{Y : Y \subset X, Y \text{ convex and } w^\alpha\text{-closed}\} \\ &= \bigcap \{Y : Y \subset X, Y \text{ convex and } \tau_P\text{-closed}\} \\ &= \tau_P\text{-cl co}(Y). \end{aligned} \quad (4.12) \quad \square$$

Remark 4.6. We can define the asymmetric polar of a subset W of the dual X^b of an asymmetric locally convex space (X, P) by

$$W_\alpha = \{x \in X : \forall \varphi \in W, \varphi(x) \leq 1\}. \quad (4.13)$$

Since, for $\varphi \in X^b$, a set of the form $\{x \in X : \varphi(x) \leq 1\}$ is not necessarily τ_P -closed, the set W_α need not be τ_P -closed. Therefore, an asymmetric analog of the bipolar theorem (see [19, Satz VIII.3.9]), asserting that

$$(A^\circ)_\circ = -\text{cl co } (A \cup \{0\}) \quad (4.14)$$

for any subset A of a locally convex space (X, Q) , does not hold in the asymmetric case.

References

- [1] C. Alegre, J. Ferrer, and V. Gregori, *Quasi-uniformities on real vector spaces*, Indian J. Pure Appl. Math. **28** (1997), no. 7, 929–937.
- [2] ———, *On the Hahn-Banach theorem in certain linear quasi-uniform structures*, Acta Math. Hungar. **82** (1999), no. 4, 325–330.
- [3] A. R. Alimov, *The Banach-Mazur theorem for spaces with nonsymmetric distance*, Uspekhi Mat. Nauk **58** (2003), no. 2(350), 159–160 (Russian), translated in Russian Math. Surveys **58** (2003), no. 2, 367–369.
- [4] A. Andrikopoulos, *Completeness in quasi-uniform spaces*, Acta Math. Hungar. **105** (2004), no. 1–2, 151–173.
- [5] P. A. Borodin, *The Banach-Mazur theorem for spaces with an asymmetric norm and its applications in convex analysis*, Mat. Zametki **69** (2001), no. 3, 329–337 (Russian), translated in Math. Notes **69** (2001), no. 3–4, 298–305.

- [6] S. Cobzaş, *Separation of convex sets and best approximation in spaces with asymmetric norm*, Quaest. Math. **27** (2004), no. 3, 275–296.
- [7] S. Cobzaş and C. Mustăta, *Extension of bounded linear functionals and best approximation in spaces with asymmetric norm*, Rev. Anal. Numér. Théor. Approx. **33** (2004), no. 1, 39–50.
- [8] M. Fabian, P. Habala, P. Hájek, V. Montesinos Santalucía, J. Pelant, and V. Zizler, *Functional Analysis and Infinite-Dimensional Geometry*, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, vol. 8, Springer, New York, 2001.
- [9] J. Ferrer, V. Gregori, and C. Alegre, *Quasi-uniform structures in linear lattices*, Rocky Mountain J. Math. **23** (1993), no. 3, 877–884.
- [10] P. Fletcher and W. F. Lindgren, *Quasi-Uniform Spaces*, Lecture Notes in Pure and Applied Mathematics, vol. 77, Marcel Dekker, New York, 1982.
- [11] L. M. García-Raffi, S. Romaguera, and E. A. Sánchez Pérez, *Sequence spaces and asymmetric norms in the theory of computational complexity*, Math. Comput. Modelling **36** (2002), no. 1–2, 1–11.
- [12] ———, *On Hausdorff asymmetric normed linear spaces*, Houston J. Math. **29** (2003), no. 3, 717–728.
- [13] ———, *The dual space of an asymmetric normed linear space*, Quaest. Math. **26** (2003), no. 1, 83–96.
- [14] ———, *Weak topologies on asymmetric normed linear spaces and non-asymptotic criteria in the theory of complexity analysis of algorithms*, J. Anal. Appl. **2** (2004), no. 3, 125–138.
- [15] R. E. Megginson, *An Introduction to Banach Space Theory*, Graduate Texts in Mathematics, vol. 183, Springer, New York, 1998.
- [16] C. Mustăta, *A Phelps type theorem for spaces with asymmetric norms*, Bul. Ştiinț. Univ. Baia Mare Ser. B Fasc. Mat.-Inform. **18** (2002), no. 2, 275–280.
- [17] ———, *On the uniqueness of the extension and unique best approximation in the dual of an asymmetric normed linear space*, Rev. Anal. Numér. Théor. Approx. **32** (2003), no. 2, 187–192.
- [18] S. Romaguera and M. Schellekens, *Duality and quasi-normability for complexity spaces*, Appl. Gen. Topol. **3** (2002), no. 1, 91–112.
- [19] D. Werner, *Funktionalanalysis [Functional Analysis]*, 3rd, revised and extended ed., Springer, Berlin, 2000.

S. Cobzaş: Department of Mathematics, Faculty of Mathematics and Computer Science, Babeş-Bolyai University, 400084 Cluj-Napoca, Romania

E-mail address: scobzas@math.ubbcluj.ro

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk