

ELLIPTIC GW INVARIANTS OF BLOWUPS ALONG CURVES AND SURFACES

JIANXUN HU AND HOU-YANG ZHANG

Received 27 January 2004

We established a relation between elliptic Gromov-Witten invariants of a symplectic manifold M and its blowups along smooth curves and surfaces.

1. Introduction

Over the last few years, many mathematicians contributed their efforts to establish the mathematical foundation of the theory of quantum cohomology or Gromov-Witten (GW) invariants. In 1995, Ruan and Tian [13, 15] first established for the semipositive symplectic manifolds. Recently, the semipositivity condition has been removed by many authors. Now, the focus turned to the calculations and applications. Many Fano manifolds were computed. We think that it is important to study the change of GW invariants under surgery.

Some recent research indicated that there is a deep amazing relation between quantum cohomology and birational geometry. The quantum minimal model conjecture [14] leads to attempt to find quantum cohomology of a minimal model without knowing minimal model. This problem requires a thorough understanding of blowup type formula of GW invariants and quantum cohomology.

According to McDuff [11], the blowup operation in symplectic geometry amounts to a removal of an open symplectic ball followed by a collapse of some boundary directions. Lerman [8] gave a generalization of blowup construction, “the symplectic cut.”

Let M be a compact symplectic manifold of dimension $2n$, \tilde{M} the blowup of M along a smooth submanifold. Denote by $p : \tilde{M} \rightarrow M$ the natural projection. Denote by $\Psi_{(A,g)}^M(\alpha_1, \dots, \alpha_m)$ the genus g GW invariant of M . The authors refer the interested reader to [15] for the definition of GW invariants.

In [5, 6], we mainly concentrated on the changes of genus zero GW invariants under blowup along smooth curves and surfaces. In this paper, we mainly concentrate on the elliptic GW invariants. These invariants were first discussed in [1] and since then have been studied in various contexts, see [3, 12]. An elegant recursion was predicted in [2] using the method of Virasoro constraints. In this paper, we will generalize some results

82 Elliptic GW invariants of blowups along curves and surfaces

about the cases of blowing up along curves and surfaces to the case of genus-one. Our main results are the following.

THEOREM 1.1. *Suppose that C is a smooth curve in M such that either its genus $g_0 \geq 2$ or $g_0 \leq 1$ and $C_1(M)(C) \geq 0$, where $C_1(M)$ denotes the first Chern classes of M . $A \in H_2(M)$ such that $p!(A) = PDp^*PD(A)$ is a nonexceptional class in $H_2(\tilde{M})$, $\alpha_i \in H^*(M)$, $1 \leq i \leq m$ satisfy either $\deg \alpha_i \neq 1$ or $\deg \alpha_i = 1$ and support away from C . Then,*

$$\Psi_{(A,1)}^M(\alpha_1, \dots, \alpha_m) = \Psi_{(p!(A),1)}^{\tilde{M}}(p^*\alpha_1, \dots, p^*\alpha_m). \quad (1.1)$$

About the changes of GW invariants of blow up of symplectic manifold along a smooth surface, in this paper, we will assume that the symplectic manifold is semipositive. Our result is the following theorem.

THEOREM 1.2. *Suppose that M is a semipositive compact symplectic manifold and S is a smooth surface in M . If $A \in H_2(M)$ such that $p!(A) = PDp^*PD(A)$ is a nonexceptional class in $H_2(\tilde{M})$, $\alpha_i \in H^*(M)$, $1 \leq i \leq m$, satisfy either $\deg \alpha_i > 2$ or $\deg \alpha_i \leq 2$ and support away from S . Then, (1.1) holds.*

2. General review of gluing formula

The proof of our result is an application of the gluing formula for symplectic cutting developed by Li and Ruan [9]. In algebraic geometry, Li [10] proved a completely analogous degeneration formula of GW invariants. Another symplectic version of the gluing formula of GW-invariants is due to Ionel and Parker [7].

Suppose that $H : M^0 \rightarrow \mathbf{R}$ is a proper Hamiltonian function such that the Hamiltonian vector field X_H generates a circle action, where $M^0 \subset M$ is an open domain. By adding a constant, we may assume that 0 is a regular value. Then, $H^{-1}(0)$ is a smooth submanifold preserved by the circle action. The quotient $Z = H^{-1}(0)/S^1$ is the famous symplectic reduction. Namely, it has an induced symplectic structure. We can cut M along $H^{-1}(0)$. Suppose that we obtain two disjoint components M^\pm which have the boundary $H^{-1}(0)$. We can collapse the S^1 -action on $H^{-1}(0)$ to obtain \overline{M}^\pm containing a real codimension two submanifold $Z = H^{-1}(0)/S^1$, see [9, Section 2 and pages 156–157] for details. There is a map

$$\pi : M \longrightarrow \overline{M}^+ \cup_Z \overline{M}^-, \quad (2.1)$$

where $\overline{M}^+ \cup_Z \overline{M}^-$ is the union of \overline{M}^\pm along Z .

To formulate the gluing formula, we need the terminology of relative GW invariants. Here, we copy the definition of [9, Section 4 and page 157]. We define a relative GW invariant $\Psi^{(M,Z)}$ by counting the number of relative stable holomorphic maps intersecting Z at finitely many points with prescribed tangency. Let $T_m = (t_1, \dots, t_m)$ be a set of nonnegative integers such that $\sum_i t_i = Z^*(A)$, where Z^* is the Poincare dual of Z and $A \in H_2(M)$. We order them such that $t_1 = \dots = t_l = 0$ and $t_i > 0$ for $i > l$. Consider the moduli space $\mathcal{M}_A(g, T_m)$ of genus g pseudoholomorphic maps f such that f has marked points (x_1, \dots, x_m) with the property that f is tangent to Z at x_i with order t_i . Here, $t_i = 0$ means that there is no intersection. Then, we compactify $\mathcal{M}_A(g, T_m)$ by $\overline{\mathcal{M}}(g, T_m)$,

the space of relative stable maps. We have evaluation map

$$e_i : \overline{\mathcal{M}}_A(g, T_m) \longrightarrow M \quad (2.2)$$

for $i \leq l$ and

$$e_j : \overline{\mathcal{M}}_A(g, T_m) \longrightarrow Z \quad (2.3)$$

for $j > l$. Roughly, the relative GW-invariants are defined as

$$\Psi_{(A, g, T_m)}^{(M, Z)}(\alpha_1, \dots, \alpha_l; \beta_{l+1}, \dots, \beta_m) = \int_{\overline{\mathcal{M}}_A(g, T_m)}^{\text{vir}} \Pi_i e_i^* \alpha_i \wedge \Pi_j e_j^* \beta_j. \quad (2.4)$$

Let $u = (u^+, u^-) : (\Sigma^+, \Sigma^-) \rightarrow (M^+, M^-)$ be J -holomorphic curves such that u^+ and u^- have ν ends and they converge to the same periodic orbits at each end. Suppose that Σ^\pm have m^\pm marked points, respectively. Here, Σ^\pm may not be connected, see [9, Section 4]. Suppose that $\Sigma = \Sigma^+ \vee \Sigma^-$ has genus g and $[u(\Sigma)] = A$. If we consider the index of the operators $D_{u^\pm} = D\partial_J u^\pm$, see [9, Section 4] for its definition, then we have the following proposition.

PROPOSITION 2.1 (see [9, Theorem 5.1]).

$$\text{Ind } D_{u^+} + \text{Ind } D_{u^-} = 2(n-1)\nu + 2C_1(A) + 2(n-3)(1-g) + 2m, \quad (2.5)$$

where C_1 is the first Chern class of M and $m = m^+ + m^-$.

Suppose that the homology classes of u^+, u^-, u are A^+, A^-, A , respectively. If (u^+, u^-) is another representative and glue to u' , by [9, Lemma 2.11], we have $[u'] = [u]$. Denote by $C = \{A^+, g^+, K^+; A^-, g^-, K^-\}$ the gluing component, where $K^\pm = (0, \dots, 0, k_1, \dots, k_\nu)$ with m^\pm zeros. The gluing formula of Li and Ruan counted the contribution of the gluing components to GW invariant of M . Denote by Ψ_C the contribution of C .

Choose a homology basis $\{\beta_b\}$ of $H^*(Z, \mathbf{R})$. Let (δ_{ab}) be its intersection matrix. For the gluing component $C = \{A^+, g^+, K^+; A^-, g^-, K^-\}$, we have the following gluing formula.

PROPOSITION 2.2 (see [9, Theorem 5.8]). *Let α_i^\pm be differential forms with $\deg \alpha_i^+ = \deg \alpha_i^-$ even. Suppose that $\alpha_i^+|_Z = \alpha_i^-|_Z$ and hence $\alpha_i^+ \cup_Z \alpha_i^- \in H^*(\bar{M}^+ \cup_Z \bar{M}^-; \mathbf{R})$. Let $\alpha_i = \pi^*(\alpha_i^+ \cup_Z \alpha_i^-)$. The following gluing formula holds:*

$$\Psi_C(\alpha_1, \dots, \alpha_{m^+ + m^-}) = |K| \sum_{I, J} \Psi_{(A^+, g^+, K^+)}^{(\bar{M}^+, Z)}(\alpha^+, \beta_I) \delta^{IJ} \Psi_{(A^-, g^-, K^-)}^{(\bar{M}^-, Z)}(\alpha^-, \beta_J), \quad (2.6)$$

where $\beta_i \delta^{i,j} \beta_j$ is associated to every intersection with Z and $|K| = k_1 \dots k_\nu$, $\delta^{IJ} = \delta^{i,j} \dots \delta^{i_\nu, j_\nu}$, $\Psi_{(A^\pm, g^\pm, K^\pm)}^{(\bar{M}^\pm, Z)}(\alpha^\pm, \beta_I)$ denote the product of relative invariants corresponding to each component.

PROPOSITION 2.3 (see [9, Remark 5.5]). *For $C = \{A^\pm, g^\pm, (0, \dots, 0)\}$,*

$$\Psi_C(\alpha_i^\pm) = \Psi_{(A^\pm, g^\pm, (0, \dots, 0))}^{(\bar{M}^\pm, Z)}(\alpha_i^\pm). \quad (2.7)$$

3. Proof of main theorem

Proof of Theorem 1.1. Since C is a smooth curve in M , the normal bundle N_C is a symplectic vector bundle. By symplectic neighborhood theorem, there is a tubular neighborhood $\mathcal{N}_\delta(C)$ of C which is symplectomorphic to the normal bundle N_C . We perform the symplectic cutting as in [5, Section 2.1]. We obtained

$$\overline{M}^+ = \mathbf{P}(N_C \oplus \mathbb{O}), \quad \overline{M}^- = \tilde{M}. \quad (3.1)$$

From the divisor property, the skew symmetry of GW invariants and our assumptions, if we choose a sufficiently small $\delta > 0$, without loss of generality, we may assume $\alpha_i^+ = 0$.

Similar to the proof of [5, Theorem 1.2], by the gluing formula of GW invariant, we first consider the contribution of each component to the GW invariants. Therefore, we consider the gluing component

$$C = \{A^+, g^+, K^+; A^-, g^-, K^-\}, \quad (3.2)$$

where $K^\pm = (0, \dots, 0, k_1, \dots, k_\nu)$ with m^\pm many zeros, respectively. From Proposition 2.1, we have

$$\text{Ind}D_{u^+} + \text{Ind}D_{u^-} = 2(n-1)\nu + 2C_1(A) + 2m, \quad (3.3)$$

since $g = 1$ in our case.

We will use the same convention as [5]. We assume that $u^\pm : \Sigma^\pm \rightarrow M^\pm$ may have l^\pm connected components $u_i^\pm : \Sigma_i^\pm \rightarrow M^\pm$, $i = 1, \dots, l^\pm$. Suppose Σ_i^\pm have arithmetic genus g_i^\pm , $g^\pm = \sum g_i^\pm$, with m_i^\pm marked points and $m^\pm = \sum m_i^\pm$. From [5, Remark 2.3], it is not difficult to see that \tilde{u}_i^+ can be identified as a stable J -holomorphic curve h_i^+ in \overline{M}^+ . Then, from [5, Proposition 2.4], we have

$$\begin{aligned} \text{Ind}D_{u^+} &= \sum_{i=1}^{l^+} \text{Ind}D_{u_i^+} = \sum_{i=1}^{l^+} \text{Ind}D_{\tilde{u}_i^+} \\ &= 2 \sum_{i=1}^{l^+} C_1[h_i^+] + 2(n-3)(l^+ - g^+) \\ &\quad + 2\nu - 2 \sum k_i + 2m^+. \end{aligned} \quad (3.4)$$

Now, we want to calculate $C_1[h_i^+]$ in two cases of our theorem.

Case 1. The genus $g_0 \geq 2$.

In this case, we claim that all stable J -holomorphic maps h_i^+ can only stay in fibers of $\overline{M}^+ = \mathbf{P}(N_C \oplus \mathbb{O})$. Otherwise, suppose that there is a stable J -holomorphic curve $h_i^+ : \Sigma \rightarrow \overline{M}^+$ which does not stay in a fiber. Denote by $\pi : \mathbf{P}(N_C \oplus \mathbb{O}) \rightarrow C$ the projection of the projective bundle. Then, we have a stable J -holomorphic map $\pi \circ h_i^+ : \Sigma \rightarrow C$ satisfying $[\pi \circ h_i^+] \neq 0$. We can perform pregluing as in [9, Section 5, pages 208–209] and obtain a system of small perturbed J -holomorphic curves $f_n : \Sigma_n \rightarrow C$ which represent the class $[\pi \circ h_i^+]$ and satisfy the perturbed Cauchy-Riemann equation $\bar{\partial}_J f_n = \nu_n$, here

Σ_n is a smooth Riemann surface. Actually, we can choose $\nu_n \rightarrow 0$ as $n \rightarrow \infty$. Therefore, by Gromov compactness theorem, we have that f_n weakly converges to a (possibly reducible) J -holomorphic curve $u = (u^1, \dots, u^N)$ and $[\pi \circ h_i^+] = \sum_{j=1}^N [u^j] \neq 0$. Therefore, we have a nonconstant J -holomorphic curve $f : \Sigma_1 \rightarrow C$ and Σ_1 has genus less than 2. It is well known that if $f' : S \rightarrow S'$ is a holomorphic map between compact Riemann surfaces, then the genus of S and S' satisfies $g(S) \geq g(S')$ unless f' is constant (see [4, page 219]). Since $g(C) = g_0 \geq 2$, we have a contradiction. So, our claim is true.

A simple index calculation [5] shows that $C_1[h_i^+] = n \sum k_j$, where summation runs over ends of component u_i^+ . In this case, we have

$$\text{Ind} D_{u^+} = 2(n-3)(l^+ - g^+) + 2(n-1) \sum k_i + 2\nu + 2m^+. \quad (3.5)$$

Case 2. $g_0 \leq 1$ and $C_1(M)(C) \geq 0$.

A simple calculation shows, that $C_1(\mathbf{P}(N_C \oplus \mathbb{O})) = C_1(C) + C_1(N_C) + n\xi = C_1(M) + n\xi$, here ξ is the class of infinite section in $\mathbf{P}(N_C \oplus \mathbb{O})$ over C . Therefore, from our positive assumption, an intersection multiplicity calculation shows that

$$\sum_{i=1}^{l^+} C_1[h_i^+] \geq n \sum k_i. \quad (3.6)$$

In this case, we have

$$\text{Ind} D_{u^+} \geq 2(n-3)(l^+ - g^+) + 2(n-1) \sum k_i + 2\nu + 2m^+. \quad (3.7)$$

Summarizing the above two cases, from (3.4), we have

$$\begin{aligned} \text{Ind} D_{u^+} &\geq 2(n-3)(l^+ - g^+) + 2(n-1) \sum k_i + 2\nu + 2m^+, \\ \text{Ind} D_{u^-} &\leq 2C_1(A) + 2(n-3)(g^+ - l^+) \\ &\quad + 2(n-1)(\nu - \sum k_i) - 2\nu + 2m^-. \end{aligned} \quad (3.8)$$

Since $\alpha_i^+ = 0$, $1 \leq i \leq m$, if $m^+ > 0$, we have for any $\beta_b \in H^*(Z)$,

$$\Psi_{(A^+, g^+, K^+)}^{(\overline{M}^+, Z)}(\alpha_i^+, \beta_b) = 0. \quad (3.9)$$

This implies $\Psi_C = 0$ except $m^- = m$. So, we may assume $m^- = m$. We also may assume

$$\sum \deg \alpha_i = 2C_1(A) + 2m. \quad (3.10)$$

Otherwise, Theorem 1.1 follows from the degree reason. Therefore, we have

$$\begin{aligned} \sum \deg(\alpha_i^-) &= 2C_1(A) + 2m \\ &> 2C_1(A) + 2(n-3)(g^+ - l^+) \\ &\quad + 2(n-1)(\nu - \sum k_i) - 2\nu + 2m^- \\ &\geq \text{Ind} D_{u^-}, \end{aligned} \quad (3.11)$$

86 Elliptic GW invariants of blowups along curves and surfaces

since $g^+ \leq g = 1$, $\nu > 0$, $k_i > 0$, $n \geq 3$. Therefore, by the definition of relative GW invariants, we have for any $\beta_b \in H^*(Z)$,

$$\Psi_{(A^-, g^-, K^-)}^{(\overline{M}^-, Z)}(\alpha_i^-, \beta_b) = 0. \quad (3.12)$$

Therefore, $\Psi_C = 0$ except $C = \{A^-, 1, m\}$.

So, now it remains to show that

$$\Psi_{(p!(A), 1)}^{\tilde{M}}(p^* \alpha_1, \dots, p^* \alpha_m) = \Psi_{(A^-, 1, m)}^{(\overline{M}^-, Z)}(\alpha_1^-, \dots, \alpha_m^-). \quad (3.13)$$

To prove this, we perform the symplectic cutting for \tilde{M} around the exceptional divisor E . Therefore, we have

$$\overline{\tilde{M}}^+ = \mathbf{P}(N_E \oplus \mathbb{O}), \quad \overline{\tilde{M}}^- \cong \tilde{M}. \quad (3.14)$$

Now, we use the gluing formula to prove the contribution of stable J -holomorphic curves in \tilde{M} which touch the exceptional divisor E to the GW invariants of \tilde{M} is zero. We consider the gluing component

$$C = \{p!(A)^+, g^+, K^+; p!(A)^-, g^-, K^-\}. \quad (3.15)$$

Since $\alpha_i^+ = 0$, $1 \leq i \leq m$, we have $\Psi_C = 0$ except

$$K^+ = (k_1, \dots, k_\nu), \quad K^- = \left(\underbrace{0, \dots, 0}_m, k_1, \dots, k_\nu \right). \quad (3.16)$$

From Proposition 2.1, we have (3.3), where C_1 denotes the first Chern class of M .

We assume that $u^\pm : \Sigma^\pm \rightarrow M^\pm$ has l^\pm connected components $u_i^\pm : \Sigma_i^\pm \rightarrow M^\pm$, $i = 1, \dots, l^\pm$. From [5, Remark 2.3], it is not difficult to see that \tilde{u}_i^+ can be identified as stable J -holomorphic curve h_i^+ in \tilde{M} . Then, from [5, Proposition 2.4], we have

$$\text{Ind } D_{u^+} = \sum_{i=1}^{l^+} \text{Ind } D_{\tilde{u}_i^+} = (2n-6)l^+ + 2 \sum_{i=1}^{l^+} C_1[h_i^+] + 2\nu - 2 \sum k_i, \quad (3.17)$$

where C_1 is the first Chern class of $\overline{\tilde{M}}^+$.

Let V be a complex rank r vector bundle over X , and $\pi : \mathbf{P}(V) \rightarrow X$ the corresponding projective bundle. Let ξ_V be the first Chern class of the tautological line bundle in $\mathbf{P}(V)$. A simple calculation shows that

$$C_1(\mathbf{P}(V)) = \pi^* C_1(X) + \pi^* C_1(V) - r\xi_V. \quad (3.18)$$

Note that $\overline{\tilde{M}}^+ = \mathbf{P}(N_E \oplus \mathbb{O})$ and $E = \mathbf{P}(N_C)$. Applying (3.18) to $\overline{\tilde{M}}^+$ and E , we obtain

$$\begin{aligned} C_1(\overline{\tilde{M}}^+) &= C_1(E) + C_1(N_E) - 2\xi \\ &= C_1(C) + C_1(N_C) - (n-1)\xi_1 + C_1(N_E) + 2\xi, \end{aligned} \quad (3.19)$$

where ξ_1 and ξ are the first Chern classes of the tautological line bundles in $\mathbf{P}(N_C)$ and $\mathbf{P}(N_E \oplus \mathbb{O})$, respectively. Here, we denote Chern class and its pullback by the same symbol. It is well known that the normal bundle to E in \tilde{M} is just the tautological bundle on $E \cong \mathbf{P}(N_C)$. Therefore, $C_1(N_E) = \xi_1$. So, we have

$$C_1(\tilde{M}^+) = C_1(M) - (n-2)\xi_1 - 2\xi. \quad (3.20)$$

We know that \tilde{M} is a projective bundle over E with fiber \mathbf{P}^1 . Let L be the class of a line in the fiber \mathbf{P}^1 and e the class of a line in the fiber \mathbf{P}^{n-2} in $E = \mathbf{P}(N_C)$. Denote by $[h_i^+]^C$ the homology class of the projection in C of the curve h_i^+ . Denote by $[h_i^+]^F$ the difference of $[h_i^+]$ and $[h_i^+]^C$, that is, $[h_i^+]^F = [h_i^+] - [h_i^+]^C$. Then, it is easy to know $[h_i^+]^F = aL + be$. Since $\xi \cdot [h_i^+] = \sum k_j$, where the summation runs over ends of u_i^+ , and $E \cdot [h_i^+] = 0$, so we have $\xi \cdot [h_i^+]^F = a = \sum k_j$ and $E \cdot [h_i^+]^F = a - b = 0$. Therefore, we have $a = b = \sum k_j$. So, we have $[h_i^+]^F = \sum k_j(L + e)$. For Case 1, we have $[h_i^+]^C = 0$. Therefore, we have

$$\sum_{i=1}^{l^+} C_1[h_i^+] = 2(n-1) \sum k_i. \quad (3.21)$$

For Case 2, since $C_1(C) + C_1(N_C) \geq 0$, we have

$$\sum_{i=1}^{l^+} C_1[h_i^+] \geq 2(n-1) \sum k_i. \quad (3.22)$$

Plugging in (3.17), we have

$$\text{Ind}D_{u^+} \geq 2(n-3)(l^+ - g^+) + 2(2n-3) \sum k_i + 2\nu. \quad (3.23)$$

Therefore,

$$\begin{aligned} \text{Ind}D_{u^-} &\leq 2C_1(A) + 2(n-3)(g^+ - l^+) \\ &\quad + (2n-2)\left(\nu - \sum k_i\right) - 2(n-2) \sum k_i + 2m. \end{aligned} \quad (3.24)$$

From the degree reason, we also may assume

$$\sum \deg(p^* \alpha_i) = 2C_1(A) + 2m. \quad (3.25)$$

Then,

$$\begin{aligned} \sum \deg(p^* \alpha_i) &= 2C_1(A) + 2m \\ &> 2C_1(A) + 2(n-3)(g^+ - l^+) \\ &\quad + (2n-2)\left(\nu - \sum k_i\right) - 2(n-2) \sum k_i + 2m \\ &\geq \text{Ind}D_{u^-}, \end{aligned} \quad (3.26)$$

since $\nu > 0$, $k_i > 0$. Therefore, by the definition of relative GW invariants, we have for any $\beta_b \in H^*(Z)$,

$$\Psi_{(p!(A)^-, 1, K^-)}^{(\overline{M}^-, Z)}((p^* \alpha_i)^-, \beta_b) = 0. \quad (3.27)$$

Therefore, the contribution of J -holomorphic curves to the GW invariant is nonzero only if it does not touch the exceptional divisor E , that is, $C = \{p!(A)^-, 1, m\}$. Therefore, we have

$$\Psi_{(p!(A), 1)}^{\tilde{M}}(p^* \alpha_1, \dots, p^* \alpha_m) = \Psi_{(p!(A)^-, 1, m)}^{(\overline{M}^-, Z)}((p^* \alpha_1)^-, \dots, (p^* \alpha_m)^-). \quad (3.28)$$

Since $\overline{M}^- = \tilde{M} = \overline{M}^-$ and $p!(A)$ is a nonexceptional class, we may identify the homology class $p!(A)^-$ with A^- . Hence, Theorem 1.1 follows. \square

Proof of Theorem 1.2. Since S is a smooth surface, the normal bundle N_S is a symplectic vector bundle. By symplectic neighborhood theorem, there is a tubular neighborhood $\mathcal{N}_\delta(S)$ of S which is symplectomorphic to the normal bundle N_S . We perform the symplectic cutting as in [5, Section 2.1]. We obtain

$$\overline{M}^+ = \mathbf{P}(N_S \oplus \mathbb{O}), \quad \overline{M}^- = \tilde{M}. \quad (3.29)$$

We may assume $\alpha_i^+ = 0$ if we choose a sufficiently small $\delta > 0$ because of the assumption of α_i .

Similar to the proof of Theorem 1.1, we first consider the contribution of each gluing component to the GW-invariants. Therefore, we consider the component (3.2). From Proposition 2.1, we have (3.3).

We assume $u^\pm : \Sigma^\pm \rightarrow \overline{M}^\pm$ has l^\pm connected components $u_i^\pm : \Sigma_i^\pm \rightarrow \overline{M}^\pm$, $i = 1, \dots, l^\pm$. Suppose Σ_i^\pm has arithmetic genus g_i^\pm , $g^\pm = \sum g_i^\pm$ with m^\pm marked points and $m^\pm = \sum m_i^\pm$.

The similar calculation to that in the proof of Theorem 1.1 shows that

$$\text{Ind } D_{u^+} = 2(n-3)(l^+ - g^+) + 2(n-2) \sum k_i + 2\nu + 2m^+. \quad (3.30)$$

Therefore, we have

$$\text{Ind } D_{u^-} = 2C_1(A) + 2(n-3)(g^+ - l^+) + 2(n-2) \left(\nu - \sum k_i \right) + 2m^-. \quad (3.31)$$

The same argument as in the proof of Theorem 1.1 shows that the contribution of the component C to the GW-invariant of M is nonzero only if C has (3.16). We also assume

$$\sum \deg \alpha_i = 2C_1(A) + 2m. \quad (3.32)$$

The same argument as in the proof of Theorem 1.1 shows that there is a symplectic cutting such that $\Psi_C = 0$ except $C = \{A^-, 1, m\}$. From the gluing theorem, we have

$$\Psi_{(A, 1)}^M(\alpha_1, \dots, \alpha_m) = \Psi_{(A^-, 1, m)}^{(\overline{M}^-, Z)}(\alpha_1^-, \dots, \alpha_m^-). \quad (3.33)$$

Now, it remains to prove

$$\Psi_{(p!(A), 1)}^{\tilde{M}}(p^* \alpha_1, \dots, p^* \alpha_m) = \Psi_{(A^-, 1, m)}^{(\overline{M}^-, Z)}(\alpha_1^-, \dots, \alpha_m^-). \quad (3.34)$$

To prove this, we perform the symplectic cutting for \tilde{M} around E as in the proof of Theorem 1.1. Therefore, we have (3.14).

We also use the gluing theorem to prove that the contribution of stable J -holomorphic curves in \tilde{M} which touch the exceptional divisor E to the GW invariant of \tilde{M} is zero. We consider the component (3.15).

Since $\alpha_i^+ = 0$, $1 \leq i \leq m$, we have $\Psi_C = 0$ except (3.16). Similar calculation to that in the proof of Theorem 1.1 shows that

$$\begin{aligned} \text{Ind } D_{u^+} &= 2(n-3)(l^+ - g^+) + 2\nu + 2(2n-5) \sum k_i, \\ \text{Ind } D_{u^-} &= 2C_1(A) + 2(n-3)(g^+ - l^+) \\ &\quad + 2(n-2)\left(\nu - \sum k_i\right) - 2(n-3) \sum k_i + 2m. \end{aligned} \quad (3.35)$$

The same argument as in the proof of Theorem 1.1 shows (3.28).

The rest of the proof is the same as that of the proof of Theorem 1.1. So, we omit it. This completes the proof of Theorem 1.2. \square

Acknowledgments

The first author would like to thank Professor Yongbin Ruan for his many suggestive discussions and encouragement. The first author also thanks Max-Planck-Institut für Mathematik in Bonn for providing an excellent research environment and support. This work was partially supported by NSF of China (1017114 and 10231050).

References

- [1] M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, *Holomorphic anomalies in topological field theories (with an appendix by S. Katz)*, Nuclear Phys. B **405** (1993), no. 2-3, 279–304.
- [2] T. Eguchi, K. Hori, and C.-S. Xiong, *Quantum cohomology and Virasoro algebra*, Phys. Lett. B **402** (1997), no. 1-2, 71–80.
- [3] E. Getzler, *Intersection theory on $\overline{\mathcal{M}}_{1,4}$ and elliptic Gromov-Witten invariants*, J. Amer. Math. Soc. **10** (1997), no. 4, 973–998.
- [4] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Wiley-Interscience [John Wiley & Sons], New York, 1978.
- [5] J. Hu, *Gromov-Witten invariants of blow-ups along points and curves*, Math. Z. **233** (2000), no. 4, 709–739.
- [6] ———, *Gromov-Witten invariants of blow-ups along surfaces*, Compositio Math. **125** (2001), no. 3, 345–352.
- [7] E.-N. Ionel and T. H. Parker, *Gromov-Witten invariants of symplectic sums*, Math. Res. Lett. **5** (1998), no. 5, 563–576.
- [8] E. Lerman, *Symplectic cuts*, Math. Res. Lett. **2** (1995), no. 3, 247–258.
- [9] A.-M. Li and Y. Ruan, *Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds*, Invent. Math. **145** (2001), no. 1, 151–218.
- [10] J. Li, *A degeneration formula of GW-invariants*, J. Differential Geom. **60** (2002), no. 2, 199–293.

90 Elliptic GW invariants of blowups along curves and surfaces

- [11] D. McDuff, *Blow ups and symplectic embeddings in dimension 4*, Topology **30** (1991), no. 3, 409–421.
- [12] R. Pandharipande, *A geometric construction of Getzler’s elliptic relation*, Math. Ann. **313** (1999), no. 4, 715–729.
- [13] Y. Ruan, *Topological sigma model and Donaldson-type invariants in Gromov theory*, Duke Math. J. **83** (1996), no. 2, 461–500.
- [14] ———, *Surgery, quantum cohomology and birational geometry*, Northern California Symplectic Geometry Seminar (Y. Eliashberg, D. Fuchs, T. Ratiu, and A. Weinstein, eds.), Amer. Math. Soc. Transl. Ser. 2, vol. 196, American Mathematical Society, Rhode Island, 1999, pp. 183–198.
- [15] Y. Ruan and G. Tian, *A mathematical theory of quantum cohomology*, J. Differential Geom. **42** (1995), no. 2, 259–367.

Jianxun Hu: Department of Mathematics, Zhongshan University, Guangzhou 510275, China
E-mail address: stsjxhu@zsu.edu.cn

Hou-Yang Zhang: Department of Mathematics, The Science College, Beijing Jiaotong University, Beijing 100044, China
E-mail address: lpshao@center.njtu.edu.cn

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru