

# K-BESSEL FUNCTIONS ASSOCIATED TO A 3-RANK JORDAN ALGEBRA

HACEN DIB

*Received 2 January 2005 and in revised form 8 September 2005*

Using the Bessel-Muirhead system, we can express the  $K$ -Bessel function defined on a Jordan algebra as a linear combination of the  $J$ -solutions. We determine explicitly the coefficients when the rank of this Jordan algebra is three after a reduction to the rank two. The main tools are some algebraic identities developed for this occasion.

## 1. Introduction

Bessel functions of matrix argument appeared as a subject of studies in the work of Herz [6]. One can find in the multivariate statistics literature some applications of these functions (see [11]). The main tool of Herz's work was the Laplace transform and its inverse in the space of real symmetric matrices. He obtained several properties but a "good" differential system was lacking. This was the major contribution of Muirhead in [10] when he characterized them by a system of second-order partial differential equations and proved the uniqueness (up to multiplicative constant) of the solution which is analytic at 0. Later, Faraut and Travaglini [5] gave a generalization of these functions to a Jordan algebra. An extensive study was done in [2]. However, the explicit resolution of the Bessel-Muirhead system in general rank remains an open problem. Nevertheless, in [9] Mahmoud wrote down an explicit basis of the solutions in the rank 2 and 3 using series of one-variable Bessel functions. On the other hand, the  $K$ -Bessel function of matrix argument was defined earlier in Herz's paper cited above and his conjecture was that there must be a linear relation between this kind of Bessel functions and the  $J$ 's one as known in the one-variable theory. The first result concerning this conjecture was established by the author in [3] for the rank two. In this paper, we continue our work and prove that a similar result for the  $K$ -Bessel function is also true when the Jordan algebra is of rank 3. In this case, there are four nonequivalent classes of real simple and Euclidean Jordan algebra:  $\text{Herm}(3, \mathbb{F})$  the algebra of  $3 \times 3$  Hermitian matrices, where  $\mathbb{F}$  is the field of real, complex, quaternionic, or Cayley (octaves) numbers. In [3] we intended to perform a case-by-case calculation. In this way, a serious difficulty arises in the evaluation of some integral over the automorphism group of the Jordan algebra. However, a unified treatment is possible

by a reduction to the rank-2 case using some algebraic identities. This is what we will present here.

## 2. Preliminaries

Let us recall the situation and some useful results in [3].

*Definition 2.1.* Bessel-Muirhead operators are defined by

$$B_i = x_i \frac{\partial^2}{\partial x_i^2} + (\nu + 1) \frac{\partial}{\partial x_i} + 1 + \frac{d}{2} \sum_{j \neq i} \frac{1}{x_i - x_j} \left( x_i \frac{\partial}{\partial x_i} - x_j \frac{\partial}{\partial x_j} \right), \quad 1 \leq i \leq r, \quad (2.1)$$

where  $r$  is the rank of the system,  $\nu$  a complex parameter, and  $d$  an integer (which will be interpreted below in Section 3). A Bessel function is a symmetric function  $f$  which is a solution of the Bessel-Muirhead system  $B_i f = 0$ ,  $i = 1, 2, \dots, r$ .

Denote by  $t_1, t_2, \dots, t_r$  the elementary symmetric functions, that is,

$$t_p = \sum_{1 \leq i_1 < i_2 < \dots < i_p \leq r} x_{i_1} x_{i_2} \cdots x_{i_p} \quad (2.2)$$

with  $t_0 = 1$  and  $t_p = 0$  if  $p < 0$  or  $p > r$ . The Bessel-Muirhead system is then equivalent to the system (see [2, 7])  $Z_k g = 0$ ,  $1 \leq k \leq r$ , where

$$Z_k = \sum_{i,j=1}^r A_{ij}^k \frac{\partial^2}{\partial t_i \partial t_j} + \left( \nu + 1 + \frac{r-k}{2} d \right) \frac{\partial}{\partial t_k} + \delta_k^1, \quad (2.3)$$

$$A_{ij}^k = \begin{cases} t_{i+j-k} & \text{if } i, j \geq k, \\ -t_{i+j-k} & \text{if } i, j < k, i+j \geq k, \\ 0 & \text{elsewhere.} \end{cases}$$

Here  $\delta_k^1$  is the Kronecker symbol and  $g(t_1, t_2, \dots, t_r) = f(x_1, x_2, \dots, x_r)$ . When  $r = 2$ , we have (see [3]) a fundamental set of solutions given by  $J_\nu^{[2,1]}$ ,  $J_\nu^{[2,2]}$ ,  $t_2^{-\nu} J_{-\nu}^{[2,1]}$ , and  $t_2^{-\nu} J_{-\nu}^{[2,2]}$ , where

$$J_\nu^{[2,1]}(t_1, t_2) = \sum_{m_1, m_2 \geq 0} \frac{(-1)^{m_1}}{(1)_{m_1} (1)_{m_2} (1+\nu)_{m_2} \left( 1 + \nu + d/2 \right)_{m_1+2m_2}} t_1^{m_1} t_2^{m_2}, \quad (2.4)$$

$$J_\nu^{[2,2]}(t_1, t_2) = \sum_{m_1, m_2 \geq 0} \frac{(-1)^{m_1}}{(1-\nu-d/2)_{m_1} (1)_{m_2} (1+\nu)_{m_2} (1)_{m_1+2m_2}} t_1^{m_1-\nu-d/2} t_2^{m_2}.$$

Here  $(a)_k$  is the classical Pochhammer symbol, that is,  $(a)_k = a(a+1) \cdots (a+k-1)$ . In this case, the  $K$ -Bessel function (in two variables) can be expressed (in the J-basis) as follows (see [3]):

$$K_\nu^{[2]}(t_1, t_2) = \sum_{j=1}^2 a_\nu^j J_\nu^{[2,j]}(-t_1, t_2) + b_\nu^j t_2^{-\nu} J_{-\nu}^{[2,j]}(-t_1, t_2), \quad (2.5)$$

where

$$\begin{aligned} a_\nu^1 &= b_{-\nu}^1 = (2\pi)^{(n-2)/2} \Gamma(-\nu) \Gamma\left(-\nu - \frac{n-2}{2}\right), \\ a_\nu^2 &= b_{-\nu}^2 = (2\pi)^{(n-2)/2} \Gamma(-\nu) \Gamma\left(\nu + \frac{n-2}{2}\right) \end{aligned} \quad (2.6)$$

(for the meaning of  $n$ , see Section 3 below). Observe that this expression reduces (up to constant factor) to the classical one-variable formula when  $t_2 = 0$  and generic  $\nu$  (i.e.,  $\nu \notin \mathbb{Z}$  and  $\nu \pm (n-2)/2 \notin \mathbb{Z}$ ). Now, for  $r = 3$ , we have eight linearly independent  $J$ -solutions  $J_\nu^{[3,1]}, J_\nu^{[3,2]}, J_\nu^{[3,3]}, J_\nu^{[3,4]}, t_3^{-\nu} J_{-\nu}^{[3,1]}, t_3^{-\nu} J_{-\nu}^{[3,2]}, t_3^{-\nu} J_{-\nu}^{[3,3]},$  and  $t_3^{-\nu} J_{-\nu}^{[3,4]}$ , where

$$\begin{aligned} J_\nu^{[3,1]}(t_1, t_2, t_3) &= \sum_{m_1, m_2, m_3 \geq 0} \frac{(-1)^{m_1+m_3}}{(1)_{m_1} (1)_{m_2} (1)_{m_3} (1+\nu)_{m_3} (1+\nu+d/2)_{m_2+2m_3}} \\ &\quad \times \frac{(1+2\nu+d)_{m_1+2m_2+4m_3}}{(1+\nu+d)_{m_1+2m_2+3m_3} (1+2\nu+d)_{m_1+2m_2+3m_3}} t_1^{m_1} t_2^{m_2} t_3^{m_3}, \\ J_\nu^{[3,2]}(t_1, t_2, t_3) &= \sum_{m_1, m_2, m_3 \geq 0} \frac{(-1)^{m_1+m_3}}{(1-\nu-d)_{m_1} (1)_{m_2} (1)_{m_3} (1+\nu)_{m_3} (1+\nu+d/2)_{m_2+2m_3}} \\ &\quad \times \frac{(1+\nu)_{m_1+2m_2+4m_3}}{(1)_{m_1+2m_2+3m_3} (1+\nu)_{m_1+2m_2+3m_3}} t_1^{m_1-\nu-d} t_2^{m_2} t_3^{m_3}, \\ J_\nu^{[3,3]}(t_1, t_2, t_3) &= \sum_{m_1, m_2, m_3 \geq 0} \frac{(-1)^{m_1+m_3}}{(1)_{m_1} (1-\nu-d/2)_{m_2} (1)_{m_3} (1+\nu)_{m_3} (1)_{m_2+2m_3}} \\ &\quad \times \frac{(1)_{m_1+2m_2+4m_3}}{(1-\nu)_{m_1+2m_2+3m_3} (1)_{m_1+2m_2+3m_3}} t_1^{m_1} t_2^{m_2-\nu-d/2} t_3^{m_3}, \\ J_\nu^{[3,4]}(t_1, t_2, t_3) &= \sum_{m_1, m_2, m_3 \geq 0} \frac{(-1)^{m_1+m_3}}{(1+\nu)_{m_1} (1-\nu-d/2)_{m_2} (1)_{m_3} (1+\nu)_{m_3} (1)_{m_2+2m_3}} \\ &\quad \times \frac{(1+\nu)_{m_1+2m_2+4m_3}}{(1)_{m_1+2m_2+3m_3} (1+\nu)_{m_1+2m_2+3m_3}} t_1^{m_1+\nu} t_2^{m_2-\nu-d/2} t_3^{m_3}. \end{aligned} \quad (2.7)$$

Observe also that when  $t_3 = 0$  (and  $\nu \notin \mathbb{Z}$ ,  $\nu \pm d/2 \notin \mathbb{Z}$ ), these functions reduce to

$$J_\nu^{[3,1]}(t_1, t_2, 0) = J_{\nu+d/2}^{[2,1]}(t_1, t_2), \quad (2.8)$$

$$J_\nu^{[3,2]}(t_1, t_2, 0) = J_{\nu+d/2}^{[2,2]}(t_1, t_2), \quad (2.9)$$

$$J_\nu^{[3,3]}(t_1, t_2, 0) = t_2^{-\nu-d/2} J_{-\nu-d/2}^{[2,1]}(t_1, t_2), \quad (2.10)$$

$$J_\nu^{[3,4]}(t_1, t_2, 0) = t_2^{-\nu-d/2} J_{-\nu-d/2}^{[2,2]}(t_1, t_2). \quad (2.11)$$

This behaviour and the expression of  $K_\nu^{[2]}$  are an important step to the final result.

### 3. Some algebraic identities

For the general theory of Jordan algebra, one can see [4], but what we will develop is somehow specific to the rank three. So, let  $A$  be a real simple and Euclidean Jordan

algebra with rank 3 and real dimension  $n$ . We know that  $n = 3 + 3d$ , where  $d = 1, 2, 4$ , or 8. The integer  $d$  is the real dimension of some decomposition spaces which appear in the Peirce decomposition of the algebra  $A$  (see [4]). Let  $\{c_1, c_2, c_3\}$  be a complete system of mutually orthogonal primitive idempotents, that is,  $c_i c_j = \delta_i^j c_i$ ,  $c_1 + c_2 + c_3 = e$  the unit of  $A$  and none of the  $c_j$ 's can split into a sum of two idempotents. We have a Cayley-Hamilton-like theorem  $x^3 - a_1(x)x^2 + a_2(x)x - a_3(x)e = 0$  and a spectral decomposition  $x = k \cdot (\lambda_1 c_1 + \lambda_2 c_2 + \lambda_3 c_3)$  with  $k$  an element of the automorphism group of  $A$  and  $\lambda_i$  reals such that

$$\begin{aligned} a_1(x) &= \lambda_1 + \lambda_2 + \lambda_3 := \text{tr}(x), \\ a_3(x) &= \lambda_1 \lambda_2 \lambda_3 := \det(x), \\ a_2(x) &= \lambda_1 \lambda_2 + \lambda_1 \lambda_3 + \lambda_2 \lambda_3 = \frac{1}{2} [\text{tr}(x)^2 - \text{tr}(x^2)]. \end{aligned} \tag{3.1}$$

The inner product is defined then by  $(x, y) := \text{tr}(xy)$ . The operators  $L(x)$  and  $P(x)$  are defined by  $L(x)y = xy$  and  $P(x) = 2L^2(x) - L(x^2)$ . Let us consider the Peirce decomposition with respect to the idempotent  $c_3$ , that is,  $A = A_0 \oplus A_{1/2} \oplus A_1$ , where  $A_\alpha$  is the eigenspace of  $L(c_3)$  with respect to the eigenvalue  $\alpha$ .  $A_0$  and  $A_1$  are Jordan subalgebras of rank 2 and 1, respectively, and  $A_{1/2}$  is a subspace of dimension  $2d$ . Put  $n_0 = \dim A_0 = 2 + d$  and  $e_0 = c_1 + c_2$  the unit of  $A_0$ . We have  $A_0 A_1 = \{0\}$ ,  $(A_0 \oplus A_1) A_{1/2} \subset A_{1/2}$  and  $A_{1/2} A_{1/2} \subset A_0 \oplus A_1$ . If we write  $\text{tr}(z)$  or  $\det(z)$  of an element of  $A_0$ , this will mean trace and determinant with respect to the subalgebra  $A_0$ . We denote by  $\Omega_3$  the cone of positivity of  $A$ , that is,  $\Omega_3 = \{x \in A / \lambda_i > 0, i = 1, 2, 3\} = \{x \in A / a_i(x) > 0, i = 1, 2, 3\}$  and by  $\Omega_2$  the cone of  $A_0$ . Every  $x$  in  $\Omega_3$  (resp., in  $\Omega_2$ ) admits a unique square root in  $\Omega_3$  (resp., in  $\Omega_2$ ) and is invertible.

LEMMA 3.1. *If  $y = e_0 + \xi + tc_3$  with  $\xi \in A_{1/2}$  and  $t \in \mathbb{R}$ , then*

$$\det(y) = t - \frac{1}{2} \|\xi\|^2. \tag{3.2}$$

*Proof.* The projection onto  $A_1$  is  $P(c_3)$ , so  $P(c_3)\xi = 0$  and therefore  $0 = \text{tr}(P(c_3)\xi) = (e, P(c_3)\xi) = (c_3, \xi) = (1/2) \text{tr}(\xi)$ . By the same argument  $\text{tr}(\xi^3) = 0$ . Now by Cayley-Hamilton  $\xi^3 + a_2(\xi)\xi - \det(\xi)e = 0$  which implies  $\det(\xi) = 0$ . On the other hand,  $\xi^2 = u + \tau c_3$  with  $u \in A_0$  and  $\tau \in \mathbb{R}$ . We have  $\tau = (c_3, \xi^2) = (\xi c_3, \xi) = (1/2) \|\xi\|^2$  and  $\text{tr}(u) = (1/2) \|\xi\|^2$ . From  $\det(\xi^2) = \tau \det(u)$ , we deduce that  $\det(u) = 0$ . So by Cayley-Hamilton (in  $A_0$ ) we can write  $u^2 = (1/2) \|\xi\|^2 u$ . Now  $\xi^2 = u + \tau c_3 \Rightarrow \xi^4 = u^2 + \tau^2 c_3 = (1/2) \|\xi\|^2 u + (1/4) \|\xi\|^4 c_3 = (1/2) \|\xi\|^2 \xi^2$  and then  $\xi^3 = (1/2) \|\xi\|^2 \xi$ . Therefore,  $u\xi = (1/4) \|\xi\|^2 \xi$ . Then we have

$$\begin{aligned} y &= e_0 + \xi + tc_3, \\ y^2 &= (e_0 + u) + (1 + t)\xi + \left(t^2 + \frac{1}{2} \|\xi\|^2\right) c_3, \\ y^3 &= (e_0 + (2 + t)u) + \left(1 + t + t^2 + \frac{1}{2} \|\xi\|^2\right) \xi + \left(t^2 + t \|\xi\|^2 + \frac{1}{2} \|\xi\|^2\right) c_3, \end{aligned} \tag{3.3}$$

and also  $\text{tr}(y) = 2 + t$ ,  $a_2(y) = (1/2)[\text{tr}(y)^2 - \text{tr}(y^2)] = 1 + 2t - (1/2)\|\xi\|^2$ . The result is a consequence of

$$y^3 - \text{tr}(y)y^2 + a_2(y)y = \left(t - \frac{1}{2}\|\xi\|^2\right)e. \quad (3.4)$$

□

LEMMA 3.2. If  $y = z + \xi + tc_3$  with  $z \in \Omega_2$ ,  $\xi \in A_{1/2}$ , and  $t \in \mathbb{R}$ , then

$$\det(y) = \det(z)[t - (z^{-1}\xi, \xi)]. \quad (3.5)$$

*Proof.* The application

$$\begin{aligned} \rho : A_0 &\longrightarrow \text{End}(A_{1/2}), \\ u &\longrightarrow \rho(u) \text{ defined by } \rho(u)\xi = 2L(u)\xi \end{aligned} \quad (3.6)$$

is a representation of  $A_0$  in the space  $A_{1/2}$  (for more details, see [4]). This means that  $2\rho(uv) = \rho(u)\rho(v) + \rho(v)\rho(u)$  and identically  $P(u)\xi = 0$ . So

$$\begin{aligned} P(u + c_3)y &= P(u + c_3)z + P(u + c_3)\xi + tP(u + c_3)c_3 \\ &= P(u)z + 2L(u)\xi + tc_3. \end{aligned} \quad (3.7)$$

Now if  $u = z^{-1/2}$ , we derive the desired result thanks to (3.2) and the fact that

$$\begin{aligned} \det[P(z^{-1/2} + c_3)y] &= \det(z^{-1/2} + c_3)^2 \det(y) \\ &= \det(z^{-1}) \det(y) = \det(z)^{-1} \det(y). \end{aligned} \quad (3.8)$$

□

COROLLARY 3.3. If  $y = z + \xi + tc_3 \in \Omega_3$ , then

$$\text{tr}(y^{-1}) = \frac{2\det(z) + 2t\text{tr}(z) - \|\xi\|^2}{2\det(z)[t - (z^{-1}\xi, \xi)]}. \quad (3.9)$$

*Proof.* First we have  $y^3 - \text{tr}(y)y^2 + a_2(y)y - \det(y)e = 0$ . Then

$$y^{-1} = \frac{1}{\det(y)}[y^2 - \text{tr}(y)y + a_2(y)e] \quad (3.10)$$

and therefore

$$\begin{aligned} \text{tr}(y^{-1}) &= \frac{1}{\det(y)}[\text{tr}(y^2) - \text{tr}(y)^2 + 3a_2(y)] \\ &= \frac{1}{2\det(y)}[\text{tr}(y)^2 - \text{tr}(y^2)] \\ &= \frac{1}{2\det(y)}[\text{tr}(z)^2 + 2t\text{tr}(z) - \text{tr}(z^2) - \|\xi\|^2] \\ &= \frac{1}{2\det(y)}[2\det(z) + 2t\text{tr}(z) - \|\xi\|^2]. \end{aligned} \quad (3.11)$$

□

#### 4. $K$ -Bessel function

The  $K$ -Bessel function is defined by (see [2, 4])

$$K_{\nu}^{[3]}(x) = \int_{\Omega_3} e^{-\text{tr}(y^{-1})-(x,y)} (\det y)^{\nu-n/3} dy. \quad (4.1)$$

This function is well defined at least for  $x \in \Omega_3$  and any complex  $\nu$ . The convergence of the previous integral is assured by the good behaviour of  $e^{-\text{tr}(y^{-1})-(x,y)}$  near infinity (in  $\Omega_3$ ) and  $\partial\Omega_3$ . We can extend it to the tube  $\Omega_3 \oplus iA$  in the complexification  $A_{\mathbb{C}} = A \oplus iA$  of  $A$  as a holomorphic function in  $x$ . As a function of  $\nu$ , it is an entire function (see [4, Proposition XVI.3.1]). After a change of variables, one can show that

$$K_{\nu}^{[3]}(x) = (\det x)^{-\nu} K_{-\nu}^{[3]}(x). \quad (4.2)$$

Following [2] where it is proved that  $K_{\nu}^{[3]}$  is a solution of a differential system similar to (2.1), we can write

$$K_{\nu}^{[3]}(x) = \sum_{j=1}^4 a_{\nu}^j J_{\nu}^{[3,j]}(-x) + b_{\nu}^j t_3^{-\nu} J_{-\nu}^{[3,j]}(-x). \quad (4.3)$$

According to (4.2), we have  $a_{\nu}^j = b_{-\nu}^j$  for  $j = 1, 2, 3, 4$ . For  $\text{Re}(\nu) < -d$ , the following limit holds (see [4] for more information on  $\Gamma_{\Omega_3}$ , the Gamma function of the cone  $\Omega_3$ ):

$$\lim_{\substack{x \rightarrow 0 \\ x \in \Omega_3}} K_{\nu}^{[3]}(x) = \Gamma_{\Omega_3}(-\nu) = (2\pi)^{3d/2} \Gamma(-\nu) \Gamma\left(-\nu - \frac{d}{2}\right) \Gamma(-\nu - d), \quad (4.4)$$

so

$$a_{\nu}^1 = b_{-\nu}^1 = (2\pi)^{3d/2} \Gamma(-\nu) \Gamma\left(-\nu - \frac{d}{2}\right) \Gamma(-\nu - d) \quad (4.5)$$

according to the behaviour of the solutions  $J_{\nu}^{[3,j]}$ . To determine the other coefficients, we take  $x \neq 0$  on the boundary of  $\Omega$ . So if  $x = x_1 c_1 + x_2 c_2$ , then the integral representation of  $K_{\nu}^{[3]}$  takes the explicit form

$$\begin{aligned} K_{\nu}^{[3]}(x_1 c_1 + x_2 c_2) &= \int_{\Omega_2} \iint_E \exp - \left[ \frac{2 \det(z) + 2t \text{tr}(z) - \|\xi\|^2}{2 \det(z) [t - (z^{-1} \xi, \xi)]} \right] \\ &\quad \times \exp -(x, z) \det(z)^{\nu-d-1} [t - (z^{-1} \xi, \xi)]^{\nu-d-1} dz d\xi dt, \end{aligned} \quad (4.6)$$

where  $E = \{(t, \xi) \in \mathbb{R} \times A_{1/2} / t > (z^{-1} \xi, \xi)\}$  and  $y = z + \xi + tc_3$ . We change  $t$  into  $t + (z^{-1} \xi, \xi)$ . The integral over  $E$  becomes

$$\begin{aligned} I &= \int_0^{+\infty} \int_{A_{1/2}} e^{-1/t} \exp - \left[ \frac{\text{tr}(z)(z^{-1} \xi, \xi) - (1/2) \|\xi\|^2}{t \det(z)} \right] t^{\nu-d-1} d\xi dt \\ &= \int_0^{+\infty} \int_{A_{1/2}} e^{-1/t} e^{-(B \cdot \xi, \xi)} t^{\nu-d-1} d\xi dt, \end{aligned} \quad (4.7)$$

where the operator  $B = \rho(\nu)$  with  $\nu = (\text{tr}(z)/2t\det(z))z^{-1} - (1/2t\det(z))e_0$ . Note that  $\nu \in \Omega_2$  because  $\text{tr}(\nu) = (\text{tr}(z)^2 - 2\det(z))/2t(\det(z)^2) > 0$  and  $\det(\nu) = 1/4t^2(\det(z))^2 > 0$ . Also  $\det B = \det(\nu)^d = 2^{-2d}t^{-2d}\det(z)^{-2d}$ . But

$$\int_{A_{1/2}} e^{-(B \cdot \xi, \xi)} d\xi = \pi^d (\det B)^{-1/2}, \quad (4.8)$$

so

$$I = (2\pi)^d \det(z)^d \int_0^{+\infty} e^{-1/t} t^{\nu-1} dt = (2\pi)^d \Gamma(-\nu) \det(z)^d. \quad (4.9)$$

Now

$$\begin{aligned} K_\nu^{[3]}(x_1 c_1 + x_2 c_2) &= (2\pi)^d \Gamma(-\nu) \int_{\Omega_2} e^{-\text{tr}(z)/\det(z)} \exp -(x, z) \det(z)^{\nu-1} dz \\ &= (2\pi)^d \Gamma(-\nu) \int_{\Omega_2} e^{-\text{tr}(z^{-1})} \exp -(x, z) \det(z)^{\nu+d/2-d/2-1} dz \\ &= (2\pi)^d \Gamma(-\nu) K_{\nu+d/2}^{[2]}(x_1 c_1 + x_2 c_2). \end{aligned} \quad (4.10)$$

**THEOREM 4.1.** *It holds that*

$$K_\nu^{[3]}(t_1, t_2, t_3) = \sum_{j=1}^4 a_\nu^j J_\nu^{[3,j]}(-t_1, t_2, -t_3) + b_\nu^j t_3^{-\nu} J_{-\nu}^{[3,j]}(-t_1, t_2, -t_3) \quad (4.11)$$

with

$$\begin{aligned} a_\nu^1 = b_{-\nu}^1 &= (2\pi)^{3d/2} \Gamma(-\nu) \Gamma\left(-\nu - \frac{d}{2}\right) \Gamma(-\nu - d), \\ a_\nu^2 = b_{-\nu}^2 &= (2\pi)^{3d/2} \Gamma(-\nu) \Gamma\left(-\nu - \frac{d}{2}\right) \Gamma(\nu + d), \\ a_\nu^3 = b_{-\nu}^3 &= (2\pi)^{3d/2} \Gamma(-\nu) \Gamma\left(\nu + \frac{d}{2}\right) \Gamma(\nu), \\ a_\nu^4 = b_{-\nu}^4 &= (2\pi)^{3d/2} \Gamma(-\nu) \Gamma\left(\nu + \frac{d}{2}\right) \Gamma(-\nu). \end{aligned} \quad (4.12)$$

*Proof.* We suppose first that  $\text{Re}(\nu) < -d$ . So the left-hand side of the formula (4.3) is nothing but  $(2\pi)^d \Gamma(-\nu) K_{\nu+d/2}^{[2]}(x_1 c_1 + x_2 c_2)$  according to (4.10). We use then the expansion established in the case of rank-2 Jordan algebra (see [3]) and the reduction formulas (2.8) to (2.11) for the right-hand side of (4.3).  $\square$

## 5. Conclusion

In light of the above calculations, we think that the complete answer to Herz's conjecture (higher ranks) can be fulfilled in two steps. First a recurrence formula between  $K$ -Bessel functions of rank  $r$  and  $r+1$  like formula (4.3). This is quite easy because the lemmas above can be written in higher ranks using some Jordan algebra identities. But what seems to be very difficult is the second step: the complete and explicit resolution

of the Bessel-Muirhead system. Here the recurrence does not work because the series method yields a very complicated recurrence system on the coefficients.

### Acknowledgment

The author is grateful to the referees for their relevant and helpful remarks.

### References

- [1] T. E. Bengtson, *Bessel functions on  $P_n$* , Pacific J. Math. **108** (1983), no. 1, 19–29.
- [2] H. Dib, *Fonctions de Bessel sur une algèbre de Jordan [Bessel functions on a Jordan algebra]*, J. Math. Pures Appl. (9) **69** (1990), no. 4, 403–448 (French).
- [3] ———, *K-Bessel functions in two variables*, Int. J. Math. Math. Sci. **2003** (2003), no. 14, 909–916.
- [4] J. Faraut and A. Korányi, *Analysis on Symmetric Cones*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 1994.
- [5] J. Faraut and G. Travaglini, *Bessel functions associated with representations of formally real Jordan algebras*, J. Funct. Anal. **71** (1987), no. 1, 123–141.
- [6] C. S. Herz, *Bessel functions of matrix argument*, Ann. of Math. (2) **61** (1955), no. 3, 474–523.
- [7] A. T. James, *A generating function for averages over the orthogonal group*, Proc. Roy. Soc. London Ser. A **229** (1955), 367–375.
- [8] A. Korányi, *Transformation properties of the generalized Muirhead operators*, Colloq. Math. **60/61** (1990), no. 2, 665–669.
- [9] N. H. Mahmoud, *Bessel systems for Jordan algebras of rank 2 and 3*, J. Math. Anal. Appl. **234** (1999), no. 2, 372–390.
- [10] R. J. Muirhead, *Systems of partial differential equations for hypergeometric functions of matrix argument*, Ann. of Math. Statistics **41** (1970), no. 3, 991–1001.
- [11] ———, *Aspects of Multivariate Statistical Theory*, Wiley Series in Probability and Mathematical Statistics, John Wiley & Sons, New York, 1982.

Hacen Dib: Department of Mathematics, Faculty of Sciences, University of Tlemcen, BP 119, Tlemcen 13000, Algeria

*E-mail address:* h\_dib@mail.univ-tlemcen.dz

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)